

#### U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:	Date of Issuance:

34704-1101

6/30/16

\	(	١(	ГΤ	C	F.	$\mathbf{O}$	F	P	ES	Ľ	T	C	П	D	Ē٠	

\_X\_Registration \_\_\_Reregistration (under FIFRA, as amended) Term of Issuance:
Conditional

Name of Pesticide Product:

**Equity VIP** 

Name and Address of Registrant (include ZIP Code):

Lisa Nichols Loveland Products Inc. P.O. Box 1286 Greeley, CO 80632-1286

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/registration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:	Date:
VE	6/30/16
Venus Eagle, Product Manager 01 Invertebrate-Vertebrate Branch 3, Registration Division (7505P)	

EPA Form 8570-6

Page 2 of 2 EPA Reg. No. 34704-1101 Decision No. 517220

- 2. You are required to comply with the data requirements described in the DCI Order identified below:
  - a. Thiamethoxam GDCI-060106-1309
  - b. Fludioxonil GDCI-071503-1243
  - c. Thibendazole GDCI-060101-1453

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <a href="http://www.epa.gov/oppsrrd1/contacts\_prd.htm">http://www.epa.gov/oppsrrd1/contacts\_prd.htm</a>

- 3. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, "EPA Reg. No. 34704-1101."
- 4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

• Basic CSF dated 05/05/2016

If you have any questions, please contact Dee Colby by phone at 703-347-8657, or via email at colby.deanna@epa.gov.

Enclosure: Stamped label







#### INSECTICIDE WITH FUNGICIDES

A seed treatment product for protection against damage from listed: insects, seed-borne diseases, and seedling diseases on soybean

#### **ACTIVE INGREDIENTS:**

Thiamethoxam <sup>1</sup>		22.40%
Mefenoxam <sup>2</sup>		
Thiabendazole <sup>3</sup>		
Fludioxonil <sup>4</sup>		
Sedaxane <sup>5</sup>		1.12%
OTHER INGREDIENTS:		<u>69.77%</u>
	TOTAL:	100.00%

<sup>1</sup>CAS No. 153719-23-4

<sup>2</sup>CAS No. 70630-17-0 and CAS No. 69516-34-3

<sup>3</sup>CAS No. 148-79-8 <sup>4</sup>CAS No. 131341-86-1 <sup>5</sup>CAS No. 874967-67-6

One gallon of EQUITY VIP contains 2.16 lb thiamethoxam, 0.32 lb mefenoxam,—

-0.22 lb

thiabendazole, 0.11 lb fludioxonil and 0.11 lb sedaxane.

## CAUTION

See additional precautionary statements and directions for use in booklet.

EPA REG. NO. 34704-XXXX EPA EST. XXXXX

Net Contents: 15 gallons (56.78 liters)

ACCEPTED

Jun 30, 2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under

EPA Reg. No. 34704-1101

SCPSR-CPS-1559A-L1 1015

4060020

Product ID: 57388

Equity is a registered trademark of Loveland Products, Inc.

FIRST AID					
If swallowed • Call a poison control center or doctor immediately for treatment advice.					
Have person sip a glass of water if able to swallow.					
<ul> <li>Do not induce vomiting unless told to by a poison control center or doctor.</li> </ul>					
Do not give anything to an unconscious person.					
Have the product container or label with you when calling a poison control center or doctor or going for treatment.					
FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565					

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling, and before eating, drinking, chewing gum, using tobacco, or using the toilet.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

#### Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants.
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- · Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **ENGINEERING CONTROL STATEMENTS**

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **USER SAFETY RECOMMENDATIONS**

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to wildlife, freshwater and estuarine/marine fish and highly toxic to aquatic invertebrates. Runoff may be hazardous to aquatic organisms in neighboring areas. Exposed treated seed may be hazardous to wildlife. Cover seed spilled during loading. Do not contaminate water when disposing of equipment washwater or rinsate.

#### **Pollinator Precautions**

Thiamethoxam is highly toxic to bees exposed to direct treatment, and effects may be possible as a result of exposure to translocated residues in blooming crops.

#### **Groundwater Advisory**

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of label use. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### **Physical and Chemical Hazards**

Do not use, pour, spill or store near heat or open flame. Do not store near or use with oxidizing agents.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treatment facilities. Use is also permitted as an end-use seed treatment on agricultural establishments at planting, or immediately before planting, as specified in the Specific Crop Use Directions. This product is to be used in liquid or slurry treaters only.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils
- Shoes plus socks

### FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT AND/OR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat a quantity of seed using equipment similar to that planned for treating the total seed lot. Prior to treatment, conduct germination tests on a portion of seed before committing the total seed lot to a selected seed treatment.

Due to seed quality, crop or variety sensitivity, and seed storage conditions beyond the control of Loveland Products, Inc., no claims are made to guarantee the germination of seed or propagating material for all crop seed when treated with EQUITY VIP.

#### **USE INFORMATION**

EQUITY VIP is a seed treatment product containing the active ingredients thiamethoxam (insecticide) and fludioxonil, thiabendazole, mefenoxam and sedaxane (fungicides). EQUITY VIP protects against damage from listed early-season insects, soil-borne and seed-borne diseases of soybeans.

Thiamethoxam is a systemic seed treatment insecticide belonging to the neonicotinoid class of chemistry. Thiamethoxam protects against listed chewing and sucking insects through contact and ingestion.

Mefenoxam fungicide is active against *Pythium*, *Phytophthora* and systemic downy mildew.

Thiabendazole fungicide is active for protection against phomopsis spp. to improve germination and early season seedling health. Thiabendazole also suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

Fludioxonil fungicide is active against *Fusarium* and *Rhizoctonia*, and suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

Sedaxane fungicide is active against seed decay, seedling blight and damping-off caused by *Rhizoctonia* species.

#### RESISTANCE MANAGEMENT

GROUP 4A INSECTICIDE

EQUITY VIP contains thiamethoxam, a Group 4A insecticide. Thiamethoxam is a systemic insecticide belonging to the neonicotinoid class of chemistry which includes nicotinic acetylcholine receptor (nAChR) agonists.

Insect populations may contain individuals naturally resistant to Group 4A insecticides and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, use sound resistance management strategies established for the crop and use area.

Base seed treatment on an integrated pest management program that includes field sanitation, historical information related to pesticide use, careful selection of pest-tolerant crop varieties, scouting, and management practices which optimize populations of natural enemies of insect pests such as within-field refugia (untreated areas). Sound management programs also consider cultural and biological control practices.

#### In order to maintain susceptibility to this class of chemistry:

- Use products at their full, specified doses.
- Use appropriate, well-maintained equipment. Use specified water volumes and apply at optimal temperatures in order to obtain optimal treatment.
- When rate ranges are given, use the higher rate within the listed rate range when insect pressure is expected to be high.
- Avoid using a single active ingredient or mode of action (same insecticide group) exclusively for season long control of
  insect species with more than one generation per crop season.
- For insect species with successive or overlapping generations, use a treatment window approach. A treatment window is a
  period of time defined by the stage of crop development and the biology of the pests of concern. Within the treatment
  window, depending on the length of residual activity, single or consecutive applications may be made using seed, infurrow, or foliar treatments unless otherwise excluded by product labels. Do not exceed the maximum amount of this
  insecticide's mode of action allowed per growing season.
- Following a treatment window of this insecticide's mode of action, rotate to a treatment window of effective products with a different mode of action before making additional applications of this insecticide.

If resistance to this product develops in your area, this product or other products with a similar mode of action may not provide adequate control. If poor performance cannot be attributed to improper application or weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for the crop and use area.

GROUP 1 4 7 12 FUNGICIDES

EQUITY VIP contains mefenoxam, a Group 4 fungicide; thiabendazole, a Group 1 fungicide, fludioxonil, a Group 12 fungicide; and sedaxane, a Group 7 fungicide. Thiabendazole belongs to the methyl-benzimidazole carbamate class of chemistry which disrupts β-tubulin assembly in mitosis. Mefenoxam belongs to the phenylamide class of chemistry which

interferes with fungal RNA synthesis. Sedaxane is a succinate dehydrogenase inhibitor (SDHI) and belongs to the carboxamide class of chemistry which disrupts cellular respiration and energy generation. Fludioxonil belongs to the phenylpyrrole class of chemistry which interferes with a MAP/histidine protein kinase in osmotic signal transduction.

Fungal populations may contain individuals naturally resistant to Group 1, 4, 7, and 12 fungicides, and if used repeatedly in the same fields, then resistant members may eventually dominate the population. Because resistance development cannot be predicted, the use of this product should conform to sound resistance management strategies such as alternation with fungicides with a different mode of action and/or tank mixes established for the crop and use area.

Use should be based on an IPM program that includes field sanitation, scouting, historical information related to pesticide use, and crop rotation. The IPM program should also consider cultural, biological, and other chemical control practices.

Loveland Products, Inc. encourages responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

#### For additional information on Insecticide or Fungicide Resistance Management:

- Contact Loveland Products, Inc. representatives at 1-888-574-2878
- Contact your local Cooperative Extension Service specialist, pest control advisor, or certified crop advisor
- Visit the Visit the Insecticide Resistance Action Committee (IRAC) on the web at: http://www.irac-online.org or the Fungicide Resistance Action Committee (FRAC) on the web at: http://www.frac.info

#### **CROP ROTATIONAL RESTRICTIONS**

In the event of a crop failure or harvest of a crop grown from EQUITY VIP treated seed, the field may be replanted according to the following schedule:

Immediate Plantback	Minimum 30-Day Plantback Interval	Minimum 120-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye,	Alfalfa	All Other Crops
Triticale, and Wheat	Brassica (Cole) Leafy Vegetables	
Peas, Dried Shelled: Chickpea (Garbanzo Bean), Field Pea, Lentil, and Pigeon Pea	Cereal Grains (buckwheat, corn, pearl millet, proso, popcorn, rice (dry- seeded), sorghum, teosinite, wild rice	
Soybean	Cotton	
	Cucurbit Vegetables	
	Fruiting Vegetables	
	Globe Artichoke	
	Leafy Vegetables	
	Legume Vegetables (Succulent or Dried)	
	Mint: Peppermint and Spearmint	
	Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower	
	Peanut	
	Root Vegetables	
	Strawberry	
	Sunflower	
	Tobacco	

For any other crop, the minimum plant back interval is 120 days from the date the EQUITY VIP treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

#### MIXING PROCEDURES

**Important:** Always re-circulate EQUITY VIP thoroughly before using.

Consult the manufacturer of the application equipment you plan to use for suitability for this application and for instructions on operation and calibration of the equipment. Follow the manufacturer application instructions for the seed treatment equipment being used.

Use an EPA approved dye/colorant that imparts an unnatural color to the seed as required in 40CFR 153.155(c).

Apply EQUITY VIP as a water-based slurry utilizing standard slurry or direct inject seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the specified amount of EQUITY VIP into the required amount of water or liquid inoculant for the slurry treater and dilution rate to be used. For direct product inject systems, add the required amount of water or liquid inoculant during application.

Certain crops require addition of inoculants when the seed is treated or planted. EQUITY VIP is compatible with several liquid inoculant products. Consult the maker of the inoculant product and a Loveland Products, Inc. representative for directions before applying EQUITY VIP with inoculants.

The total application volume must be sufficient to provide desired level of coverage. Dilution is typically done with water or liquid inoculants. The minimum slurry volume to achieve minimum adequate coverage is 4.0 fluid ounces per 100 pounds of seed assuming an average seed size of 3,000 seeds per pound (slurry volume will vary depending on seed size). More diluent may be required to obtain complete coverage.

Continuous agitation or mixing of the slurry mixture is necessary to prevent settling out of the solution.

Allow seed to dry before bagging or reprocessing directly after treating operation.

Follow planter manufacturer specifications for use of talc or other hopper box additives at planting. Seed must be completely dry before adding to planter.

#### SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing treated seeds shall be labeled with the following statements:

- This seed has been treated with thiamethoxam insecticide and fludioxonil, mefenoxam, thiabendazole and sedaxane fungicides.
- Do not use for feed, food, or oil purposes.

In addition, the U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with EQUITY VIP:

Ground Water Advisory:

Thiamethoxam has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Mefenoxam is known to leach through soil into ground water under certain conditions as a result of agricultural use. This chemical may leach into the ground water if used in areas where soils are permeable, particularly where the water table is shallow.

- Pollinator Precautions:
  - Thiamethoxam is highly toxic to bees, and effects are possible as a result of exposure to translocated residues in blooming crops.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in the ethanol by-products that are used in agronomic practice.
- Do not allow children, pets, or livestock to have access to treated seed.
- Store away from feeds and foodstuffs.
- Wear long-sleeved shirt, long pants and chemical resistant gloves when handling treated seed.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.
- Treated seed must be planted into the soil at a depth greater than 1 inch.
- Dispose of all excess treated seed. Leftover treated seed may be doublesown around the headland or buried away from water sources in accordance with local requirements.
- Do not contaminate water bodies when disposing of planting equipment wash waters.
- Dispose of seed packaging in accordance with local requirements.
- Do not graze or feed livestock on treated areas for 45 days after planting seed treated with EQUITY VIP.
- In the event of a crop failure or harvest of a crop grown from EQUITY VIP treated seed, the field may be replanted according to the following schedule.

Immediate Plantback	Minimum 30-Day Plantback Interval	Minimum 120-Day Plantback Interval
Cereal Grains: Barley, Oats, Rye,	Alfalfa	All Other Crops
Triticale, and Wheat	Brassica (Cole) Leafy Vegetables	
Peas, Dried Shelled: Chickpea (Garbanzo Bean), Field Pea, Lentil, and Pigeon Pea	Cereal Grains (buckwheat, corn, pearl millet, proso, popcorn, rice (dry- seeded), sorghum, teosinite, wild rice	
Soybean	Cotton	
	Cucurbit Vegetables	
	Fruiting Vegetables	
	Globe Artichoke	
	Leafy Vegetables	
	Legume Vegetables (Succulent or Dried)	
	Mint: Peppermint and Spearmint	
	Oilseeds: Borage, Crambe, Flax Seed, Mustard Seed, and Safflower	
	Peanut	
	Root Vegetables	
	Strawberry	
	Sunflower	
	Tobacco	

• For any other crop, the minimum plant back interval is 120 days from the date the EQUITY VIP treated seed was planted. A cover crop other than the crops listed above that is planted for erosion control or soil improvement may be planted sooner than the 120 day interval; however, the crop may not be grazed or harvested for food or feed.

- Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.8 grams ai/A) per acre, 0.004 lb fludioxonil (2.0 grams ai/A) per acre and 0.013 lb mefenoxam (5.7 grams ai/A) per acre per calendar year as a seed treatment application.
- This seed has been treated with 0.0756 mg ai thiamethoxam, 0.0113 mg mefenoxam, and 0.0038 mg fludioxonil per seed.
- Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with EQUITY VIP.
- Do not use more than 0.266 lb ai/A of thiamethoxam per calendar year, including all thiamethoxam products and application methods (for example, seed treatment, in-furrow, and foliar).

#### **CROP USE DIRECTIONS**

When applied according to the **EQUITY VIP RATE TABLE**, EQUITY VIP provides early season protection against injury by aphids, bean leaf beetle, grape colaspis, leaf miners, leaf hoppers, Mexican bean beetle, seed corn maggot, three-cornered alfalfa hopper, thrips, white grubs, and wireworm.

EQUITY VIP provides protection against damping-off and seed borne rots due to *Pythium, Phytophthora, Fusarium, Rhizoctonia* species and early season *Phytophthora* root rot. EQUITY VIP also suppresses seed-borne *Sclerotinia* and *Phomopsis* species.

#### **EQUITY VIP Rate Table<sup>1</sup>**

Crop	Rate of EQUITY VIP					
Soybean, including soybean, vegetable	fl oz per 100 lb seed or	grams ai per 100 kg seed	mg ai per seed			
<b>,</b> , <b>,,</b>	fl oz per 140,000 seeds		<b>F</b>			
	2.96 fl oz	Thiamethoxam 50 g	Total = 0.102			
	or	Mefenoxam 7.5 g				
	1.38 fl oz per 140,000 seeds	Thiabendazole 5.0 g				
		Fludioxonil 2.5 g				
		Sedaxane 2.5 g				

<sup>&</sup>lt;sup>1</sup>The mg ai per seed and fl oz EQUITY VIP per 100 lb seed rates are based on 3,000 seeds per pound.

#### **USE RESTRICTIONS**

- Do not allow children, pets, or livestock to have access to treated seed.
- Do not graze or feed livestock on treated areas for 45 days after planting seed treated with EQUITY VIP.
- Do not use at a rate that will result in more than 0.083 lb thiamethoxam (37.8 grams ai/A) per acre, 0.004 lb fludioxonil (2.0 grams ai/A) per acre and 0.013 lb mefenoxam (5.7 grams ai/A) per acre per calendar year as a seed treatment application.
- This seed has been treated with 0.0756 mg ai thiamethoxam, 0.0113 mg mefenoxam, and 0.0038 mg fludioxonil per seed.
- Do not use more than 0.266 lb ai/A of thiamethoxam per calendar year, including all thiamethoxam products and application methods (for example, seed treatment, in-furrow, and foliar).
- Do not apply a neonicotinoid insecticide within 45 days of planting seed treated with EQUITY VIP.

#### STORED GRAIN PROTECTION

When treated according to the directions for post-planting protection against listed pests, EQUITY VIP will also provide protection during post treatment storage of the soybean seed against damage from the following stored grain insects: Indian Meal Moth (*Plodia interpunctella*), Rice Weevil (*Sitophilus oryza*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*).

If the soybean seed to be treated has existing infestations of stored grain insects, fumigate the seed with a registered product approved for such use prior to treating with EQUITY VIP and bagging.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, or disposal, or cleaning of equipment.

#### **Pesticide Storage**

Store in a cool, dry place. Storage for extended periods above 90°F is not recommended.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

#### **Pesticide Disposal**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

#### Container Handling [less than or equal to 5 gallons]

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

#### Container Handling [greater than 5 gallons]

**Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**BEFORE BUYING OR USING THIS PRODUCT**, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. The buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL

AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. Box 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

For non-emergency (e.g., current product information), call Loveland Products, Inc. 1-888-574-2878.

Equity is a registered trademark of Loveland Products, Inc. ©2016 Loveland Products, Inc.

Formulated for: Loveland Products, Inc. P.O. Box 1286 Greeley, Colorado 80632-1286