



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

34704-1157

Date of Issuance:

8/3/20

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

LPI.A011

Name and Address of Registrant (include ZIP Code):

Robert Avalos
 Manager of Registrations
 Loveland Products, Inc.
 P.O. Box 1286
 Greeley, CO 80632-1286

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Continued on page 2

Signature of Approving Official:

Mindy Ondish, Product Manager 23
 Herbicide Branch, Registration Division (7505P)

Date:

8/3/20

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

The alternate brand name, "Zoran 26.2 EC" has been added to the product record.

Please also note that the record for this product currently contains the following CSF:

- Basic CSF dated 02/25/2020

If you have any questions, please contact Julia Kerr by phone at 703-347-0386, or via email at kerr.julia@epa.gov.

Enclosure

{BOOKLET FRONT PANEL LANGUAGE}

FLUROXYPYR

GROUP

4

HERBICIDE

LPI.A011^[TM]**[Alternate Brand Name: Zoran 26.2 EC]**

SELECTIVE POSTEMERGENCE CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS AND VOLUNTEER POTATOES IN BARLEY, OATS, WHEAT, FIELD CORN, SWEET CORN, GRAIN SORGHUM, FALLOW CROPLAND, AND ON-FARM NON-CROPLAND.

MAY ALSO BE USED FOR SELECTIVE POSTEMERGENCE CONTROL OF ANNUAL AND PERENNIAL BROADLEAF WEEDS AND WOODY BRUSH IN: RANGELAND AND PERMANENT PASTURES, NON-CROPLAND AREAS INCLUDING INDUSTRIAL SITES, NON-IRRIGATION DITCH BANKS, AND RIGHTS-OF-WAY (ELECTRICAL POWER LINES, COMMUNICATION LINES, PIPELINES, ROADSIDES AND RAILROADS INCLUDING GRAZED AREAS WITHIN THESE SITES) AND PINE PLANTATIONS.

NOT FOR SALE, SALE INTO, DISTRIBUTION AND/OR USE IN NASSAU AND SUFFOLK COUNTIES OF NEW YORK STATE

Active Ingredient:	(% by weight)
Fluroxypyr: 1-methylheptyl ester: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy)	
acetic acid, 1-methylheptyl ester	26.2%
Other Ingredients	73.8%
Total	100.0%
Contains 1.5 pounds fluroxypyr acid per gallon (18.2%)	
Contains Petroleum Distillates	

**KEEP OUT OF REACH OF CHILDREN
WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside label booklet for First Aid, Precautionary Statements and Directions for Use.


EPA Reg. No.: 34704-1157

EPA Est. No.:

Net Weight:

[Label Print Date Code]

MANUFACTURED FOR:
LOVELAND PRODUCTS, INC.
P.O. BOX 1286
GREELEY, COLORADO 80632-1286



{ LANGUAGE INSIDE BOOKLET }

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. • Do not give any liquid to the person
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Note to Physician: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.	

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate or Viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agriculture pesticides 40 CFR 170.607(d-e), the handler PPE requirements may be reduced or modified as specified in the WPS

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

Physical Chemical Hazards:

Do not mix or allow to come into contact with an oxidizing agent. A hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water are:

- Coveralls
- Chemical-resistant gloves made of barrier laminate or Viton \geq 14 mils
- Shoes plus socks
- Protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. **Do not enter treated areas until sprays have dried.**

WEED RESISTANCE MANAGEMENT

For resistance management, **LPI.A011** is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to **LPI.A011** and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of **LPI.A011** or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures from a different group if such use is permitted; where information on resistance in target weeds species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance contact Loveland Products, Inc. retailer, representative or call 1-888-574-2878. You can also contact your pesticide distributor or university extension specialist to report resistance.

PRODUCT INFORMATION

Crop: This emulsified concentrate product provides selective postemergence control of annual and perennial broadleaf weeds and volunteer potatoes in on-farm non-cropland, fallow cropland, wheat, barley, or oats not under seeded with a legume, field corn, sweet corn and grain sorghum.

Non-Crop: This product is a selective postemergence product for control of annual and perennial broadleaf weeds and woody brush in rangeland and permanent pastures, pine plantations, and non-crop areas (including grazed areas within these sites) industrial sites, non-irrigation ditch banks, and rights-of-way (electrical power lines,

communication lines, pipelines, roadsides and railroads).

Restrictions

This product is persistent and may be present in plant materials for over 30 days after application. Do not use treated plant material or manure from animals that have grazed or consumed forage from treated areas for compost, mulch, or mushroom spawn until 30 days after application.

Animals that have been fed fluroxypyr treated forage must be fed forage free of fluroxypyr for at least 3 days before they are moved off the treated property.

AVOIDING DRIFT AND RUN-OFF TO SURFACE WATER OR ADJACENT LAND

To minimize off-site exposure and potential effects on non-target plants and aquatic organisms, this product should be used strictly in accordance with the run-off and drift precautions below. The applicator is responsible for all spray drift produced during application and care should be taken to minimize off-target movement of spray during application. Using a coarser spray category nozzle set-up is the most desirable method for reducing spray drift, however, a drift control agent suitable for agricultural use can be used. If drift control agents are used, be sure to follow all applicable precautions and use directions on the manufacturer's label.

Avoiding Runoff:

This product may have a potential to run-off to surface water or adjacent land under certain conditions. To minimize water run-off, use of vegetation filter strips or treatment setbacks along rivers, creeks, streams, wetlands, etc., or on the downhill side of treated areas where run-off could occur is recommended.

AVOIDING INJURY TO NON-TARGET PLANTS

Aerial Application:

Restrictions:

- In non-cropland areas (including rights-of-way), this product may be applied aerially only by helicopter.
- Do not apply this product to non-cropland areas using fixed-wing aircraft.
- This product may be applied to rangeland, permanent pastures and pine plantations using either fixed wing aircraft or helicopter equipment.
- Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Rangeland, Permanent Pastures and Pine Plantations:

To minimize spray drift, apply this product in a total spray volume of 3 or more gallons per acre. Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high potential for temperature inversion. Spray drift from aerial application can be minimized by applying a coarse spray as per USDA-ARS/PAASS at spray boom pressure no greater than 30 psi; by using straight-stream nozzles directed straight back; and by using a spray boom no longer than 75% the wing span or 90% of rotor width for the aircraft. For fixed wing aircraft, maximum speed during application is limited to 140 mph and application height above the vegetation canopy should not exceed 10 feet. Spray pattern and droplet size distribution can be evaluated by applying sprays containing a water-soluble dye marker or appropriate drift control agents over a paper tape (adding machine tape). Mechanical flagging devices may also be used.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

- **IMPORTANCE OF DROPLET SIZE**

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide

target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- **Volume** - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- **Pressure** - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- **Spray Nozzle** - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- **Adjust Nozzles** - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

- **BOOM HEIGHT - Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

- **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

- **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

- **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

- **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

- **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

- **Boom-less Ground Applications:**

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

- **Handheld Technology Applications:**

Take precautions to minimize spray drift.

MIXING INSTRUCTIONS

Note: When adding ingredients to the mixture, allow time for each ingredient to be thoroughly mixed before adding the next. Be sure to agitate spray mixture before use if allowed to stand after mixing.

1. Fill spray tank with water equal to 1/2 to 3/4 of the required spray volume and start agitation.
2. Add the required labeled amount of this product.
3. Add any surfactants, adjuvants or drift control agents according to the respective manufacturer's instructions.

Maintain agitation during final filling of the spray tank with water and maintain sufficient agitation during application to ensure uniformity of the spray mixture.

Tank Mixing:

This product may be tank mixed with other products at labeled rates as long as tank mixing with products containing fluroxypyr is not prohibited by the label(s) of the tank mix partner products and the tank mix partner products are labeled for the timing and method of application for the use site to be treated. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Tank Mixing Precautions:

- Be sure to follow all applicable use directions, precautions, and limitations on the respective product labels.
- When using injection equipment, note that 2,4-D amine concentrates are not compatible with undiluted **LPI.A011** and cannot be mixed together in the same supply tank. However, 2,4-D ester is compatible with this product when using injection equipment.
- Prior to final use, perform a (jar) test to verify the compatibility of tank mix partner products (see instructions below).

Tank Mixing Restrictions:

- Do not exceed labeled application rates.
- Do not tank mix with other pesticide products that contain the same active ingredient as this product unless the label of either mix partner specifies the maximum dosages that may be applied.
- If packaged in water soluble packaging, do not tank mix with products containing boron or mix in equipment previously used to apply a mixture containing boron unless the tank and spray has been adequately cleaned. (Refer to the Sprayer Cleanup section).

Tank Mix Compatibility Testing (Jar Test):

The following jar test is recommended prior to tank mixing to ensure the compatibility of this product with other tank mix partner products:

1. Mix the desired tank mix ingredients in their relative proportions in a clear glass quart jar with lid.
2. Invert the jar containing the mixture several times and observe the mixture for approximately 1/2 hour.
3. If the mixture balls-up, forms flakes, sludges, gels, oily films or layers, or other precipitates, it is not compatible and the tank mix combinations should not be used.

Tank Mixing Instructions:

1. Fill spray tank with water to 1/4 to 1/3 of the required spray volume.
2. Start agitation and maintain agitation continuously during mixing, final filling and while applying.
3. Add different formulation types in the following order being sure to allow sufficient time for each product to completely mix and dispersion after addition (**Note:** This product is an emulsifiable concentrate (EC) formulation):
 - a. Dry flowables
 - b. Wettable powders
 - c. Aqueous suspensions
 - d. Flowables or liquids
4. Maintaining agitation, fill spray tank to 3/4 of total spray volume and then add this product, other

emulsifiable concentrates, and any solutions.

5. Finish filling the spray tank.
6. While spraying, the tank mix ingredients may settle out of suspension if agitation is stopped before the spray tank is empty. The settled materials must be resuspended before any spraying is resumed and a sparger agitator works particularly well in this situation. Note that settled material may be more difficult to resuspend than when originally mixed.

Application Rates:

In general, the application rates at the lower end of the labeled rate range will be efficacious when applied to susceptible weed species with young, succulent growth. Use the higher rates within the rate range when applying to less sensitive species, perennials, and under conditions where control is more difficult (e.g., when plants are stressed due to drought or extreme temperatures, in dense weed stands and/or the weeds are larger). Higher labeled rates will also be needed to control or suppress weeds in areas where competition from crops is not present (e.g., fallow land).

Sprayer Cleanup:

To avoid injury to desirable plants, before applying other chemicals with the equipment used to apply this product, all equipment must be thoroughly cleaned.

1. After applying this product, flush and rinse application equipment with water thoroughly, disposing of the water according to the disposal instructions in this label. All rinse water must be disposed of in compliance with local, state and federal guidelines.
2. Hose down the interior surfaces of the tank, flushing the tank, hoses, boom and nozzles with clean water for 10 minutes.
3. Fill the tank with water and recirculate for 15 minutes.
4. Spray part of the mixture through the hoses, boom and nozzles and drain the tank.
5. Remove the nozzles and screens and clean separately.
6. If the spray equipment will be used on crops other than those labeled for this product, repeat steps 1 and 2 and thoroughly wash the outside of spray tank and the boom.

Management of Kochia Biotypes:

Research indicates many biotypes of kochia may occur within a single field and while kochia biotypes can vary in their susceptibility to this product, in general all biotypes will be suppressed or controlled at the labeled rate of 2/3 pint per acre. Application of this product at rates lower than 2/3 pint per acre can result in a shift to more tolerant biotypes within a field.

Best Practices for Resistance Management:

Extensive populations of dicamba-tolerant kochia have been identified in certain small grain and corn production regions (such as Chouteau, Fergus, Liberty, Toole, and Treasure counties in the state of Montana). For optimal control of dicamba-tolerant kochia in these counties, apply this product at the labeled rate of 2/3 pint per acre.

To minimize selection pressure and preserve the utility of this product for control of dicamba-tolerant kochia biotypes, this product should be rotated with products that do not contain dicamba.

Application Timing:

Only weeds that have emerged at the time of application will be controlled so be sure to apply to actively growing weeds. Weed control may be reduced and the risk of crop injury (at all stages of growth) may increase if extreme growing conditions (such as drought or near-freezing temperatures) occur prior to, at, or following application. Control may be decreased if target plant foliage is wet at the time of application. Applications of this product are rain-fast within 1 hour after application.

Effect of Temperature on Herbicidal Activity:

The herbicidal activity of this product is influenced by weather conditions. Optimum herbicidal activity requires active plant growth and temperature between 55°F to 75°F. Reduced efficacy will occur when temperatures are below 45°F

or above 85°F. Weed control and crop tolerance may be reduced if frost occurs before or shortly after application (3 days).

Spray Coverage:

Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. For best results (and to minimize spray drift), apply in a spray volume of 5 gallons or more per acre by ground and 3 or more gallons of total spray volume per acre by air. Spray volume should be increased as weed density and vegetative canopy increase in order to obtain equivalent weed control; however, do not exceed 40 gallons per acre total spray volume. Rather than increasing boom pressure, decreased spraying speed or larger nozzle tips should be used to increase spray volume.

Use only nozzle types and spray equipment designed for herbicide application. To reduce spray drift, be sure to follow the precautions in the section titled "Avoiding Injury to Non-Target Plants".

CROP USES AND ON FARM NON-CROPLAND USES

- Volunteer potatoes in Barley, Oats and Wheat
- Field corn
- Sweet corn
- Grain sorghum
- Fallow cropland
- On-farm non-cropland

Crop Use Restrictions:

- See individual crop use directions for additional restrictions including maximum application rates.
- When applying this product, do not contaminate water used for domestic purposes or irrigation ditches.
- Do not allow spray drift to come in contact with or apply this product directly to susceptible broadleaf plants or broadleaf crops, including but not limited to the following: alfalfa, canola, cotton, edible beans, grapes, lentils, lettuce, mustard, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tobacco or tomatoes.
- Do not apply this product through any type of irrigation system (i.e., chemigation).
- If replanting is required within 120 days after application plant only crops listed on this label.

APPLICATION INFORMATION

Broadleaf Weeds Controlled or Suppressed for Crop Uses and On Farm Non-Cropland Uses		
Bedstraw (cleavers)	Grape spp.	Mustard spp. [†]
Bindweed, field [†]	Hemp dogbane	Nightshade spp. [†]
Bindweed, hedge	Horsetail, field [†]	Pennycress, field [†]
Buckwheat, wild [†]	Horseweed (marestail) [†]	Potato, volunteer [†]
Canola, volunteer [†]	Jimsonweed	Prickly lettuce
Chickweed	Knotweed [†]	Puncturevine
Clover, white	Kochia ¹	Purslane, common
Cocklebur	Mallow, common [†]	Ragweed, common
Coffeeweed	Mallow, Venice	Sunflower
Devilsclaw [†]	Marshelder [†]	Thistle, Russian [†]
Flax, volunteer	Morning glory	Velvetleaf

[†] **Indicates Suppression Only** – Suppression is a reduction in weed competition (i.e., a reduction in population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

¹ Includes biotypes that are herbicide resistant or tolerant.

Adjuvants

To improve weed control, a high-quality adjuvant labeled for use on growing crops may be used. An adjuvant can optimize herbicidal activity when applications are made at lower carrier volumes, under conditions of cool temperature, low relative humidity or drought, or to small, heavily pubescent kochia.

Spot Treatments for Crop Uses and On Farm Non-Cropland Uses

Only apply using a calibrated boom sprayer or with a hand sprayer using the following directions:

When using hand-held sprayers for spot applications, be sure to uniformly apply a rate equivalent to a broadcast application. Application rates in the table below are based on an area of 1,000 square feet.

Mix the amount of this herbicide (fluid ounces or ml) corresponding to the desired broadcast rate in one or more gallons of spray. To calculate the amount of this herbicide required for larger areas, multiply the table value (fluid ounces or ml) by the area to be treated in “thousands” of square feet. An area of 1,000 square feet is approximately 10.5 x 10.5 yards (strides) in size.

For example: If the area to be treated is 3,500 sq ft, multiply the table value by 3.5 (calc. 3,500 ÷ 1,000 = 3.5).

Broadcast Rate Conversions for Spot Treatments for Crop Uses and On Farm Non-Cropland Uses	
Broadcast Rate (Pints per Acre)	LPI.A011 per Gallon (Fluid Ounces)
2/3	0.25 (7.25 ml)
1	0.37 (11 ml)
1 – 1/3	0.50 (14.5 ml)

BARLEY, OATS AND WHEAT

Restrictions:

- Do not harvest treated forage or allow livestock to graze treated areas within 7 days of application.
- Do not apply more than 1-1/3 pints (21.3 fluid ounces) of this product per acre per growing season.
- Do not apply within 40 days prior to harvesting grain and straw or within 14 days prior to cutting hay.
- The risk of crop injury at all stages of growth and poor weed control is increased if the application is made and extreme growing conditions (such as drought or near freezing temperatures) occur prior to, at, and following the application.

Application Timing:

To control listed broadleaf weeds, apply as a postemergence broadcast treatment to actively growing wheat, barley or oats from the 2 leaf stage up to and including flag leaf emergence (Zadoks scale 39). Because only weeds that have emerged at the time of application will be controlled, be sure to apply when weeds are actively growing but before weeds are 8 inches tall or vining.

For perennial weeds (such as Canada thistle), apply when the majority of the basal leaves have emerged from the soil up to bud stage to obtain season-long control.

To suppress volunteer potatoes, apply before potato plants are 4 inches tall.

Broadcast Application Rates:

For a complete listing of weeds controlled or suppressed, refer to the “Weeds Controlled or Suppressed” section.

For seedlings of susceptible species < 4 inches tall: Apply 1/2 pint per acre

For seedlings of susceptible species 4 to 8 inches tall or vining: Apply 2/3 pint per acre

For volunteer potatoes: Apply 1-1/3 pints per acre

Note: Kochia seedlings less than 4 inches tall (including ALS resistant biotypes) will be controlled using the 1/2 pint per acre rate. However, when conditions for control are less favorable, such as under drought or cool temperature, a rate of 2/3 pints per acre will provide more consistent control of kochia seedlings 1 to 4 inches tall. For more consistent control of small kochia, apply when the plants are at least 1 inch tall. A rate of 2/3 pints per acre should be used for optimal control of dicamba tolerant kochia populations (refer to the “Management of Kochia Biotypes” section).

Spot Applications:

Spot applications may be made using rates and spray volumes equivalent to a broadcast application (refer to the “Spot Treatments for Crop Uses and On Farm Non-Cropland Uses” section above for additional information).

FIELD CORN

Restrictions:

- Do not broadcast apply to field corn with 6 fully exposed leaf collars (V6 growth stage).
- Do not apply within 90 days prior to harvest of grain or stover.
- Do not harvest forage or allow livestock to graze treated areas within 47 days of application.
- Do not make more than two applications or apply more than 1-1/3 pints of this product per acre per growing season.

Application Timing:

Apply to field corn as a broadcast or band treatment before or when 5 fully exposed leaf collars have developed (the V5 growth stage) using a rate of 2/3 pint per acre. This product may be applied to field corn beyond the V5 growth stage only as a directed spray using drop nozzles.

Apply when broadleaf weeds are less than 8 inches tall and actively growing. Apply to wild buckwheat before the vining stage of growth. To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged up to bud stage.

Preplant Burndown:

Apply this product alone or in tank mix combination with a labeled herbicide prior to planting to control emerged weeds in no-till or burndown situations. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Options for Suppression or Control of Volunteer Potatoes:

For pre-plant suppression applications, apply 2/3 pint of this product per acre when the majority of volunteer potato plants are 4 to 8 inches tall. For best results, leave the soil undisturbed and plant the field corn two weeks after application.

For post emergence suppression applications, apply 2/3 pint of this product per acre when the majority of volunteer potato plants are 4 to 8 inches tall.

For pre-plant and postemergence control applications, a pre-plant application of 2/3 pint of this product per acre may be followed by a postemergence application of 2/3 pint per acre.

Restriction: Do not make more than two applications in a single growing season.

Crop Tolerance Precaution:

When this product is applied as a broadcast treatment, some corn hybrids or lines may experience crop injury (e.g., stem curvature, stunting and brace root injury). In particular, hybrids or lines that are susceptible to phenoxy injury may also be susceptible to injury from this product.

Tank Mixtures for Field Corn:

Unless tank mixing is specifically prohibited by the label of the desired tank mix partner product, this product may be applied in tank mixes containing other herbicides registered for post-emergence application in field corn. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Refer to the "Tank Mixing Precautions" section under MIXING INSTRUCTIONS section of this label for additional information.

Adjuvants:

To improve weed control (especially in hot, dry conditions), using a high quality adjuvant is recommended.

SWEET CORN

Restrictions:

- Do not broadcast apply to sweet corn with 5 fully exposed leaf collars (V5 growth stage).
- Do not apply this product with crop oil concentrates, petroleum-based oils or methylated seed oils.
- Do not apply within 31 days prior to harvesting ears.
- Do not apply within 90 days prior to harvest of grain or stover.
- Do not harvest forage or allow livestock to graze treated areas within 31 days of application.
- Do not make more than two applications or apply more than 1-1/3 pints of this product per acre per growing season.

Application Timing:

Apply to sweet corn as a broadcast or band treatment before or when 4 fully exposed leaf collars have developed (the V4 growth stage) using a rate of 2/3 pint per acre. This product may be applied to sweet corn beyond the V4 growth stage only as a directed spray using drop nozzles.

Apply when broadleaf weeds are less than 8 inches tall and actively growing. Apply to wild buckwheat before the vining stage of growth. To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged up to bud stage.

Preplant Burndown:

Apply this product alone or in tank mix combination with a labeled herbicide prior to planting to control emerged weeds in no-till or burndown situations. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Options for Suppression or Control of Volunteer Potatoes:

For pre-plant suppression applications, apply 2/3 pint of this product per acre when the majority of volunteer potato plants are 4 to 8 inches tall. For best results, leave the soil undisturbed and plant the sweet corn two weeks after application.

For post emergence suppression applications, apply 2/3 pint of this product per acre when the majority of volunteer potato plants are 4 to 8 inches tall.

For pre-plant and postemergence control applications, a pre-plant application of 2/3 pint of this product per acre may be followed by a postemergence application of 2/3 pint per acre.

Restriction: Do not make more than two applications in a single growing season.

Crop Tolerance Precaution:

When this product is applied as a broadcast treatment, some corn hybrids or lines may experience crop injury (e.g., stem curvature, stunting and brace root injury). In particular, hybrids or lines that are susceptible to phenoxy injury may also be susceptible to injury from this product. For further information, consult current sweet corn company herbicide management guidelines.

Tank Mixtures for Sweet Corn:

Unless tank mixing is specifically prohibited by the label of the desired tank mix partner product, this product may be applied in tank mixes containing other herbicides registered for post-emergence application in sweet corn. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Refer to the "Tank Mixing Precautions" section under MIXING INSTRUCTIONS section of this label additional information.

Using Spray Adjuvants in Tank Mixes:

When applying this product alone, spray adjuvants are not recommended because using an adjuvant may increase effectiveness on weeds but can also reduce selectivity to the crop, particularly under conditions of plant stress such as drought or cold temperatures. Be sure to follow all manufacturer guidelines if a tank mix partner requires the addition of an adjuvant.

GRAIN SORGHUM (MILO)**Restrictions:**

- Do not apply after the boot stage.
- Do not make more than two applications per growing season.
- Do not apply more than 1-1/3 pints (21.3 fluid ounces) of this product per acre per growing season.
- Do not apply in combination with metsulfuron methyl.
- Do not allow livestock to graze treated areas or harvest treated forage within 40 days of application.

Application Timing:

Apply to Grain Sorghum as a pre- or post-emergence broadcast treatment using a rate of 2/3 pint per acre. Apply when broadleaf weeds are less than 8 inches tall and actively growing. If wild buckwheat is present, apply before the vining stage of growth. Apply to wild buckwheat before the vining stage of growth. To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged up to bud stage.

For pre-emergence burndown or no-till applications, broadcast apply this product to emerged weeds after planting but prior to grain sorghum emergence.

For post-emergence applications, broadcast apply this product between the 3-leaf and 7-leaf growth stages. From the 8-leaf to boot stage, use drop nozzles and directed spray ONLY. To reduce the potential for crop injury, drop nozzles should direct the spray toward the soil surface to avoid spray contact with grain sorghum foliage.

A pre-emergence application may be followed by a post-emergence application to control heavy weed populations.

Tank Mixtures for Grain Sorghum:

Unless tank mixing is specifically prohibited by the label of the desired tank mix partner product, this product may be applied at labeled rates in tank mixes containing other herbicides registered for post-emergence application in grain sorghum. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Refer to the "Tank Mixing Precautions" section under MIXING INSTRUCTIONS section additional information.

Restriction: Do not apply in combination with metsulfuron methyl.

Adjuvants:

To improve weed control (especially in hot, dry conditions), using a high quality adjuvant is recommended.

MILLET GROWN FOR GRAIN, FORAGE OR HAY

Restrictions:

- Do not apply during boot, flowering or seed development stages if grass crop is to be harvested for seed.
- Do not make more than two applications per growing season.
- Do not apply more than 1-1/3 pints (21.3 fluid ounces) of this product per acre per growing season.
- Do not apply within 14 days of harvesting millet hay or within 40 days of harvesting millet grain and straw.
- Animals to be slaughtered for meat must be removed from treated forage areas at least two days before slaughter.

Application Timing:

Apply to established millet in the spring as a broadcast postemergence treatment using ground equipment or by air.

Apply to new millet plantings from the 2 true leaf stage to just before early boot stage of growth. The potential for injury may be increased if applications are made during or after the boot stage.

Apply when weeds are actively growing, but before weeds are 4 inches tall or vining. For control of late-emerging Canada thistle or kochia, a preharvest treatment may be made after grass seed is fully developed. Postharvest treatments in the fall may be made to actively growing Canada thistle after the majority of basal leaves have emerged. Less consistent control may occur if Canada thistle is at the bud stage or later or kochia greater than 8 inches tall is treated.

Broadcast Application Rates:

For a complete listing of weeds controlled or suppressed, refer to the "Weeds Controlled or Suppressed" section.

For seedlings of susceptible species < 4 inches tall: Apply 1/2 pint per acre

For seedlings of susceptible species 4 to 8 inches tall or vining: Apply 2/3 pints per acre

Note: Kochia seedlings less than 4 inches tall (including ALS resistant biotypes) will be controlled using the 1/2 pint per acre rate. However, when conditions for control are less favorable, such as under drought or cool temperature, a rate of 2/3 pints per acre will provide more consistent control of kochia seedlings 1 to 4 inches tall. For more consistent control of small kochia, apply when the plants are at least 1 inch tall. A rate of 2/3 pints per acre should be used for optimal control of dicamba tolerant kochia populations (refer to the "Management of Kochia Biotypes" section).

A single retreatment may be made a minimum of 14 days after the first treatment.

Tank Mixtures for Millet:

Unless tank mixing is specifically prohibited by the label of the desired tank mix partner product, this product may be applied at labeled rates in tank mixes containing other herbicides registered for post-emergence application in millet. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Refer to the "Tank Mixing Precautions" section under MIXING INSTRUCTIONS for additional information.

GRASSES GROWN FOR SEED, FORAGE OR HAY

This product may be applied for broadleaf weed control in the following Grasses Grown for Seed, Forage or Hay: Bermudagrass, Bluegrass (perennial and annual), Bromegrass, Fescue, Hay Grazer, Orchardgrass, Ryegrass (perennial and annual), Redtop Cane, Sorghum, Sorghum-Sudan, Sudan, Sudex, and Timothy. This product may be applied for broadleaf weed control in the following Grasses Grown for Hay or Forage: Sorghum and Triticale.

Apply this product as a broadcast postemergent treatment by air or using ground equipment. A second application may be made a minimum of 14 days after the first application. This product may be applied in tank mix combination at specified labeled rates with other herbicides registered for these uses. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Application Restriction: Do not apply more than 1-1/3 pints per acre of this product per growing season.

Grazing Restriction: There are no grazing restrictions for lactating or non-lactating dairy animals.

Harvesting Restriction: Do not harvest grass for hay or silage from treated areas within 7 days of application.

Slaughter Restriction: Meat animals must be withdrawn from treated forage at least 2 days before slaughter.

Application Timing:

Apply to established grasses in the Spring when weeds are actively growing and before weeds are 8 inches tall. Only emerged weeds at the time of application will be controlled. New plantings of grass crops may be treated for the 2-true leaf stage of growth prior to early boot stage. Do not apply during boot, flowering, or seed development stages of growth if grass is to be harvested for seed.

BROADCAST APPLICATION RATES	
Weed Size or Species ¹	Application Rate (Pints/Acre)
Susceptible broadleaf weed seedlings less than 4 inches tall ²	1/2
Susceptible broadleaf weed seedlings less than 8 inches tall or vining	2/3

¹ Refer to the "Broadleaf Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.

² The 1/2 pint per acre rate will generally provide satisfactory control of Kochia seedlings less than 4 inches tall (including ALS resistant biotypes). However, when conditions for control are less favorable, such as under drought or cool temperatures; the 2/3 pint per acre rate will provide more consistent control of Kochia seedlings between 1 to 4 inches tall. Control of small Kochia with reduced rates will be more consistent if Kochia is at least 1 inch tall. The 2/3 pint per acre rate should be used for optimal control of Dicamba tolerant Kochia populations (see "Management of Kochia Biotypes" section).

POSTEMERGENCE BROADLEAF WEED CONTROL IN POME FRUIT

(All Crops from Group 11 Including Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear and Quince)

Restrictions:

- Do not make more than one treatment per crop year.
- Do not apply more than 2-2/3 pints per acre per crop year.
- Do not apply within 14 days of harvest.
- Do not apply to trees less than 4-years-old.
- Do not apply during bloom.

Application Timing and Rate:

Apply this product at the rates indicated in the table below as a uniform broadcast postemergence treatment using ground equipment in a minimum of 10 gallons of water per acre. Avoid contact with tree foliage by applying when air temperatures are between 50°F to 80°F and there is no wind.

Note: If this product comes in contact with the tree foliage, leaves and affected portions of the tree may show symptoms or die but the remainder of the tree will remain healthy.

BROADCAST APPLICATION RATES	
Weeds Controlled or Suppressed	Application Rate (Pints/Acre)
Bedstraw (cleavers), Common Purslane, Hairy Buttercup, Hemp Dogbane, Kochia ^{1,2,3} , Marshelder ² , Sericea Lespedeza ² , Tropic Croton	2/3 – 1-1/3
Chickweed, Cocklebur, Coffeeweed, Common Ragweed, Curly Dock, Cutleaf Primrose, Dandelion, Dogfennel, Grape, Horseweed/Marestail, Morningglory, Prickly Lettuce, Puncturevine, Stinging Nettle, Sunflower, Vetch, Velvetleaf, Venice Mallow, Western Ragweed, White Clover, White Cockle	1-1/3
Field Bindweed [†] , Blackberry, Buckhorn Plantain [†] , Carolina Geranium [†] , Catsear, Common Mallow [†] , Common Mullein [†] , Field Cudweed [†] , Giant Ragweed, Goldenrod, Henbane, Field Horsetail [†] , Hop Clover, Horsenettle, Ironweed, Leafy Knotweed [†] , Lantana, Musk Thistle, Mustard [†] , Narrowleaf Plantain [†] , Nightshade spp. [†] , Pennycress [†] , Spiny Amaranth [†] , Spotted Knapweed, Spurge [†] , Wild Buckwheat [†] , Wild Carrot, Yellow Thistle [†]	2-2/3

[†] **Indicates Suppression Only** – Suppression is a reduction in weed competition (i.e., a reduction in population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

¹ Includes herbicide resistant or tolerant biotypes. Apply when weeds are 1 to 4 inches tall for best results.

² Use the higher rate in the range listed when controlling these weeds.

³ To control larger and more advanced kochia, apply 1.5 to 2.0 pints of this product, or tank mix with the labeled rate of 2,4-D Ester and 1 to 2 quarts per acre of methylated seed oil. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. For best results, apply during the 1 to 4 leaf stage of growth (before vining).

Tank Mixtures for Pome Fruit:

Unless tank mixing is specifically prohibited by the label of the desired tank mix partner product, this product may be applied in tank mixes containing other herbicides registered for post-emergence application in pome fruit. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Refer to the "Tank Mixing Precautions" section under MIXING INSTRUCTIONS section for additional information.

FALLOW CROPLAND

Restriction:

- Do not plant any crops within 120 days of application.

To control susceptible broadleaf weeds and volunteer potatoes, apply 2/3 to 1-1/3 pints of this product as a single broadcast treatment using ground or aerial equipment (refer to the **APPLICATION INFORMATION** section for specific information on weeds controlled). Apply when weeds are less than 8 inches tall or not vining and actively growing. This product may be applied alone or in tank-mix combination with other herbicides registered for use on fallow croplands (See "Tank Mixing Precautions" under **MIXING INSTRUCTIONS** section). It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If weeds are under stress from drought or extreme temperatures, control may be reduced. For light to moderate weed infestations and under good growth conditions, use the lower rates listed. For moderate to heavy infestations and to compensate for less than ideal growth conditions, use the higher rates listed.

ON-FARM NON-CROPLAND

This product may be applied as a single broadcast treatment or spot treatment to control susceptible broadleaf weeds in on-farm non-cropland areas (fencerows, building perimeters, around irrigation equipment and on-farm private roadways). Apply at the rate of 2/3 to 1-1/3 pints per acre when weeds are actively growing but less than 8 inches tall or not vining.

Spot treatments should be applied at rates and spray volumes equivalent to broadcast application. Refer to the instructions for "Spot Treatments for Crop Uses and On Farm Non-Cropland Uses" in the **APPLICATION INFORMATION** section for more information.

Refer to the **APPLICATION INFORMATION** section of this label for a complete listing of weeds controlled or suppressed.

CRP ACRES**Restrictions:**

- Do not use on CRP acres that are under seeded with desirable legumes, clovers, or other sensitive broadleaf plants.
- Grazing or haying of treated CRP acres is prohibited.

Application Instructions:

To control susceptible broadleaf weeds apply this product as a single broadcast treatment using ground or aerial equipment. Apply at the rate of 2/3 to 1-1/3 pints per acre when weeds are actively growing but less than 8 inches tall or not vining.

Spot treatments should be applied at rates and spray volumes equivalent to broadcast application. Refer to the instructions for "Spot Treatments for Crop Uses and On Farm Non-Cropland Uses" in the **APPLICATION INFORMATION** section for more information.

Refer to the **APPLICATION INFORMATION** section of this label for a complete listing of weeds controlled or suppressed.

NON-CROP USE

- Rangeland and permanent pastures
- Non-cropland areas including industrial sites, non-irrigation ditch banks, and rights-of-way (electrical power lines, communication lines, pipelines, roadsides and railroads including grazed areas within these sites)
- Pine plantations

Non-Crop Use Restrictions:

- Do not use on plants grown for agricultural or commercial production (such as designated grazing areas) in the state of Arizona.
- See individual use directions for additional restrictions, including maximum application rates.
- When applying this product, do not contaminate water used for domestic purposes or irrigation ditches.
- Do not apply this product through any type of irrigation system (i.e., chemigation).
- Do not apply this product in or around greenhouses.
- Do not allow spray drift to come in contact with or apply this product directly to susceptible broadleaf plants or broadleaf crops, including but not limited to the following: alfalfa, canola, cotton, edible beans, grapes, lentils, lettuce, melons, mustard, peas, potatoes, radishes, safflower, soybeans, sugar beets, sunflower, tobacco and tomatoes or other vegetable crops, flowers, fruit trees, ornamentals and shade trees.
- Do not store or handle other agricultural chemicals using this products container.
- Do not apply other agricultural chemicals or pesticides with equipment used to apply this product until the equipment has been thoroughly cleaned (refer to the “Sprayer Cleanup” section under MIXING INSTRUCTIONS for additional information.).
- Do not harvest grass for hay or silage from treated areas within 7 days of application.
- Animals to be slaughtered for meat must be removed from treated forage areas at least two days before slaughter.

APPLICATION INFORMATION

Broadleaf Weeds Controlled or Suppressed for Non-Crop Use			
Weed	Application Rate (Pints per Acre)	Weed	Application Rate (Pints per Acre)
Bedstraw (cleavers)	2/3 to 1-1/3	*Knotweed [†]	2-2/3
Blackberry	2-2/3	Kochia ^{1, 2, 3}	2/3 to 1-1/3
Buckhorn plantain [†]	2-2/3	Lantana	2-2/3
Carolina geranium [†]	2-2/3	Leafy spurge [†]	2-2/3
Catsear	2-2/3	Marshelder ²	2/3 to 1-1/3
Chickweed	1-1/3	Morningglory	1-1/3
Cocklebur	1-1/3	Musk thistle	2-2/3
Coffeeweed, common	1-1/3	Mustard [†]	2-2/3
*Common mallow [†]	2-2/3	Narrowleaf plantain [†]	2-2/3
Common mullein [†]	2-2/3	Nightshade spp. [†]	2-2/3
Common purslane	2/3 to 1-1/3	Prickly lettuce	1-1/3
Cudweed [†]	2-2/3	*Puncturevine	1-1/3
Curly dock	1-1/3	Ragweed	1-1/3
Cutleaf primrose	1-1/3	Sericea lespedeza ²	2/3 to 1-1/3
Dandelion	1-1/3	Spiny amaranth [†]	2-2/3
Dogfennel	1-1/3	*Spotted knapweed	2-2/3
Field bindweed [†]	2-2/3	*Stinging nettle	1-1/3
Field horsetail [†]	2-2/3	Sunflower	1-1/3

Field pennycress [†]	2-2/3	Tropic croton	2/3 to 1-1/3
Giant ragweed	2-2/3	Velvetleaf	1-1/3
Goldenrod	2-2/3	Venice mallow	1-1/3
Grape spp.	1-1/3	Vetch	1-1/3
Hairy buttercup	2/3 to 1-1/3	Western ragweed	1-1/3
Hemp dogbane	2/3 to 1-1/3	White clover	1-1/3
Henbane	2-2/3	White cockle	1-1/3
Hop clover	2-2/3	Wild buckwheat [†]	2-2/3
Horsenettle	2-2/3	Wild carrot	2-2/3
Horseweed / marestalk	1-1/3	Yellow thistle [†]	2-2/3
Ironweed	2-2/3		

[†] **Indicates Suppression Only** – Suppression is a reduction in weed competition (i.e., a reduction in population or vigor) as compared to untreated areas. The degree of weed control and duration of effect may vary with weed size, density, application rate, coverage, and growing conditions before, during and after treatment.

¹ Includes biotypes that are herbicide resistant or tolerant.

² To control these weeds, use the higher rates in the range listed.

³ When controlling kochia, adding a methylated seed oil surfactant (i.e., MSO or ESO) at a rate of 1 to 2 quarts per acre is recommended. Improved control for infestations of larger kochia plants at more advanced growth stages may be achieved by increasing the rate of this product to 1.5 to 2.0 pints per acre OR adding 1 to 2 quarts of 2,4-D Ester and 1 to 2 quarts of methylated seed oil per acre.

***Not for this weed in California**

Spot Treatments for Non-Crop Use:

Only apply using a calibrated boom sprayer or with a hand sprayer using the following directions:

When using hand-held sprayers for spot applications, be sure to uniformly apply a rate equivalent to a broadcast application. Application rates in the table below are based on an area of 1,000 square feet.

Mix the amount of this product corresponding to the desired broadcast rate in one or more gallons of spray. To calculate the amount of this product required for larger areas, multiply the table value (fluid ounces or ml) by the area to be treated in “thousands” of square feet. An area of 1,000 square feet is approximately 10.5 x 10.5 yards (strides) in size.

For example: If the area to be treated is 3,500 square feet, multiply the table value by 3.5 (calc. $3,500 \div 1,000 = 3.5$).

Broadcast Rate Conversions for Spot Treatments for Non-Crop Use	
Broadcast Rate (Pints per Acre)	LPI.A011 per Gallon (Fluid Ounces)
2/3	0.25 (7.25 ml)
1	0.37 (11 ml)
1-1/3	0.50 (14.5 ml)
2	0.74 (22 ml)
2-2/3	1.0 (30 ml)

RANGELAND AND PERMANENT GRASS PASTURES

Note: There are no grazing restrictions for livestock (including lactating or non-lactating dairy animals).

Restrictions:

- Do not apply more than 2-2/3 pints of this product per acre per year.
- Do not apply if injury to legumes cannot be tolerated as this product may injure or kill legumes. Legumes may be less sensitive to herbicide injury after plant growth is mature and seed has set.
- Do not allow meat animals to graze on treated forage within 2 days before slaughter.
- Do not harvest grass for hay or silage from treated areas within 7 days of application.
- Only forage grasses, wheat, barley, oats, field corn, sweet corn and grain sorghum may be planted in treated fields within 120 days following application of this product.

Application Timing:

Apply this product when weeds are actively growing (but prior to bud stage of weed growth) as a sequential postemergence broadcast treatment or as a single broadcast treatment using ground or aerial application equipment. Use a total spray volume of 5 or more gallons per acre for ground broadcast application or 3 or more gallons per acre by air.

Note: Only weeds that have emerged at the time of application will be controlled.

Tank Mixtures:

For control of additional weeds and woody plants, this product may be applied as a tank mix with other foliar-applied herbicides labeled for use on rangeland and permanent pastures. When tank mixing, be sure to follow the most restrictive use directions, precautions and limitations on each product label. Rates for tank mixes containing either triclopyr or picloram are listed in the following table:

Tank Mix	Rates (per Acre)	Additional Weeds / Brush Controlled		
LPI.A011 + Triclopyr	1/2 pint + Labeled rate	Buttercup, hairy Croton Dogbane, hemp Kochia	Lespedeza, sericea Marshelder Ragweed Sunflower	Thistle, musk Vetch
LPI.A011 + Triclopyr	2/3 pint + Labeled rate	Dandelion Dock, curly Dogfennel	Goldenrod Horseweed / Marestail Ironweed	Lantana Plantain
LPI.A011 + Triclopyr	1 pint + Labeled rate	Blackberry Persimmon Rose, multiflora	Tropical soda apple Wax myrtle	
LPI.A011 + Picloram	2/3 pint + Labeled rate	Bindweed, Field Broomweed, annual Buttercup, hairy Cocklebur Croton Dogbane, hemp	Dogfennel Goldenrod Horsenettle Horseweed Kochia Lespedeza, sericea	Marshelder Mullein Ragweed Sneezeweed, bitter Sunflower Thistle, musk Vetch
LPI.A011 + Picloram	1-1/3 pint + Labeled rate	Blackberry Locust Plum, wild	Prickly pear cactus Rose, Cherokee Rose, Macartney	Rose, multiflora Sumac

NON-CROPLAND AND PINE PLANTATIONS

(Includes the following sites and grazed areas within them: Industrial sites, non-irrigation ditch banks, and rights-of-way (power lines, communication lines, pipelines, roadsides and railroads))

Restrictions:

- Do not apply this product to pine plantations as an over-the-top broadcast treatment during active terminal growth (from initiation of budbreak/growth flush until seasonal terminal growth has hardened off and overwintering buds have formed). Directed spray applications may be made to pine plantations during periods of active growth, avoid spray contact with actively growing foliage.
- Do not apply this product to pine plantations in a tank mix unless the tank mix partner product is labeled for application using the desired method to control weed or brush in pines. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Application Timing:

Broadcast apply this product when weeds are small and / or actively growing using a rate of 2/3 to 2-2/3 pints per acre (refer to the "Weeds Controlled or Suppressed" section of this label for specific rate directions). This product may be split applied in a single year as long as the total amount of this product applied does not exceed the maximum labeled rate of 2-2/3 pints per acre. For spot treatments, apply at rates and spray volumes equivalent to broadcast application (refer to the instructions for "Spot Treatments for Non-Crop Use" in the APPLICATION INFORMATION section for additional information).

Tank Mixtures:

To improve control of pine species, shingle oak, red maple, red oak and other woody species, this product may be tank-mixed with triclopyr, picloram and glyphosate as indicated in the following table:

Tank Mix	Rates (per Acre)	Additional Brush / Trees Controlled	
LPI.A011 + Triclopyr	32 to 42 fluid ounces + Labeled rate	Bay spp. Black cherry Dogwood	Water oak Willow oak
LPI.A011 + Triclopyr + Picloram	32 to 42 fluid ounces + Labeled rate + Labeled rate	Pine spp. Red maple Red oak	Shingle oak Virginia pine Water oak
LPI.A011 + Glyphosate	32 to 42 fluid ounces + Labeled rate	Dogwood Gallberry	Pine spp. Wax myrtle

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store above 10°F or warm and agitate before use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

Nonrefillable Containers (≤ 5 gallons): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Containers (> 5 gallons): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Refillable Containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the entire Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way.

Follow the Directions for Use of this product carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop or other plant injury, ineffectiveness, or other unintended consequences may result from such risks as weather or crop conditions, mixture with other chemicals not specifically identified in this product's label, or use of this product contrary to the label instructions, all of which are beyond the control of LOVELAND PRODUCTS, INC. and the seller. To the extent consistent with applicable law, the buyer or user of this product assumes all such inherent risks.

Subject to the foregoing inherent risks, LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use when the product is used in strict accordance with such Directions for Use under normal conditions of use. EXCEPT AS WARRANTED IN THIS LABEL AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS PRODUCT IS SOLD "AS

IS," AND LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE.

IN THE UNLIKELY EVENT THAT BUYER OR USER BELIEVES THAT LOVELAND PRODUCTS, INC. HAS BREACHED A WARRANTY CONTAINED IN THIS LABEL AND TO THE EXTENT REQUIRED BY APPLICABLE LAW, BUYER OR USER MUST SEND WRITTEN NOTICE OF ITS CLAIM TO THE FOLLOWING ADDRESS: LOVELAND PRODUCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT, INCLUDING BUT NOT LIMITED TO CLAIMS OF BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, STRICT LIABILITY, OR OTHER TORTS, SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

[LPI.A011] is a trademark of Loveland Products, Inc.

{LANGUAGE ON LABEL AFFIXED TO CONTAINER}

FLUROXYPYR	GROUP	4	HERBICIDE
------------	-------	----------	-----------

LPI.A011™

Active Ingredient: (by weight)
 Fluroxypyr: 1-methylheptyl ester: ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic acid, 1-methylheptyl ester26.2%
 Other Ingredients73.8%
 Total100.0%
 Contains 1.5 pounds fluroxypyr acid per gallon (18.2%)
 Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

NOT FOR SALE, SALE INTO, DISTRIBUTION AND/OR USE IN NASAU AND SUFFOLK COUNTIES OF NEW YORK STATE.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Do not give any liquid to the person
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
Note to Physician: Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Drift or runoff from treated areas may be hazardous to aquatic organisms and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store above 10°F or warm and agitate before use. In case of spill or leak on floor or paved surfaces, soak up with sand, earth, or synthetic absorbent. Remove to chemical waste area.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING

Nonrefillable Containers (≤ 5 gallons): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Containers (> 5 gallons): Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Refillable Containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC – 1-800-424-9300.

See inside label booklet for additional Precautionary Statements and Directions for Use.

EPA Reg. No. 34704-1157

EPA Est. No.

Net Weight:

[Label ID Print Code]

MANUFACTURED FOR:

LOVELAND PRODUCTS, INC.

P.O. BOX 1286

GREELEY, COLORADO 80632-1286