



## USE DIRECTIONS

**Disinfection** — Use at 2 ounces per gallon. AN-CARE STAPH MINUS II is effective against PSEUDOMONAS AERUGINOSA PRD-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use STAPH MINUS II with a cloth or mop and apply to walls, floors and other hard surfaces.

**Sanitization** — If used for sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (2.8 ounces per 5 gallons). NO RINSE REQUIRED AT THIS LEVEL.


For heavy soil or organic matter, pre-cleaning step is recommended.

\*STAPH MINUS II when used on environmental, inanimate, hard surfaces at 2 ounces per gallon of water, is effective against INFLUENZA A<sup>2</sup>, HERPES SIMPLEX, ADENOVIRUS TYPE 2 and VICCINIA VIRUSES.

### PHENOL COEFFICIENTS:

Guaranteed	Minimum
S. AUREUS	42
S. TYPHOSA	27

Rinse empty container thoroughly with water and discard.



NET CONTENTS  
ONE U.S. GAL.

**STAPH**  
**DISINFEC**  
**DEOD**  
**BA**

ACTIVE INGREDIENTS  
N-Alkyl (60% C  
dimethyl b  
N-Alkyl (68% C  
benzyl amr  
Sodium Carbon  
Tetrasodium Ethy  
INERT INGREDIENTS

**WARNING**  
See First Aid Stat

# ANDERSON

ANDERSON CHEMICAL COMPANY