as amandod wuler 4 7 27 - 2 registered unuer 4 7 7 27 - 2

PROPOSED LABEL COPY

PRODUCT NAME:

Flea & Tick Spray

LABEL DIMENSIONS

FORMULATION/

PRODUCT NUMBER:

AUTHORED BY:

REVIEWED BY:

Under the Federal Insections of paper weight Fundanded inter and a control of the paper weight of superior and superior of the paper weight of superior and superior of the paper weight of superior and superior of the paper weight of superior of the paper weight of the paper wei

Flea & Tick Spr:

For use on cats and dogs ontrol fleas, ticks, and ear mites.

ACTIVE INGREDIENTS:

w/w

*Piperonyl Butoxide, technical

0.60% 0.06%

Pyrethrins INERT INGREDIENTS:

99.34% 100.00%

*Equivalent to 0.48% (butylcarbityl) (6-propylpiperonyl) ether and 0.12% related compounds.

FOR EXTERNAL AND VETERINARY USE ONLY

KEEP OUT OF REACH OF CHILDREN

CAUTION: See side panel for additional precautions.

EPA Req. No. 34797-2

EPA Est. No. 34797-1A-03

Manufactured by: Qualis, Inc.

Des Moines, Iowa 50321

NET CONTENTS:

16 fluid ounces

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE

For Fleas and Ticks: Hold bottle upright and point nozzle opening in desired direction of spray. Spray the animal from a distance of 3-12 inches.

Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under body. While spraying, fluff the hair so that spray will penetrate to the skin. Make sure spray wets thoroughly. Do not spray into eyes or face. Repeat as necessary.

For Ear Mites: Shake well before use; cleanse ear thoroughly. Apply 5 drops into each ear twice daily. Repeat treatment if necessary.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION:

Harmful if swallowed. Avoid inhalation of vapors. Avoid contact with skin. Wash contaminated skin properly with soap and water. Avoid contamination of food or foodstuffs. For external use only. In case of infection or skin irritation, discontinue use and consult a veterinarian.

STORAGE: Do not transport or store below 32°F. DISPOSAL: Do not reuse empty bottle. Wrap bottle and put in trash.