DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Apply undiluted to hair and scalp or any other infested areas until entirely wet. Do not use on eyelashes or eyebrows. When applying product for Head Lice, place towel across forehead. Allow treatment to remain 10 minutes, but no longer. Then wash thoroughly with warm water and shampoo or soap. Dead lice and eggs may require removal with enclosed fine-toothed comb. Repeat treatment in 7-10 days or earlier if reinfestation has occurred. Do not exceed two applications within 24 hours. To help eliminate infestations, sterilize or treat all clothing and bedding of infested person at time of treatment. When used on children, adult supervision is recommended.

STORAGE AND DISPOSAL: Store product in original container above freezing and away from children; wrap in several layers of newspaper before discarding in trash.

LICE KILLING

Shampoo

Liquid pediculicide for human use. Kills head lice, body lice, crab lice and their eggs on contact.

ACTIVE INGREDIENTS:

Pyrethrins	0.33%
*Piperonyl Butoxide, Technical	4.00%
INERT INGREDIENTS:	95.67%
	100.00%

*Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and 0.8% related compounds.

KEEP OUT OF REACH OF CHILDREN.
CAUTION: SEE OTHER PRECAUTIONS ON SIDE PANEL

Manufactured By:
Qualis, Inc.
Des Moines, Iowa 50321

Net Contents: (1 Gallon or smaller)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION:

Harmful if swallowed, absorbed through skin or inhaled. Causes eye irritation/injury. Avoid contact with skin, eyes or clothing. Avoid breathing dust (vapor or spray mist). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

USE PRECAUTIONS: This product is for external use only. Should be used with caution by ragweed sensitized persons. In case of infection or skin irritation, discontinue use and consult a physician.

STATEMENT OF PRACTICAL TREATMENT:

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If in eyes: Hold eyelids open and flush with a steady, gentle stream of water. Call a physician if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

EPA Reg. NO. 34797-83
EPA Est. NO. 34797-IA-3

MAR 12 '9

BEST AVAILABLE COPY

34797-83