



WEX-SAN

CONCENTRATED GERMICIDAL DETERGENT

SANITIZER, FUNGICIDE, DISINFECTANT,
DEODORIZER, PSEUDOMONACIDAL, VIRUCIDAL*

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chlorides	4.5%
n-Alkyl (68% C12, 32% C14) dimethyl	
ethylbenzyl ammonium chlorides	4.5%
Tetrasodium ethylene diamine tetraacetate	2.0%
Sodium Carbonate	4.0%
Essential Oils	0.2%
INERT INGREDIENTS:	84.8%
	100.0%

EPA REGISTRATION NO. 34810-1

DANGER: KEEP OUT OF REACH OF CHILDREN.

See side panel for additional precautionary and first aid statements.

Wexford Labs, Inc.

1035 South Vandeventer

St. Louis, Missouri 63110

FOR INDUSTRIAL AND INSTITUTIONAL USE

CONTENTS ONE U. S. GALLON

DIRECTIONS

Disinfecting

For all disinfection, including hospital and health care facilities, use WEX-SAN at a dilution of 1 ounce per gallon of water.

* When used at a dilution of 1 ounce per gallon of water on environmental inanimate surfaces, WEX-SAN is effective against Influenza A2, Herpes Simplex, Adenovirus Type 2, and Vaccinia viruses.

For heavy soil, or organic matter, a pre-cleaning step is recommended.

Sanitizing

For all sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (1.4 ounces per 5 gallons of hard** or soft water). Food contact surfaces must be rinsed with potable water before reuse.

WEX-SAN solution may be applied with cloth, mop, or brush, or by immersing equipment to be sanitized.

Rinse empty container thoroughly with water and discard.


** WEX-SAN when used for sanitization is effective in water up to 250 ppm of hardness calculated as CaCO3.

WEX-SAN
cally des-
uses for
surfaces.

WEX-SAN
deodorizi
suited fo
Emergen

When usi
lococcus
inosa, Pa

When WI
processin
active q
rinsed w



WEX-SAN

WEX-SAN, a Concentrated Germicidal Detergent, is specifically designed for hospitals, health care and other institutional uses for the control of cross-infection from environmental surfaces.

WEX-SAN is used for cleaning, disinfecting, sanitizing and deodorizing, floors, walls and other hard surfaces. Ideally suited for use in Operating Rooms, Intensive Care, Nurseries, Emergency Areas, Food Service Areas, Rest Rooms.

When used as directed **WEX-SAN** is effective against Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, Pathogenic fungi, and Lipophylic viruses.

When **WEX-SAN** is used for sanitizing previously cleaned food processing equipment, or food contact items, at 200 ppm of active quaternaries, these food contact surfaces must be rinsed with potable water before reuse.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food. Protect eyes and skin when handling concentrate.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Wexford Labs, Inc.

EPA ESTABLISHMENT NUMBER 34810-MO-1

1 health
1 ounce

allon of
EX-SAN
simplex,

ing step

d equip-
of active
allons of
must be

h, mop,
sanitized.

ater and

ective in
CaCO₃.