JUN 2 1994

Under the Federal Insert, 11e, Fungi de mat Patendiciae Act. as amended, for the pesticide regisored undar 4810-2



SUPER WEX-SAN

CONCENTRATED DETERGENT/DISINFECTANT Cleaner, Detergent, Deodorizer, Mildewstat, Disinfectant, Fungicide, Virucide*

ACTIVE INCREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)

n-AlkvI (68% C12, 32% C14)dimethyl INERT INGREDIENTS: 87.50%

EPA REG. NO. 34810-2

KEEP OUT OF REACH OF CHILDREN DANGER

See back panel for additional precautionary and first aid statements.

Wexford Labs, Inc.

325 Leffingwell Avenue

Kirkwood, Missouri 63122

FOR INDUSTRIAL AND INSTITUTIONAL USE

Contents: One U.S. Gallon

BACTERICIDAL ACTIVITY-

SUPER WEX-SAN Germicidal Detergent exhibits effective disinfectant activity against Pseudomonas aeruginosa PRD 10, Escherichia coli, Klebsiella pneumoniae, Salmonella schottmuelleri, Salmonella choleraesuis, Streptococcus salivarius, Brevibacterium ammoniagenes, Streptococcus faecalis, Shigella dysenteriae, Enterobacter serogenes and

FUNGICIDAL ACTIVITY:

SUPER WEX-SAN is fungicidal against the pathogenic fungi, Tricophyton mentagrophytes, in organic soil (5% blood serum) at.....

*VIRUCIDAL ACTIVITY:

SUPER WEX-SAN is effective against Influenza A2/Japan (representative of the common flu virus) Herpes Simplex Type 1 (causative agent of fever blisters), Adenovirus Type 5 (causative agent of upper respiratory infections), and Vaccinia virus (representative of the pox virus group), and HIV-1 (associated with AIDS) in organic soil (5% blood serum) on hard, inanimate sufaces at ... 1:256

SANITIZATION

To sankize previously cleaned and rinsed non-porous food comact surfaces prepare a 200 ppm active quaternary solution by adding 1 ounce of SUPER WEX-SAN to 5 gallons of water. Apply solution with cloth, mop, brush, or by immersing equipment to be sanitized. When used for sankization of previously cleaned food equipment or food contact items, limit active quaternaries to 200 ppm. At this level NO POTABLE WATER RINSE IS REQUIRED, SUPER WEX-SAN IS an effective sanitizer when diluted in water up to 600 ppm hardness (CaCO3).

STORAGE AND DISPOSAL


DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL STORAGE - Store in a dry place no lower in temperature than 50° F or higher than 120° F.

CONTAINER DISPOSAL - Do not reuse empty container. Triple tinse empty container with water. Puncture metal drums and dispose of in a sanitary landfill or by other procedures approved by State and local authorities. Plastic containers may be disposed of in a sanitary landfill, incinerated, or if allowed by local authorities, by burning, if burned stay out of smoke,

PESTICIDE DISPINSAL - Pesticide wastes are acutely hazardous, improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law, if these wastes cannot be disposed of by use according to label instructions contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.







SUPER WEX-SAN

SUPER WEX-SAN is a phorphate free, concentrated germicidal detergent specifically designed for hospitals, health-care, schools, food processing plants, and other institutional uses for the control of cross-infection from environmental surfaces. SUPER WEX-SAN, when used as directed is formulated to disinfect inanimate hard surfaces such as walls, floors, sink tops, toilet bowls, tables, chairs, telephones, and bed frames. When used in such areas as operating rooms, patient care facilities, and restrooms SUPER WEX-SAN is designed to provide both general cleaning and disinfecting. It deodorizes those areas which are generally hard to keep fresh smelling such as garbage bins and cans, toilet bowls, and other areas which are prone to odor causing microorganisms.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. To disinfect inanimate hard surfaces add 1/2 ounce SUPER WEX-SAN per gallon of water. Apply solution with a mop, cloth, sponge or brush so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove excess liquid. For heavily soiled areas, a pre-cleaning step is required. Prepare a fresh solution for each use. To disinfect toilet bowls, flush toilet, add 1/2 ounce SUPER WEX-SAN directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 10 minutes and flush.

*SUPER WEX-SAN KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health-care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the

surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS.)

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks or eye coverings. CLEANING PROCEDURES: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of SUPER WEX-SAN germicidal detergent. CONTACT TIME: Allow surface to remain wet for 10 minutes. DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS. DANGER. KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food, Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink egg whites or gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Wexford Labs, Inc.

EPA ESTABLISHMENT NUMBER 34810-MO-1

BEST AVAILABLE COPY

