



WEX-SAN-M

CONCENTRATED GERMICIDAL DETERGENT

SANITIZER, FUNGICIDE, DISINFECTANT,
DEODORIZER, PSEUDOMONACIDAL, VIRUCIDAL *

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chlorides	2.25%
n-Alkyl (68% C12, 32% C14) dimethyl	
ethylbenzyl ammonium chlorides	2.25%
Sodium Carbonate	3.00%
Tetrasodium ethylenediamine tetraacetate	1.00%
INERT INGREDIENTS:	91.50%
	100.00%

EPA REGISTRATION NO. 34810-3

WARNING

KEEP OUT OF REACH OF CHILDREN.

See side panel for additional precautionary and first aid statements.

Wexford Labs, Inc.

1035 South Vandeventer

St. Louis, Missouri 63110

FOR INDUSTRIAL AND INSTITUTIONAL USE
CONTENTS ONE U. S. GALLON

ACCEPTED

AUG 21 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 34810-3

DIRECTIONS

Disinfecting

For all disinfection, including hospital and health care facilities, use **WEX-SAN-M** at a dilution of 2 ounces per gallon of water.

*When used at a dilution of 2 ounces per gallon of water on environmental inanimate surfaces, **WEX-SAN-M** is effective against Influenza A₂, Herpes Simplex, Adenovirus Type 2, and Vaccinia viruses.

For heavy soil, or organic matter, a pre-cleaning step is recommended.

Sanitizing

For all sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (2.8 ounces per 5 gallons of hard** or soft water). Food contact surfaces must be rinsed with potable water before re-use.

WEX-SAN-M solution may be applied with cloth, mop, or brush, or by immersing equipment to be sanitized.

Rinse empty container thoroughly with water and discard.

** **WEX-SAN-M** when used for sanitization is effective in water up to 250 ppm of hardness calculated as CaCO₃.

WEX-SAN-M is specific of cross-infection from

WEX-SAN-M is used for surfaces. Ideally suited for Service Areas, Rest Rooms

When used as directed against *Pseudomonas aeruginosa*

When **WEX-SAN-M** is used on items, at 200 ppm of active quaternaries before re-use.

Keep Out of Reach of Children. Harmful if swallowed.

In case of contact, immediately call a physician. Remove egg whites, gelatin solutions

DIRECTIONS

Disinfecting

For all disinfection, including hospital and health care facilities, use **WEX-SAN-M** at a dilution of 2 ounces per gallon of water.

When used at a dilution of 2 ounces per gallon of water on environmental inanimate surfaces, **WEX-SAN-M** is effective against Influenza A₂, Herpes Simplex, Adenovirus Type 2, and Vaccinia viruses.

For heavy soil, or organic matter, a pre-cleaning step is recommended.

Sanitizing

For all sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (2.8 ounces per 5 gallons of hard** or soft water). Food contact surfaces must be rinsed with potable water before re-use.

WEX-SAN-M solution may be applied with cloth, mop, or brush, or by immersing equipment to be sanitized.

Rinse empty container thoroughly with water and discard.

WEX-SAN-M when used for sanitization is effective in water up to 250 ppm of hardness calculated as CaCO₃.

WEX-SAN-M

A CONCENTRATED GERMICIDAL DETERGENT

WEX-SAN-M is specifically designed for hospitals, health care and other institutional uses for the control of cross-infection from environmental surfaces.

WEX-SAN-M is used for cleaning, disinfecting, sanitizing and deodorizing, floors, walls and other hard surfaces. Ideally suited for use in Operating Rooms, Intensive Care, Nurseries, Emergency Areas, Food Service Areas, Rest Rooms.

When used as directed **WEX-SAN-M** is effective against Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, pathogenic and Lipophylic viruses.

When **WEX-SAN-M** is used for sanitizing previously cleaned food processing equipment, or food contact items, at 200 ppm of active quaternaries, these food contact surfaces must be rinsed with potable water before re-use.

WARNING

Keep Out of Reach of Children. Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before re-use. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

Wexford Labs, Inc.

EPA ESTABLISHMENT NUMBER 34810-MO-1