



DAIRY-WEX

**MULTI PURPOSE CLEANER and SANITIZER
FOR THE
DAIRY INDUSTRY**

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18)	
dimethyl benzyl ammonium chlorides	5.0%
n-Alkyl (68% C12, 32% C14) dimethyl	
ethylbenzyl ammonium chlorides	5.0%
Phosphoric Acid	30.0%
INERT INGREDIENTS:	60.0%
	100.0%

EPA REGISTRATION NO. 34810-4

DANGER: KEEP OUT OF REACH OF CHILDREN.

See rear panel for additional precautionary and first aid statements.

Wexford Labs, Inc.

1035 South Vandeventer

St. Louis, Missouri 63110

**FOR INDUSTRIAL AND INSTITUTIONAL USE
CONTENTS ONE U. S. GALLON**

ACCEPTED

OCT 28 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. *34810-4*

Comment

DAIRY-WEX SANITIZES — CI

DIRECTIONS

DAIRY-WEX is easily and quickly dispersed in hot or cold water to 1700 ppm of hardness (as calcium carbonate) to form a complete solution. **DAIRY-WEX** when used at a 1:500 aqueous dilution (2 quaternaries) may be used on food processing and dairy equipment with the necessity of a final potable rinse.

TO CLEAN BULK MILK TANKS:

As soon as possible after milk has been removed from bulk tank, scrub completely with cold water and allow to drain. Place 4 gallons 120° F. in tank and add 1 ounce (2 tablespoons) **DAIRY-WEX** to tank. Scrub tank, mixers and covers and then allow to drain.

TO SANITIZE MILK EQUIPMENT AND COW UDDERS

Previously cleaned milking equipment, pail, inflations and tubs should be sanitized prior to milking operations by preparing a 200 ppm solution of quaternaries by mixing 1 ounce (2 tablespoons) **DAIRY-WEX** water. Scrub or dip equipment into the solution and allow to begin milking operations.

Use a clean single service towel for each animal to wipe udders with sanitizing solution prior to milking. Never reuse towel or dip udders into solution. Always use freshly prepared solutions and rinsing utensils.

HARD WATER TOLERANCE OF **DAIRY-WEX**

This product fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1965 recommendations of the U. S. Public Health Service.

ACCEPTED

OCT 28 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 34810-4

Comments

DAIRY-WEX

DAIRY-WEX SANITIZES — CLEANS — REMOVES MILKSTONE — MIXES EASILY

DIRECTIONS

DAIRY-WEX is easily and quickly dispersed in hot or cold water and in water up to 1700 ppm of hardness (as calcium carbonate) to form a completely uniform solution. **DAIRY-WEX** when used at a 1:500 aqueous dilution (200 ppm active quaternaries) may be used on food processing and dairy equipment without the necessity of a final potable rinse.

TO CLEAN BULK MILK TANKS:

As soon as possible after milk has been removed from bulk tank, rinse tank completely with cold water and allow to drain. Place 4 gallons of warm water 120° F. in tank and add 1 ounce (2 tablespoons) **DAIRY-WEX** to the water in the tank. Scrub tank, mixers and covers and then allow to drain.

TO SANITIZE MILK EQUIPMENT AND COW UDDERS

Previously cleaned milking equipment, pail, inflators and tubing should be sanitized prior to milking operations by preparing a 200 ppm solution of active quaternaries by mixing 1 ounce (2 tablespoons) **DAIRY-WEX** in 4 gallons of water. Scrub or dip equipment into the solution and allow to drain prior to beginning of milking operations.

Use a clean single service towel for each animal to wipe udders with the above sanitizing solution prior to milking. Never reuse towel or dip used towel back into solution. Always use freshly prepared solutions and non-corrodible utensils.

HARD WATER TOLERANCE OF **DAIRY-WEX**

This product fulfills the criteria of Appendix F of the Grade A Pasteurized Milk Ordinance 1965 recommendations of the U. S. Public Health Service in waters

up to 600ppm of hardness calculated as calcium carbonate, when tested by the AOAC Germicidal and Detergent/Sanitizing Official Method. Consult your milk inspector or health officer for recommendations on your local program.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.

Rinse empty container thoroughly with water and discard.

Wexford Labs, Inc.

EPA ESTABLISHMENT NUMBER 34810-MO-1

ACCEPTED WITH COMMENTS