

ACCEPTED
 SEP 22 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 EPA Reg. No. 34810-5
 registered under



SANI-WEX

**DISINFECTANT, SANITIZER, DEODORIZER
 PSEUDOMONACIDAL, VIRUCIDAL***

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides	5%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl Ethylbenzyl ammonium chlorides	5%
INERT INGREDIENTS	90%
	100%

SANI-WEX is a concentrate and should be diluted before using.

EPA REGISTRATION NO. 34810-5

DANGER: KEEP OUT OF REACH OF CHILDREN.

See side panel for additional precautionary and first aid statements.

Wexford Labs, Inc.

1035 South Vandeventer

St. Louis, Missouri 63110

DIRECTIONS

Use SANI-WEX at a dilution of 3 ounces per 5 gallons of water for disinfecting floors, walls, sink tops, tables, desks and other inanimate surfaces in hospitals, health care facilities, schools, motels and other institutional and commercial establishments. At this dilution SANI-WEX kills *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, and pathogenic fungi, *Trichophyton interdigitale*. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

*When used at a dilution of 3 1/2 ounces per 5 gallons of water on environmental inanimate surfaces, SANI-WEX is effective against Influenza A2, Herpes Simplex, Adenovirus Type 2, and Vaccinia Viruses.

For Food Processing Equipment sanitize previously cleaned equipment and food utensils by using 1 ounce of SANI-WEX per 4 gallons of water. This provides 200 ppm of active quaternaries. NO POTABLE WATER RINSE required at this level.

For Sanitizing previously cleaned Dairy Equipment such as tanks, lines and pails, use 1 ounce of SANI-WEX to 4 gallons of water (200 ppm). At this level NO POTABLE WATER RINSE is required. Follow recommendations of local Health Board.

For Restaurant and Bar Rinse — Dishes, glassware, silverware, cooking utensils, etc. should be thoroughly washed, rinsed and then immersed in a sanitizing solution containing 1 ounce of SANI-WEX per 4 gallons of water (200 ppm) NO TERMINAL POTABLE WATER RINSE REQUIRED

Cold Disinfection

To disinfect previously cleaned barber tools, and dental equipment into a solution containing 1 1/2 ounce per gallon of water for 10 minutes. At this dilution SANI-WEX is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, and pathogenic fungi, *Trichophyton interdigitale*, Herpes Simplex, Adenovirus Type 2, and

DANGER

Keep Out of Reach of Children. Causes eye and skin damage. Do not get on clothing. Protect eyes and skin from concentrate. Harmful if swallowed food.

**FOR INDUSTRIAL AND INSTITUTIONAL USE
 CONTENTS ONE U. S. GALLON**

ED
5
icide.
icide Act.
sticida
810-5

DIRECTIONS

Use SANI-WEX at a dilution of 3 ounces per 5 gallons of water for disinfecting floors, walls, sink tops, tables, desks and other inanimate surfaces in hospitals, health care facilities, schools, hotels and other institutional and commercial establishments. At this dilution SANI-WEX kills Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, and pathogenic fungi, Tricophyton interdigitale. For heavily soiled contaminated areas, a pre-cleaning step is recommended.

When used at a dilution of 3 1/2 ounces per 5 gallons of water on environmental inanimate surfaces, SANI-WEX is effective against Influenza A2, Herpes Simplex, Adenovirus Type 2, and Vaccinia Viruses.

For Food Processing Equipment sanitize previously cleaned equipment and food utensils by using 1 ounce of SANI-WEX per 4 gallons of water. This provides 200 ppm of active ingredients. NO POTABLE WATER RINSE required at this level.

For Sanitizing previously cleaned Dairy Equipment such as tanks, lines and pails, use 1 ounce of SANI-WEX to 4 gallons of water (200 ppm). At this level NO POTABLE WATER RINSE is required. Follow recommendations of local Health Board.

For Restaurant and Bar Rinse — Dishes, glassware, silverware, cooking utensils, etc. should be thoroughly washed, rinsed and then immersed in a sanitizing solution containing 1 ounce of SANI-WEX per 4 gallons of water (200 ppm) NO TERMINAL POTABLE WATER RINSE REQUIRED

SANI-WEX

Cold Disinfection

To disinfect previously cleaned surgical instruments, barber tools, and dental equipment, submerge instruments into a solution containing 1 1/2 ounces of SANI-WEX per gallon of water for 10 minutes. At this level SANI-WEX is effective against Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, pathogenic fungi, Tricophyton interdigitale, Influenza A2, Herpes Simplex, Adenovirus Type 2, and Vaccinia Viruses.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

Wexford Labs, Inc.

EPA ESTABLISHMENT NUMBER 34810-MO-1