

LESTER-30 is a highly effective product for controlling the growth of bacteria and fungi found in industrial recirculating cooling water towers, air washers, and evaporative condensers.

Industrial Recirculating Water Cooling Towers, Evaporative Condensers and Air Washer Systems.

Dosages for industrial recirculating water systems will depend on the condition of the system prior to treatment initiation. Systems which are heavily contaminated should be cleaned first. Apply **LESTER-30** to the cleaned system or when growth is first noticed, according to the following schedule:

Initial Dose: Apply 13.06 fluid ounces of product per 1,000 gallons of water in the system (120 ppm.). This dose may be repeated once, twice or three times weekly or as required to control the growth of slime-forming organisms. **LESTER-30** should be applied at a point where the product will be uniformly mixed.

Subsequent Dose: When microbial control is evident, add 2.18 to 4.36 fluid ounces of **LESTER-30** per 1,000 gallons of water (20-40 ppm.) in the system every three days or as needed. **LESTER ALGAE-CIDE 30** should be applied at a point where the product will be uniformly mixed.

LESTER

LESTER 30 BACTERICIDE AND FUNGICIDE

Active Ingredients
Sodium Dimethyldithiocarbamate
Nabam (Disodium Ethylene Bisdithiocarbamate)
Inert Ingredients

CONTENTS: Liquid — See Markings on Top of Drum
Do not re-use empty containers. Container should be decontaminated or crushed. Discard container in a safe place.

CAUTION KEEP OUT OF REACH OF CHILDREN
SEE OTHER PRECAUTIONS ON RIGHT PANEL.

SOLD BY:

LESTER LABORATORIES

INCORPORATED

ATLANTA, GEORGIA U.S.A.



CAUTION

KEEP OUT OF REACH OF CHILDREN

Causes eye and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

CAUTION: This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water. Apply this product only as specified on this label.

INSECTICIDE
TICIDE ACT
REGISTERED

58

WEIGHT: 9.7 Lb./Gal.
E.P.A. Registration No. 337-58



WEX-CIDE

CONCENTRATED GERMICIDAL DETERGENT
CLEANS ——— DISINFECTS ——— DEODORIZES
GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL
FUNGICIDAL and VIRUCIDAL*

in soft or hard water (A.O.A.C. 400 ppm Hardness as Calcium Carbonate)

ACTIVE INGREDIENTS:

ortho-Phenylphenol 6.73%, ortho-Benzyl-para-Chlorophenol 5.76%
Essential Oils 0.50%

INERT INGREDIENTS:

87.01% Includes biodegradable detergents and no phosphates.

EPA REGISTRATION NO. 34810-8

DANGER: KEEP OUT OF REACH OF CHILDREN.

See side panel for additional precautionary and first aid statements.

Wexford Labs, Inc.

1035 South Vandeventer

St. Louis, Missouri 63110

FOR INDUSTRIAL AND INSTITUTIONAL USE
CONTENTS ONE U. S. GALLON

WEX-CIDE

CONCENTRATED GERMICIDAL DETERGENT

..... 1:256

..... 1:256

..... 1:256

WEX-CIDE is used for cleaning, disinfecting and deodorizing floors, walls, and other similar hard surfaces. Especially suited for use in Operating rooms, Intensive Care, Nurseries, Emergency Areas, Patient Rooms and Rest Rooms.

When used as directed WEX-CIDE is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, pathogenic fungi, and Influenza A₂ virus.

DIRECTIONS

For all disinfection, including hospital and health care facilities, use WEX-CIDE at a dilution of ½ ounce per gallon of water. *When used at a dilution of ½ ounce per gallon of water on environmental inanimate surfaces, WEX-CIDE is effective against Influenza A₂ virus.

WEX-CIDE solution may be applied with a cloth, mop or brush, or by immersing equipment to be disinfected.

When disinfecting surgical instruments with WEX-CIDE, these instruments must be precleaned to remove blood and serious exudates prior to immersion in, or treatment with WEX-CIDE.

Do not reuse empty container. Wash thoroughly with water and detergent before discarding or returning drum to drum reconditioner.

WEX-CIDE is a synthetic germicidal detergent with broad spectrum kill of both gram positive and gram negative micro-organisms. It is an excellent cleaner.

Contains biodegradable detergents and is phosphate free. Do not mix with other cleaners.

DANGER — Keep out of reach of children. Corrosive. Causes eye damage or skin irritation. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID — In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol and call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Wexford Labs, Inc.

EPA ESTABLISHMENT NUMBER 34810-MO-1

ACCEPTED

SEP 9 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO 34810-8