EPA Form 8570-6 (Rev. 5-76)

SIGNATURE OF APPROVING OFFICIAL

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

DATE

ACCEPTED
with COMMENTS
in EPA Letter Dated;

DEC 0 7 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

NON-CHLOR Concentrated Germicidal Detergent

CLEANS -- DISINFECTS -- DEODORIZES
GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL, FUNGICIDAL and VIRUCIDAL*
IN SOFT WATER or HARD WATER (A.O.A.C. 400 ppm hardness as
Calcium Carbonate), with 5% Serum.

outerem carbonates, when an because	
ACTIVE INGREDIENTS: o-Phenylphenol	
Includes histograndship determines and water 9	4
Includes biodegradable detergents, perfume and water 8	J.4
EPA REG. NO	
DANCED. VEED OUT OF DEACH OF CUITADEN	
DANGER: KEEP OUT OF REACH OF CHILDREN	_
See rear panel for additional precautionary and first aid stateme	nts.
WEXFORD LABS, INC. 325 Leffingwell Avenue Kirkwood, Missouri 6	3122

FOR USE AND STORAGE BY MAINTENANCE PERSONNEL ONLY Contents: One U.S. Gallon

NON-CHLOR Concentrated Germicidal Detergent

NON-CHLOR is used for cleaning, disinfecting and deodorizing floors, walls, and other hard surfaces. Especially suited for use in Operating Rooms, Intensive Care, Nurseries, Emergency Areas, Patient Rooms and Rest Rooms. When used as directed NON-CHLOR is effective against gram negative and gram positive nicroorganisms such as Staphylococcus aureus, Salmonella choleraesuis, Pseudomonas aeruginosa, Pathogenic fungi, Mycobacterium tuberculosis and *Lipophilic viruses, such as Influenza A3 (H3N2-Hong Kong), Herpes Simplex Type 2, and Vaccinia, in the presence of A.O.A.C. 400 ppm hard water and 5% serum.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For all disinfection of hard surfaces in hospitals, health care facilities, schools and nursing homes, such as floors, walls, bathroom fixtures, shower stalls, and surgical instruments, use NON-CHLOR at 1 ounce per gallon of water. NON-CHLOR use-dilution may be applied with a cloth, mop or brush on surfaces such as stainless steel, chrome, glass, vinyl, ceramic, and terrazzo. Wet all surfaces thoroughly for at least ten minutes. Use-dilution of NON-CHLOR should be changed daily or sooner if it becomes over diluted or soiled. Cat 2010.

NON-CHLOR is effective against Influenza A3 Hong Kong, Herpex Simplex Type 2, and Vaccinia viruses in 400 ppm hard water and 5% organic soil for inanimate hard surfaces. NON-CHLOR is a synthetic germicidal detergent with broad spectrum kill of both gram positive and gram negative microorganisms. Contains biodegradable detergents and is phosphate free. DO NOT MIX WITH OTHER CLEANERS. For gross filth or heavy soil a pre-cleaning step is required.

DANGER KEEP OUT OF REACH OF CHILDREN. Corrosive, causes eye damage and skin irritation. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wear safety glasses when handling concentrate. Harmful if swallowed. Avoid contamination of food.

STORAGE and DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration, if allowed by state and local authorities. Consult federal, state or local disposal authorities for approved alternative procedure. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticides, spray mixture or rinsate is a violation of Federal Law.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol and call a physician.

NOTE TO PHYSICIAN Probable muscosal damage may

contraindicate the use of gastric lavage. Measures against circulator shock, respiratory depression and convulsion may be needed.