M32 S4810-01

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

NOV 6 1995

WEXFORD LABS, INC. 325 Leffingwell Avenue Kirkwood, MO 63122

Attention: Said I. Raziq

Director of Research

Subject:

Read to Use Wexcide

EPA Registration No. 34810-21

MOU Compliance Amendment dated July 31, 1995

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, will be acceptable provided that you make the labeling change listed below before you release the product for shipment bearing the amended label.

Remove surgical instruments from the list of surfaces to be disinfected. Surgical instruments are critical medical devices and must be sterilized.

A stamped copy is enclosed for your records. Submit five (5) copies of the final printed label prior to release of the product for shipment.

If you have any questions concerning this letter, please contact Wanda Mitchell at (703) 305-6141.

Sincerely,

Ruth G. Douglas Product Manager (32) Antimicrobial Program Branch Registration Division (7505C)

SURNAME	
SURNAME)	
.	
DATE	••••••

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

12 U.S. GOVERNMENT PRINTING OFFICE 1995 - 619-457



READY TO USE WEX-CIDE

GERMICIDAL DETERGENT
CLEANS • DISINFECTS • DEODORIZES
GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL,
FUNGICIDAL AND VIRUCIDAL*

 ACTIVE INGREDIENTS:
 0.026%

 c-Phenylphenol
 0.023%

 o-Benzyl-p-Chlorophenol
 0.023%

 INERT INGREDIENTS:
 99.951%

 includies biodegradable delergents and no phosphates
 T00.000%

EPA REG. NO. 34810-21 ● EPA EST. NO. 34810-MO-1 KEEP OUT OF REACH OF CHILDREN CAUTION

GERMICIDAL ACTIVITY: (AOAC Use Dilution Confirmation in the presence of 5% serum.) Ps. aeruginosa, S. aureus, S. choleraesuis; including S. aureus (MRSA) (TNSR) (GRM) and (PR) and S. Epidermidis (MRSE). TUBERCULOCIDAL ACTIVITY: (AOAC M. tuberculosis in the presence of 5% serum at 20°C for 10 minutes) FUNGICIDAL ACTIVITY: (AOAC T. Mentagrophytes in the presence of 5% serum.) VIRUCIDAL ACTIVITY: (*Influenza A3[Hong Kong], Adenovirus Type 2, Herpes simplex Types 1 & 2, Vaccinia virus. Human Rotavirus, Respiratory Syncytial virus, in the presence of 5% serum, and HIV-1 (virus associated with AIDS) In the presence of 10% serum on inanimate surfaces:

Use for all disinfection of herd, non-porous inanimate surfaces in Operating Rooms, Intensive Care, Nurseries, Emergency Areas, Dental Operatories, Police, EMS and other emergency response

STORAGE and DISPOSAL: Store at room temperature. Do not 'eave containers of this product open during storage. Do not reuse empty container. Rinse empty container thoroughly and discard in trash.



J Wexford Labs, Inc.

325 Leffingwell Avenue

Kirkwood, Missouri 63122

Contents: One Quart

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent woof hard, non-porous inanimate surfaces, surgical instruments, stainless steel, a cloth or brush, or by immersing equipment to be disinfected. All hard surfacements wet for at least 10 minutes before drying with a cloth. Gross filth and to applying cleaning solution. Discard solution when it becomes dirty and repulse Wex-Cide is a synthetic germicidal detergent with broad spectrum kill of microorganisms. It is an excellent cleaner. DO NOT MIX WITH OTHER CLEAD DECONTAMINATION OF CRITICAL OR SEMI-CRITICAL DEV STERILIZATION/HIGH LEVEL DISINFECTION:. This product is not to be used disinfectant on any surface or instrument that (1) is introduced directly into contact with the bloodstream or normally sterile areas of the body, or (2) contact with the bloodstream or normally sterile areas of the body, or (2) contact with the bloodstream or decontaminate critical or semi-critical morning level disinfection.

*READY TO USE WEX-CIDE KILLS HIV ON PRE-CLEANED ENVIRON PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in all settings in which soiling of inanimate surfaces/objects with blood or body fluids, and in which soiled with blood or body fluids can be associated with the potential for transmivirus type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SUBBLOOD AND BODY FLUIDS WHICH MAY CONTAIN HIV-1(AIDS virus), disposable rubber or latex gloves, gowns, face masks, or eye coverings as application of these surfaces. CONTACT TIME: Leave surfaces wet for 1 minus necessary to control other common types of viruses and bacteria. CLEANING be used to remove and clean blood and other body fluids from surfaces allightly soiled areas wipe off visual soil/clean with solution, and disinfect surfaces solution. For large spills the contaminated area should be flooded with decontaminated with fresh solution. DISPOSAL OF INFECTIOUS MATER should be autoclaved and disposed of according to federal, state, and loc disposal.

NOV r 1395

Under the Ferry constructed, empiride of the table Act as the posticide code ill A Reg. No.

18-018 hE