WC-RTU-122397

W

348

7/1/

7/1/98

Dode 1

ACCEPTED JUL 1 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 1

READY TO USE WEX-CIDE

GERMICIDAL DETERGENT
CLEANS • DISINFECTS • DEODORIZES
GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL,
FUNGICIDAL AND VIRUCIDAL*

ACTIVE INGREDIENTS:

O-Phenylphenol

O-Benzyl-p-Chlorophenol

INERT INGREDIENTS:

Includes blodegradable detergents and no phosphates

100.000

EPA REG. NO. 34810-21 ● EPA EST. NO. 34810-MO-1 KEEP OUT OF REACH OF CHILDREN CAUTION

GERMICIDAL ACTIVITY: (AOAC Use Difution Confirmation in the presence of 5% serum.) Ps. aeruginosa, S. aureus, S. choleraesuls; including S. aureus (MRSA) (TNSR) (3RM) and (PR) and S. Epidemidis (MRSE), Entercooccus Faecalis (VRE). TUBERCULOCIDAL ACTIVITY: (AOAC M. tuberculosis in the presence of 5% serum at 20° C for 10 minutes) FUNGICIDAL ACTIVITY: (AOAC T. Mentegrophytes in the presence of 5% serum,) VIRUCIDAL ACTIVITY: (*Influenza A3[Hong Kong), Adenoxinus Type 2, Herpes simplex Types 1 & 2, Vaccinta vi/us, Human Rotavirus, Respiratory Syncytial virus, in the presence of 5% serum, and HIV-1 (virus associated with AIDS) in the presence of 10% serum on inanimate surfaces.)

Use for all disinfection of hard, non-porous inanimate surfaces in Operating Rooms, intensive Care, Nurserfee, Emergency Areas, Dental Operatories, Police, EMS and other emergency response units

STORAGE and DISPOSAL: Store at room temperature. Do not leave containers of this product open during storage. Do not reuse empty container. Rinse empty container incroughly, and discard in trash.

Wexford Labs, Inc.

325 Leffingwell Avenue

Kirkwood, Missouri 63122

Contents: One Quart

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For all disinfection of hard, non-porous inanimate surfaces, stainless steel, chrome, glass and vinyl, apply with a cloth or brush, or by immersing equipment to be disinfected. All hard surfaces must be well thoroughly, and remain wet for at least 10 minutes before drying with a cloth. Gross fifth and heavy soil must be removed before applying cleaning solution. Discard solution when it becomes dirty and replace with fresh solution, Ready To Use Wex-Cide is a synthetic germioidal detergent with broad spectrum kill of gram positive and gram negative microorganisms, it is an excellent cleaner. DO NOT MIX WITH OTHER CLEANERS.

DECONTAMINATION OF CRITICAL OR SEMI-CRITICAL DEVICES PRIOR TO TERMINAL STERILIZATION/HIGH LEVEL DISINFECTION:. This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

*READY TO USE WEX-CIDE KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in all settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be solled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SURFACES/OBJECTS SOILED WITH BLOOD AND BODY FLUIDS WHICH MAY CONTAIN HIV-1 (AIDS virus). PERSONAL PROTECTION: Wear disposable rubber or latex gloves, gowns, face masks, or eye coverings as appropriate, when performing manual cleaning of these surfaces. CONTACT TIME: Leave surfaces wet for 1 minute. A contact time of 10 minutes is necessary to control other common types of viruses and bacteria. CLEANING PROCEDURE: The solution may be used to remove and clean blood and other body fluids from surfaces and objects before disinfection. For lightly soiled areas wipe off visual soil/clean with solution, and disinfect surface with a second application of the solution. For large spills the contaminated area should be flooded with the solution before cleaning, then decontaminated with fresh solution. DISPOSAL OF INFECTIOUS MATERIAL: Blood and other body fluids should be autoclaved and disposed of according to federal, state, and local regulations for infectious waste disposal.