

34810-35

10/02/2007

**ACCEPTED**  
 OCT - 2 2007

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. **34810-35**

113

**CLEAN-CIDE  
 READY TO USE  
 GERMICIDAL DETERGENT  
 CLEANS • DISINFECTS • DEODORIZES  
 GERMICIDAL, PSEUDOMONACIDAL, TUBERCULOCIDAL,  
 FUNGICIDAL and VIRUCIDAL\***

**ACTIVE INGREDIENT:**

Citric Acid .....	0.60%
<b>OTHER INGREDIENTS:</b> .....	99.40%
<b>TOTAL</b>	<b>100.00%</b>

EPA Reg. No. 34810-35    EPA Est. No. 34810-MO-1

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**GERMICIDAL:** effective in 5% serum against: *Pseudomonas aeruginosa*<sup>1</sup> [ATCC 15442], *Staphylococcus aureus*<sup>1</sup> [ATCC 6538], *Salmonella enterica*<sup>1</sup> (formerly *Salmonella Choleraesuis*) [ATCC 10708], *Listeria monocytogenes*<sup>1</sup> [ATCC 19117], *Escherichia Coli* O157:H7<sup>1</sup> [ATCC 43888], *Staphylococcus aureus* - MRSA<sup>1</sup> [ATCC 33592], *Staphylococcus epidermis* - MRSE<sup>1</sup> [ATCC 51625], *Enterococcus faecalis* Vancomycin Resistant<sup>1</sup> (VRE) [ATCC 51575], *Acinetobacter baumannii*<sup>2</sup> [19606] **VIRUCIDAL:** effective in 5% serum against: Poliovirus<sup>2</sup> type 1, Canine Parvovirus<sup>2</sup> (CPV), Respiratory Syncytial Virus<sup>1</sup>, Rotavirus<sup>1</sup>, Human Immunodeficiency Virus<sup>1</sup> Type 1 (Strain HTLV-III<sub>B</sub>), Herpes Simplex<sup>1</sup> Type 1, Herpes Simplex<sup>1</sup> Type 2 (Strain G), Vaccinia Virus<sup>1</sup>, Influenza A Virus<sup>1</sup> (Strain Hong Kong), Adenovirus<sup>1</sup> Type 2, Duck Hepatitis B Virus (DHBV)<sup>2</sup>, Feline Calicivirus<sup>1</sup> (surrogate for Norovirus) **FUNGICIDAL:** effective in 5% serum against *Trichophyton mentagrophytes*<sup>2</sup> [ATCC 9533]

1 - Five minute Contact Time    2 - Ten minute Contact Time

**TUBERCULOCIDAL:** effective against *Mycobacterium bovis* [ATCC 35743] in 5 minutes at 20°C / 68°F on previously cleaned hard non-porous inanimate surfaces.

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

**FIRST AID**

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a Poison Control Center or doctor for further treatment advice.

**Wexford Labs, Inc.** 325 Leffingwell Avenue Kirkwood, Missouri 63122  
Net Contents: One Quart

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Product is ready to use. **DO NOT DILUTE.**

**TO CLEAN:** Apply Clean-Cide undiluted to hard, non-porous surfaces by wiping with a cloth or towel until clean.

**TO DISINFECT:** Apply the Clean-Cide product undiluted to hard, non-porous surfaces with a cloth, mop or brush, or by immersing equipment to be disinfected. Wet all surfaces thoroughly. The surface must remain wet with Clean-Cide for at least 10 minutes. Wipe or let air dry. Preclean areas with gross filth and heavy soil prior to disinfection.

**[AREAS OF USE:]**

- [Hospitals] [Nursing homes]
- [Health Care Facilities] [Bathrooms]
- [Medical Offices] [Institutions]
- [Dental offices] [Institutional facilities]
- [Day care centers] [Kennels]
- [Nurseries] [Grooming Facilities]
- [Schools] [Veterinary Clinics]
- [Laboratories or other small animal facilities]

**[TYPES OF SURFACES:]**

- [On washable hard, nonporous surfaces of:]
- [Bathroom fixtures] [Stainless Steel]
- [Floors] [Chrome]
- [Walls] [Glass]
- [Vinyl]

**DECONTAMINATION OF CRITICAL OR SEMI-CRITICAL DEVICES PRIOR TO TERMINAL STERILIZATION/HIGH LEVEL DISINFECTION:** This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

**\*CLEAN-CIDE KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in all settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SURFACES/OBJECTS SOILED WITH BLOOD AND BODY FLUIDS WHICH MAY CONTAIN HIV-1 (AIDS virus). PERSONAL PROTECTION:** When handling objects soiled with blood or body fluids use disposable latex gloves, gowns, mask or eye coverings. **CLEANING PROCEDURE: DO NOT DILUTE** this product. Use as is. Remove and clean blood and other body fluids from surfaces and objects before application of Clean-Cide with a mop, cloth, or sponge so as to thoroughly wet pre-cleaned surfaces. Allow surfaces to remain wet for five minutes to ensure total in-activation of HIV virus. For lightly soiled areas wipe off visual soil, clean with solution, and disinfect surface with second application of solution. For large spills, the contaminated area should be flooded with the solution before cleaning then decontaminated with fresh solution.

**Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shop / Kennels**

**Disinfection Directions:** For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with Clean-Cide for a period of 10 minutes. Wipe or allow to air dry. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

**DISPOSAL OF INFECTIOUS MATERIAL:** Blood and other fluids should be autoclaved and disposed of according to Federal, State and Local regulations for infectious waste disposal.

**STORAGE AND DISPOSAL:** Store in a cool place. Do not leave containers open during storage. Do not reuse empty container. Wastes resulting from use of this product may be poured down the drain. No special handling is required. Federal, State, or local regulations should be observed.

**Made in the USA**

MSDS #: 2130-02

Reorder #: 2130-02