

34810-36

10/02/2007

**ACCEPTED**  
 OCT - 2 2007  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. **34810-36**

1/3

**CLEAN-CIDE WIPES**  
**Cleans, Disinfects, Deodorizes**  
**Germicidal, Fungicidal, and Virucidal**

Active Ingredient: Citric Acid..... 0.6%  
 Inert Ingredients: ..... 99.4%  
 Includes biodegradable detergents  
 Total: ..... 100.0%

EPA Reg. No. 34810-36

EPA Est. No. 34810-MO-1

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

**Effective in 5% serum against:**

**GERMICIDAL:** *Pseudomonas aeruginosa*<sup>1</sup> [ATCC 15442], *Staphylococcus aureus*<sup>1</sup> [ATCC 6538], *Salmonella enterica*<sup>1</sup> (formerly *Salmonella Choleraesuis*) [ATCC 10708], *Listeria monocytogenes*<sup>1</sup> [ATCC 19117], *Escherichia Coli O157:H7*<sup>1</sup> [ATCC 43888], *Staphylococcus aureus - MRSA*<sup>1</sup> [ATCC 33592], *Staphylococcus epidermis - MRSE*<sup>1</sup> [ATCC 51625], *Enterococcus faecalis Vancomycin Resistant*<sup>1</sup> (VRE) [ATCC 51575], *Acinetobacter baumannii*<sup>2</sup> [ATCC 19606] **VIRUCIDAL:** Poliovirus<sup>2</sup> type 1, Canine Parvovirus<sup>2</sup> (CPV), Respiratory Syncytial Virus<sup>1</sup>, Rotavirus<sup>1</sup>, Human Immunodeficiency Virus<sup>1</sup> Type 1 (Strain HTLV-III<sub>B</sub>), Herpes Simplex<sup>1</sup> Type 1, Herpes Simplex<sup>1</sup> Type 2 (Strain G), Vaccinia Virus<sup>1</sup>, Influenza A Virus<sup>1</sup> (Strain Hong Kong), Adenovirus<sup>1</sup> Type 2, Duck Hepatitis B Virus (DHBV)<sup>2</sup>, Feline Calicivirus<sup>1</sup> (surrogate for Norovirus) **FUNGICIDAL:** *Trichophyton mentagrophytes*<sup>2</sup> [ATCC 9533]

- 1 - Five minutes Contact Time
- 2 - Ten minutes Contact Time

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

**FIRST AID**

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a Poison Control Center or doctor for further treatment advice.

**Wexford Labs, Inc.** 325 Leffingwell Avenue Kirkwood, Missouri 63122

**NET CONTENTS:**

X Pre-measured Towelettes (X)" x (X)"  
 (e.g. 100 Pre-measured Towelettes 7"x 8")

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Product is ready to use.

**DISPENSER DIRECTIONS:** Wear gloves. To start feed, remove large cover. Pull up the towelette corner from the center of the towelette roll, twist it into a point and thread it through the hole in the container cover. Pull through about one inch. Replace cover. Pull out first towelette and snap off at a 90° angle. Remaining towelettes feed automatically ready for their next use. When through using, keep small center cap closed to prevent moisture loss.

**TO CLEAN:** Unfold premoistened cloth and wipe area to be cleaned for 30 seconds or until clean.

**TO DISINFECT:** Unfold premoistened cloth and wipe area to be cleaned and disinfected. Use enough wipes for treated surface to remain visibly wet for 10 minutes. Wipe or let air dry. For gross filth or heavy soil, use a wipe to clean and then a second wipe to disinfect the surface. **DO NOT MIX WITH OTHER CLEANERS.**

**[AREAS OF USE:]**

[Hospitals]	[Bathrooms]
[Health Care Facilities]	[Institutions]
[Medical Offices]	[Institutional facilities]
[Dental offices]	[Veterinary Clinics]
[Day care centers]	[Kennels]
[Nurseries]	[Grooming Facilities]
[Schools]	[Zoos]
[Nursing homes]	[Pet Shops]
[Laboratories or other small animal facilities]	

**[TYPES OF SURFACES:]**

[On washable hard, nonporous surfaces of:]

[Bathroom fixtures]	[Stainless Steel]
[Floors]	[Chrome]
[Walls]	[Glass]
[Vinyl]	

**DECONTAMINATION OF CRITICAL OR SEMI-CRITICAL DEVICES PRIOR TO TERMINAL STERILIZATION/HIGH LEVEL DISINFECTION:**

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

**\*CLEAN-CIDE WIPES KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in all settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS).

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SURFACES/OBJECTS SOILED WITH BLOOD AND BODY FLUIDS WHICH MAY CONTAIN HIV-1 (AIDS virus):**

**Personal Protection:** When handling objects soiled with blood or body fluids use disposable latex gloves, gowns, mask, or eye coverings.

**Cleaning Procedure:** All blood and body fluids must be cleaned from surfaces before disinfection by the germicidal cloth. Use one germicidal cloth to remove gross soil.

**Contact Time:** Use a second germicidal cloth to thoroughly wet surface. Allow to remain wet 5 minutes, let air dry.

**Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shop / Kennels Disinfection**

**Directions:** For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent. Empty rinse with water. Unfold premoistened cloth and wipe area to be cleaned and disinfected. Use enough wipes for treated surface to remain visibly wet for 10 minutes. Wipe or allow to air dry. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

**Disposal of Infectious Material:** Dispose of used towelette in accordance with local regulations for infectious waste disposal.

**STORAGE AND DISPOSAL:** Store in a cool place. Do not leave containers open during storage. Do not reuse empty container. No special handling is required. Dispose of used towelette in accordance with Federal, State, or local regulations for infectious waste disposal.

**Made in the USA**

MSDS #: 3130-XX

Reorder #: 3130-XX