



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, DC 20460

APR - 6 2010

OFFICE OF
PREVENTION, PESTICIDE
AND TOXIC SUBSTANCES

Mary Anne Auer Wexford Labs, Inc. 325 Leffingwell Avenue Kirkwood, MO 63122

Subject:

CleanCide Wipes

EPA Registration No. 34810-36 Notification Date: March 12, 2010 EPA Receipt Date: March 15, 2010

Dear Ms. Auer:

This letter acknowledges receipt of your notification submitted under the provision of PR Notice 98-10.

Proposed Notification

• Adding new surface (CuVerro Antimicrobial Copper Surface)

General Comment:

Based on a review of the submitted materials, your notification for the addition of a new surface is acceptable and a part of the records on file:

Should you have any questions concerning this letter, please contact Heather Garvie at (703) 308-0034 or by email address at garvie.heather@epa.gov.

Sincerely.

acqueline McFarlane

(Acting) Product Manager (34) Regulatory Management Branch II Antimicrobials Division (7510P) March 12, 2010

Ms. Jacqueline McFarlane
(Acting) Product Manager (34)
Regulatory Management Branch II
United States Environmental Protection Agency
Antimicrobial Division (7510P)
2777 Crystal Drive
Potomac Yard 1
Arlington, VA 22202

Re: Letter Dated: February 2, 2010

Label revision for CleanCide Wipes (EPA # 34810-36)

Dear Ms. McFarlane:

I have enclosed the following:

- Copy of your letter dated February 2, 2010
 - Three copies of our CleanCide Wipes (EPA # 34810-36) label revised per your letter dated February 2, 2010
 - Form 8570-1 Notification for CleanCide Wipes (EPA # 34810-36)

We appreciate your assistance with this process.

Anne auer

Sincerely,

WEXFORD LABS, INC.

Mary Anne Auer

CEO

Enclosures

CLEAN-CIDE WIPES Cleans, Disinfects, Deodorizes Germicidal, Fungicidal, and Virucidal*

NOTIFICATION Date Reviewed: 310.

Active Ingredient: Citric Acid	0.6%
Inert Ingredients:	
Includes biodegradable detergents	
Total:	100.0%

EPA Reg. No. 34810-36

EPA Est. No. 34810-MO-1

KEEP OUT OF REACH OF CHILDREN CAUTION

Effective in 5% serum against:

GERMICIDAL: Pseudomonas aeruginosa¹ [ATCC 15442], Staphylococcus aureus¹ [ATCC 6538], Salmonella enterica¹ (formerly Salmonella Choleraesuis) [ATCC 10708], Listeria monocytogenes¹ [ATCC 19117], Escherichia Coli O157:H7¹ [ATCC 43888], Staphylococcus aureus - MRSA¹ [ATCC 33592], Staphylococcus epidermis - MRSE¹ [ATCC 51625], Enterococcus faecalis Vancomycin Resistant¹ (VRE) [ATCC 51575], Acinetobacter baumannii² [ATCC 19606] VIRUCIDAL: Poliovirus² type 1, Canine Parvovirus² (CPV), Respiratory Syncytial Virus¹, Rotavirus¹, Human Immunodeficiency Virus¹ Type 1 (Strain HTLV-III_B), Herpes Simplex¹ Type 1, Herpes Simplex¹ Type 2 (Strain G), Vaccinia Virus¹, Influenza A Virus¹ (Strain Hong Kong), Adenovirus¹ Type 2, Duck Hepatitis B Virus (DHBV)², Hepatitis C (HCV)² (utilizing Bovine Viral Diarrhea Virus as a surrogate for Human Hepatitis C), Feline Calicivirus¹ (surrogate for Norovirus) FUNGICIDAL: Trichophyton mentagrophytes² [ATCC 9533]

- 1 Five minutes Contact Time
- 2 Ten minutes Contact Time

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a Poison Control Center or doctor for further treatment advice.

Wexford Labs, Inc. 325 Leffingwell Avenue Kirkwood, Missouri 63122

NET CONTENTS: X lbs, X oz.

X Pre-moistened Towelettes (X)" x (X)" (e.g. 100 Pre-moistened Towelettes 7"x 8")

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Product is ready to use.

DISPENSER DIRECTIONS: Wear gloves. To start feed, remove large cover. Pull up the towelette corner from the center of the towelette roll, twist it into a point and thread it through the hole in the container cover. Pull through about one inch. Replace cover. Pull out first towelette and snap off at a 90° angle. Remaining towelettes feed automatically ready for their next use. When through using, keep small center cap closed to prevent moisture loss. **TO CLEAN/DEODORIZE:** Unfold premoistened cloth and wipe area to be cleaned for 30 seconds or until clean. **TO DISINFECT:** Unfold premoistened cloth and wipe area to be cleaned and disinfected. Use enough wipes for treated surface to remain visibly wet for 10 minutes. Wipe or let air dry. For gross filth or heavy soil, use a wipe to clean and then a second wipe to disinfect the surface. DO NOT MIX WITH OTHER CLEANERS.

[AREAS OF USE:]

[Hospitals]

[Bathrooms]

[Health Care Facilities]

[Institutions]

[Medical Offices]
[Dental offices]

[Institutional facilities] [Veterinary Clinics]

[Day care centers]

[Kennels]

[Nurseries]

[Grooming Facilities]

[Schools]
[Nursing homes]

[Zoos] [Pet Shops]

[Laboratories or other small animal facilities]

[TYPES OF SURFACES:]

[On washable hard, nonporous surfaces of:]

[Bathroom fixtures]

[Stainless Steel]

[Floors]

[Chrome]

[Walls]

[CuVerro_{TM} Antimicrobial Copper Surfaces]

[Vinyl]

[Glass]

DECONTAMINATION OF CRITICAL OR SEMI-CRITICAL DEVICES PRIOR TO TERMINAL STERILIZATION/HIGH LEVEL DISINFECTION:

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or, (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

*CLEAN-CIDE WIPES KILLS HBV, HCV and HIV ON PRE-CLEANED ENVIRONMENTAL

SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in all settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Hepatitis B Virus, Hepatitis C Virus and Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING SURFACES/OBJECTS SOILED WITH BLOOD AND BODY FLUIDS WHICH MAY CONTAIN HBV, HCV AND HIV-1 (AIDS virus):

<u>Personal Protection</u>: When handling objects soiled with blood or body fluids use disposable latex gloves, gowns, mask, and eye coverings.

<u>Cleaning Procedure</u>: All blood and body fluids must be cleaned from surfaces before disinfection by the germicidal cloth. Use one germicidal cloth to remove gross soil.

Contact Time: Use a second germicidal cloth to thoroughly wet surface. Allow to remain wet 5 minutes for HIV-1, 10 minutes for HBV and HCV, let air dry.

Disposal of Infectious Material: Dispose of used towelette in accordance with local regulations for infectious waste disposal.



Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shop / Kennels Disinfection Directions:

For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Unfold premoistened cloth and wipe area to be cleaned and disinfected. Use enough wipes for treated surface to remain visibly wet for 10 minutes. Wipe or allow to air dry. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

STORAGE AND DISPOSAL: Store in a cool place. Do not leave containers open during storage. Do not reuse empty container. No special handling is required. Dispose of used towelette in accordance with Federal, State, or local regulations for infectious waste disposal.

Made in the USA

MSDS #: 3130-XX

Reorder #: 3130-XX

