

ACCEPTED

Net Contents      Pounds

JUN 17 1975

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 34858-4

## ALLADDIN ALGICIDE PELLETS

### ACTIVE INGREDIENTS:

Sodium Pentachlorophenate..... 79%

Sodium Salts of Other Chlorophenols ..... 11%

INERT INGREDIENTS: ..... 10%

### **DANGER . . .** **KEEP OUT OF REACH OF CHILDREN**

Corrosive. Causes eye damage and skin irritation. Do not get dust in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. (See First Aid and Additional Precautions on Back Panel).

This product is toxic to fish and wildlife. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Apply this product only as specified on this label. Do not reuse empty drum. Destroy when empty.

### DIRECTIONS

ALLADDIN ALGICIDE PELLETS are used to control algae and fungal and bacterial slime in recirculating industrial cooling waters. Badly fouled systems must be cleaned before treatment is begun. Begin treatment when the system is in danger of becoming affected or after cleaning systems whose efficiency is already impaired. An initial slug of 6-12 ounces of ALLADDIN ALGICIDE PELLETS per 1000 gallons of water to provide 40-80 parts per million of active ingredients based on the total weight of water in the system is recommended. Repeat until control is achieved.

Subsequent slug additions of 3-6 ounces of ALLADDIN ALGICIDE PELLETS per 1000 gallons of water (20-40 parts per million of active ingredients) should be employed every 2-5 days or as needed. Frequency of addition depends upon the severity of the microbiological problem and the relative amount of bleed-off. Slug additions should be made in the sump of water cooling towers.

Sold by

## **ALLADDIN LABORATORIES, INC.**

30 SOUTH OCEAN AVENUE  
FREEPORT, N.Y. 11520

---

---

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before re-use.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

---

---