

EPA-PR

RUSSALL GENERAL PURPOSE AQUEOUS INSECTICIDE

A concentrated PRYRENONE® Mico Emulsion
Clear - Stable - Leaves No Oily or Objectionable
Residues

Effective, Fast Acting Insecticide

For use in Homes, Institutions, Restaurants
Food Processing Plants, Dairies, Livestock and
Outdoors for Mosquito Control

Apply with Conventional Mechanical or Compressed Air Equipment

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid contamination of food and foodstuffs.

Remove pets, birds and cover fish aquaria before spraying. This product is toxic to fish. Keep out of lakes, ponds or streams. Do not contaminate water by leaching of equipment or disposal of wastes. Apply this product only as directed on the label.

TORAGE AND DISPOSAL Store only in tightly closed, original container in a secure area inaccessible to children. Do not contaminate water, food or feed by storage or disposal. Waste resulting from the use of this product may be disposed on site or at an approved waste disposal facility. Do not reuse empty container. Triple-wash empty container with water. Return metal drums to reconditioner or hauler and dispose of in a sanitary landfill or by other procedures proved by state and local authorities. Plastic containers may be disposed of in a sanitary landfill incinerated or crushed by local authorities, by burning. If burned, stay out smoke.

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxid, Technical	1.0%

INERT INGREDIENTS:

	98.9%
	100.0%

*Equivalent to 0.8% *butylcarbitol (6-propylpiperonyl) ether and 0.2% related compounds.

PYRENONE - Registered trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side panel for additional precautionary statements

EPA REGISTRATION NO.

EPA ESTABLISHMENT NO.

DO NOT TRANSPORT OR STORE UNDER 32°

34892-16

34892 N.Y.C.

NET CONTENTS

MANUFACTURED BY:
RUSSALL PRODUCTS CO., INC.
2253 LIGHT ST., BRONX, N.Y. 10466

BEST AVAILABLE COPY

FEB 20 1985

34892-16

"It is a violation of Federal law to use this product inconsistent with its labeling."

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control fleas, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, root and flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, meal worms, grain mites, and codling: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray baseboards for ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horse flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies, apply at a rate of 2 ounces per adult cow or horse sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking ticks, apply to the infected areas of the animal using a stiff brush to get the spray to the base of the hair. Repeat every 2 or 3 weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools or ponds and ...ve trees. Apply at a rate of 1½ gallons per acre (4½ ounces per 1000 square feet). Using truck-mounted conventional misting equipment, such as Rotational® models 51C, 91 or 100 HT, traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 4.5 gallons per minute.