

421853

JUL 10 1992

Hi-Yield Chemical Company
Box 460
Bonham, TX 75418

Gentlemen:

Subject: Amendment - Revise per Diazinon Registration Standard
Hi-Yield 5% Diazinon Insect Killer
EPA Reg. No. 34911-13
Your submission received February 14, 1990

The amendment referred to above, submitted in connection with a registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided the following revisions are made prior to release for shipment under the amended labeling:

- 1) The Diazinon Registration Standard stipulates that diazinon outdoor use products are classified as Restricted Use due to avian and aquatic toxicity unless they are registered for application by homeowners to the home yard and/or home garden or by commercial applicators only to home yards. You can clarify the intended use site for this product by addition of "For outdoor use on the home lawn and in the home garden only" to the front panel. A good location for this addition would be just under the golf course/sod farm prohibition.
- 2) As you are aware, there are a number of label elements with type size minimums. Please be certain that your finished labels meet all type size requirements.
- 3) Under the heading "IN GARDENS DIAZINON KILLS", delete the pests "Armyworms", "Carrot rust flies", "Corn borers", "Rootworms" since the application methods required to control these pests, in-furrow or foliar broadcast, are not being supported for reregistration of granular diazinon products and directions for control of these pests will not appear on the revised label.

- 4) In the "Precautions" under the heading "Overall Lawn Application", revise the sixth sentence to read as follows:

Where thorough watering is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected to enter the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after applying this product, however, stop watering before puddling occurs.

- 5) The spreader setting table should be relocated to appear at the end of the overall lawn application directions, i.e., just after the directions for control of Hyperodes weevil. Delete the list of insect pests that currently appears beneath the spreader settings table. In its new location this table will be near the listing of lawn pests controlled by overall lawn application of the product.

Revise the heading for the spreader setting table to read:

LAWN SPREADER SETTING NO. (To deliver 2 lbs. per 1000 sq. ft)

<u>Spreader Make and Model</u>	<u>Setting</u>
--------------------------------	----------------

- 6) Revise the heading "Spot Treatment" to read "Lawn Spot Treatment".
- 7) Revise the first sentence of the directions for aiding in control of fire ants to read as follows to clearly indicate the home-use nature of this product:

To aid in the control of fire ants in home lawns and other outdoor home recreational areas, apply 1/2 cups over, and 2 ft. around, each fire ant mound.

- 8) Under the heading "Vegetables Insect Control" replace the second paragraph with the following:

Apply to the soil at the rate shown for the type of insect pest you want to control and immediately mix thoroughly into the soil to the depth specified. Do not apply this product to food crops grown in greenhouse.

- 9) For the crop "Beans", delete the directions for control of "Lesser Cornstalk Borers". Ciba-Geigy is not supporting reregistration of granular diazinon applied post-emergence to food crops because of the hazards to birds from exposed granules.

- 10) For the crop "Carrots", delete the directions for control of "Carrot Rust Flies". Ciba-Geigy is not supporting registration of in-furrow application of granular diazinon.
- 11) Revise the variety listing for melons to read as follows:

MELONS (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, and Hybrids of these) and WATERMELONS
- 12) For the crop "Onions", delete "by disking, harrowing, or other suitable means". This instruction was a carryover from an agricultural product label and is not appropriate to a homeowner-use product.
- 13) For the crop "Potatoes", delete the "Note" about hand-harvested commercially-grown potatoes, since this is not appropriate to a product whose crop uses are limited to application to the home garden.
- 14) For the crop "Radishes", delete the directions for control of "Root Maggots". Ciba-Geigy is not supporting reregistration of in-furrow application of granular diazinon.
- 15) For the crop "Sweet Corn" delete the directions for use for control of "European Corn Borers", "Fall Armyworms", and "Southwestern Corn Borers". Ciba-Geigy is not supporting reregistration of granular diazinon applied post-emergence to food crops because of the hazards to birds from exposed granules.
- 16) For the crop "Sweet Potatoes", delete the special directions for post-emergence use in the Southeastern U. S.. Ciba-Geigy is not supporting reregistration of granular diazinon applied post-emergence to food crops because of the hazards to birds from exposed granules.
- 17) The "Hazards to Humans and Domestic Animals:" text should be revised to read as follows:

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or

measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

- 18) Based on Ciba-Geigy's toxicology data submitted in support of reregistration of their 5% diazinon granular product, and upon which you are relying for reregistration of the subject product, in the Statement of Practical Treatment add "Get medical attention." to the end of the "If on skin" advice.

A stamped copy of the label is enclosed for your records. Please submit five copies of the revised amended label prior to release for shipment under the amended label. Labeling released for shipment after the deadline imposed by the Diazinon Registration Standard, as amended, must be the revised amended label.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (H7505C)

5/21

PM 13 Reg. No. 34911-13

Hi-Yield® 5% Diazinon® Insect Killer TREATS UP TO 5000 SQUARE FEET
Granules
lawn & garden

This product must not be used
on golf courses and sod farms.

5% Diazinon® Insect Killer Granules Kills

- | | |
|--------------------|-------------------|
| Ants | Earwigs |
| Armyworms | Fire ants* |
| Bermudagrass mites | Fleas |
| Billbugs | Onion maggots |
| Brown dog ticks | Root maggots |
| Chiggers | Sod webworms |
| Chinch bugs | White grubs of: |
| Crickets | . Japanese beetle |
| Cutworms | . European chafer |
| | . Southern chafer |
| | Wireworms |
| | and others |

*Aids in control.

See Directions for Use.

Ready-to-use Granules. Contains
5% diazinon insecticide.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	5%
Inert Ingredients:	95%
Total:	100%

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements
on back of bag.

EPA Reg. No. 34911-13

EPA Est. 7401-TX-1

Ten Pounds
Net Weight

ACCEPTED
with COMMENTS
In EPA Letter Dated

JUL 10 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
34911-13

INSTRUCTIONS FOR THE USE AND WARRANTY OF ALL ANS

IMPORTANT Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Hi-Yield Co. or the Seller. All such risks shall be assumed by the Buyer.

✓ Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Hi-Yield Chemical Company warrants this product to conform to the chemical description on the label and for the purpose stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

In Lawns
5% Diazinon® Kills

In Gardens
5% Diazinon® Kills

- Armyworms
- Bermudagrass mites
- Billbugs
- Brown dog ticks
- Chiggers
- Chinch bugs
- Clover mites
- Crickets
- Cutworms
- Digger wasps
- Earwigs
- Fleas
- Fire ants*
- Lawn moths
- Leafhoppers
- Millipedes
- Sod webworms
- Sowbugs
- Springtails
- White grubs of:
 - . European chafer
 - . Japanese beetle
 - . Southern chafer

- Carrot root flies
- Corn borers
- Cutworms
- Mole crickets
- Onion maggots
- Root maggots
- Rootworms
- Wireworms

*Aids in control.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Overall Lawn Application:

5% Diazinon® controls the following lawn insects on both grass and dichondra lawns:

- Ants . Armyworms . Bermudagrass mites . Brown dog ticks . .
- Chiggers . Chinch bugs . Clover mites . Crickets . Cutworms
- Earwigs . Fleas . Lawn moths (sod webworms) . Leafhoppers .
- Millipedes . Sowbugs . Springtails (Collembola)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds. Avoid overlapping treated areas. If waterfowl, i.e., ducks, geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 0.25 inch water immediately after application, however, stop watering before puddling occurs.

Apply 5% Diazinon[®] with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry.

For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10').

Billbugs; White grubs of Japanese beetle, European chafer, Southern chafer, Ataenius spretulus (dung beetle):

Apply 2 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late Summer or early Fall if a second generation occurs.

Hyperodes weevils:

Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Spot Treatment

Ants: Apply 1 1/2 teaspoons of 5% Diazinon[®] over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.

Digger wasps: Apply 1 1/2 teaspoons of 5% Diazinon[®] in and around nest opening. Apply at dusk when wasps are not active.

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses.

- | | |
|-----------------|--------------------|
| Ants | Fleas |
| Brown dog ticks | Millipedes |
| Clover mites | Sowbugs (Pillbugs) |
| Crickets | Springtails |
| Earwigs | (Collembola) |

Apply a 5 foot band of 5% Diazinon[®] next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. Repeat as necessary.

Vegetables - Insect Control

5% Diazinon[®] controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Unless directions under "Timing/Special Directions" state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

<u>Insect</u>	<u>Depth of Mixing or Incorporation into Soil</u>
Surface cutworms*	2-3 inches
Subterranean cutworms*	3-6 inches
Mole crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

*If type of cutworm cannot be identified, mix into soil 4-6 inches deep.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
(Pinto, Red or Lima) (succulent only)			
	Lesser Cornstalk Borers	3 1/2-7 ozs. per 500 sq. ft.	Apply in a 6-inch band centered over the row when plants are 1/2-1 inch tall.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc.		
Beets, Red (Table)	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Broccoli	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

	INSECT	RATE	TIMING/SPECIAL DIRECTIONS
(Cont'd)	Crickets	sq. ft.	days before planting
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Brussels Sprouts	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Cabbage	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

			TIMING/SPECIAL DIRECTION
		sq. ft.	Before planting.
	Carrot Rust Flies	7 ozs. per 500 sq. ft.	Apply in the seed furrow at planting time.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Broccoli	Root Maggots	7-10 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Peas	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

	INSECT	RATE	TIMING/SPECIAL DIRECTIONS
		sq. ft.	before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Cucumber	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Endive (Escarole)	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
ale	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

		sq. ft.	Before planting
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, & Hybrids of these and watermelons)	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Mustard	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Onions (green and bulb)	Onion Maggots	7-14 ozs. per 500 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil by disking, harrowing, or other suitable means. NOTE: Diazinon may not control organophosphate-resistant onion maggots.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
		sq. ft.	before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Peas (succulent only)	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
NOTE: Do not use on dried pea varieties such as dried blackeyed peas.			
Peppers	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Potatoes	Cutworms (Surface & Subterranean)	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
NOTE: Do not use on commercially-grown potatoes which will be hand-harvested.			

	Pests	RATE	TIMING/SPECIAL INSTRUCTIONS
Spinach	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Apply at time of planting (2 1/2-4 5/8 ozs. per 1,000 linear feet of row). Broadcast and mix into soil 1-2 days before planting.
Spinach	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Spinach	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Squash, Summer and Winter	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Squash, Summer and Winter	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Sweet Corn	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Sweet Corn	Seed Corn Maggots	7-14 ozs. per 500 sq. ft.	Broadcast just before planting. Mix into soil 2 inches.
Sweet Corn	Wireworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
(continued)	Borers		<p>Apply over the corn so granules fall into whorls when first generation borers begin feeding.</p> <p>Make a second application 7 days later if needed.*</p> <p>For second-generation borers, apply as above when there is an average of 1 egg mass per plant.</p> <p>Do not make more than 3 applications per season.*</p>
	Fall Armyworms	3 1/2-7 ozs. per 500 sq. ft.	Apply over corn so granules fall into whorls when larvae first appear. If necessary, application may be repeated twice at 7-day interval. Do not make more than 3 applications per season.*
	South-western Corn Borers	3 1/2-7 ozs. per 500 sq. ft.	Apply over corn so granules fall into whorls when second-generation larvae first appear. If necessary, application may be repeated at 7-day intervals. Do not make more than 3 applications per season.*

*NOTES: Corn may be picked immediately after last application. Do not feed fodder to livestock for 10 days following last application.

Sweet Potatoes	Wireworms, Flea Beetles	10 1/2 ozs. per 500 sq. ft.	<p>Broadcast and mix into soil just before planting.</p> <p>Southeastern U.S. - Broadcast 10 1/2 ozs. per 500 sq. ft. over the tops of vines when roots begin to enlarge and do not mix into soil.</p>
----------------	-------------------------	-----------------------------	--

	PEST	RATE	TIMING/COMMENTS
Chard	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Tomatoes	Cutworms	7-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Turnips	Mole Crickets	3 1/2 ozs. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10 1/2-14 ozs. per 500 sq. ft.	Broadcast and mix into soil just before planting.

SPREADER SETTING NO
(per 1000 sq ft)

Name and Model

Grass or Dichondra Lawl.
2 lbs.

ferti-lome
PLUS IRON
BROADCAST

16

ferti-lome
PLUS IRON
PROF. BROADCAST

16

ferti-lome
PLUS IRON
DROP

3

Scotts
MODEL PF-2
DROP

3

ORTHO
BROADCAST

3

Ants
Armyworms
Bermudagrass Mites
Brown Dog Ticks
Chiggers
Chinch Bugs

Clover Mites
Crickets
Cutworms
Earwigs
Fleas
Hyperodes Weevil

Lawn Moths
(Sod Webworms)
Leafhoppers
Millipedes
Sowbugs
(Pill Bugs)
Springtails
(Collembola)

Corrected Page for D-2-a^(B) 6000 Lawn & Garden
Insect Control Label (5% G)
-16-

20/21

Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. May cause contact sensitization following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

Updated
Statements

Statement of Practical Treatment

If swallowed: call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

Corrected
Area

If inhaled: remove victim to fresh air and apply artificial respiration, if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Continued

Note to Physician

... symptoms ...
... by injection ...
... be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wild-life. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 0.25 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries.

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

DIAZINON[®] registered trademark of CIBA-GEIGY CORP.

Hi-Yield Chemical Co.
EPA Est. No. 7401-TX-1

Bonham, Tx. 75418
EPA Reg. No. 34911-13

NET WT. 10 LBS.