

WHITE - UNIVERSAL

CAUTION
STATEMENT OF PRACTICAL TREATMENT
 If swallowed: Get medical attention.
 If in eyes: Wash with soap and water.
 If on skin: Wash with water for 15 minutes. Contact a physician.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION
 Harmful if swallowed. Avoid breathing dust. Avoid contact with skin. Eyes and skin should be washed thoroughly after handling and before eating or drinking. Remove and wash contaminated clothing before re use.
ENVIRONMENTAL HAZARDS
 This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from treated area. Do not contaminate water by cleaning of equipment or disposal of wastes. Use of this product only as specified on this label.
DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with the label.
 1. Dust lightly at the rate of 2 ounces per 50 feet of row on first appearance of insect or insect damage.
 2. Repeat at 4 to 7 days intervals under control.
 3. Frequent light dustings are more effective than single heavy dustings.
 4. Do not apply to food crops within 3 days of harvest.
VEGETABLES
 POTATOES TOMATOES LEGUMES PEPPERS Apples Colorado Potato Beetles Flea Beetles
 BEANS BEETS Field Beans Maroon Bean Beetles Cucumber Beetles
 CABBAGE CARROTS BROCCOLI BRUSSELS SPROUTS COLLARDS KALE
 KALE Flea Beetles Imported Cabbageworms Cabbage Looper Cabbageworms
 CUCUMBER MELONS SQUASH Cucumber Beetles Pick Weevils
 ASPARAGUS Asparagus Beetle
 CARROTS Vegetable Weevils
 LETTUCE Colorado Potato Beetles
 PEAS Aphids Flea Beetles For Peas Weevils control dust on fresh podding early in podding before pods are full.
 PUMPKINS Apples Flea Beetles Vegetable Weevils
 RADISH SPINACH TURNIPS Flea Beetle
FLOWERS
 FLOWERS such as Roses Chrysanthemums Dahlias Plumes Sweet Peas Phlox
 Delphiniums Marigolds Zinnias Flea Beetles Cucumber Beetles Zabrotus Aphids
 (Green and Rust)
SMALL FRUITS
 BLACKBERRIES LOGANBERRIES RASPBERRIES For Raspberry Curc Borer and
 Raspberry Curc Borer: Dust for 10 days before plants bloom. Repeat at about 2 weeks.
 For Raspberry Fruitworm: Dust for 10 days after first blossom. Repeat in 10 days and
 again 10 days later.
DOGGS and CATS
 To control fleas: Dust thoroughly on hair and rub in with the hand. Avoid getting in
 animal's eyes, nose and mouth. Also thoroughly treat bedding, furniture and bedding.
DISPOSAL
 DO NOT REUSE EMPTY CONTAINER. WRAP CONTAINER AND PUT IN WASTE COLLECTION
 BIN.
 Buyer assumes all risks of use, storage or handling of this material in strict accord
 with directions given herein.
 HI-YIELD CHEMICAL COMPANY
 BOONHAM, TEXAS 75418
 NET WEIGHT 1 LB. (454 GRAMS)

Hi-Yield

Rotenone 100 Insecticide Dust



Controls Listed Insects On
 Vegetables, Small Fruits, Shrubs,
 Flowers and Pets

ACCEPTED

MAR 10 1980

Active Ingredients:	Rotenone	1.0
	Other Cube Resin	2.0
Inert Ingredients:	Total	97.0
		100.0

EPA Reg. No. 34911-21

EPA Reg. No. 34911-21 EPA Est. 7401-TX
KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel For Additional Cautions

34911-21

BEST DOCUMENT AVAILABLE

CAUTION
STATEMENT OF PRACTICAL TREATMENT
 Wash with soap and water.
 Flush with water for 15 minutes. Contact a physician.
PRECAUTIONARY STATEMENTS
HAZAROUS TO HUMANS AND DOMESTIC ANIMALS
CAUTION
 If swallowed. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling and before eating or smoking. Remove and wash clothing before re use.
ENVIRONMENTAL HAZARDS
 Is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where likely to occur. Do not apply when weather conditions favor drift from area. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on this label.
DIRECTIONS FOR USE
 In accordance with Federal law to use this product in a manner inconsistent with its labeling is at the rate of 2 ounces per 50 feet of row on first appearance of insect or damage.
 Repeat at 4 to 7 day intervals until under control.
 Light dustings are more effective than single heavy dustings.
 Apply to food crops within one day of harvest.
VEGETABLES
TOMATOES Eggplants, Peppers. Aphids, Colorado Beetle, Beetles.
BEETS Beetles, Mexican Bean Beetles, Cucumber Beetles.
CABBAGE Beetles, Broccoli, Brussels Sprouts, Collards, Kohlrabi.
BEETLES Imported, Colorado, and Cucumber Beetles, Vegetable Weevils.
SPINACH Spinach. Cucumber Beetles, Flea Weevils.
WEEDS - Vegetable Weevils.
WEEDS - Cabbage Looper.
APHIDS - Flea Weevils. For Flea Weevil control dust on foliage during early bloom.
WEEDS - Flea Weevils, Vegetable Weevils.
SPINACH, TURNIPS - Flea Beetle.
FLOWERS
 As such as Roses, Chrysanthemums, Dahlias, Peonies, Sweet Peas, Phlox, Marigolds, Zinnias. Flea Beetles, Cucumber Beetles, Diabrotica, Acanthos, and Rosy.
SMALL FRUITS
RASPBERRIES - Raspberries. For Raspberry Case Borer and other damage. Dust foliage just before plants bloom. Repeat at about 2 weeks.
FRUIT - Fruit worm. Dust foliage 7 days after first blossom. Repeat in 10 days and 14 days later.
RASPBERRIES - Sawflies, Flea Beetles.
DOGS AND CATS
 Fleas. Dust thoroughly into hair and rub in with the hand. Avoid getting into eyes, nose and mouth. Also thoroughly treat sleeping quarters including bedding.
DISPOSAL
 REUSE EMPTY CONTAINER. WRAP CONTAINER AND PUT IN TRASH COLLECTOR.
 For some risks of use, storage or handling of this material not in strict accordance with directions given hereon.
HI-YIELD CHEMICAL COMPANY
 SONNAN, TEXAS 75418
 NET WEIGHT 1 LB. (454 GRAMS)

Hi-Yield

Rotenone 100

Insecticide

Dust



Controls Listed Insects On
 Vegetables, Small Fruits, Shrubs,
 Flowers and Pets

ACCEPTED
 MAR 10 1980

Active Ingredients:	Rotenone	1.0%
	Other Cube Resin	2.0%
Inert Ingredients:		97.0%
Total		100.0%

EPA Reg No 34911-21 EPA Est 7401-TX-1

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See Back Panel For Additional Cautions

34911-21

BEST DOCUMENT AVAILABLE