

(Front Panel)

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 29 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
34911-27

Hi-Yield

BENOMYL*

Systemic

Fungicide

* Prevents & Controls Black Spot, Powdery
Mildew, Brown Patch, Dollar Spot, Fusarium
Patch and Gray Mold on Roses, Ornamentals,
Flowers and Lawns.

* 2½ Level Tablespoons Makes 2 Gallons of Spray
for Flowers & Ornamentals.


Active Ingredient:

Benomyl Methyl 1-(butylcarbamoyl)-2
benzimidazolecarbamate 25%
Inert Ingredients: 75%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Hi-Yield BENOMYL SYSTEMIC FUNGICIDE provides an effective and economical method for controlling those fungus diseases listed below.

ROSES, ORNAMENTALS, FLOWERS:(Field & Greenhouse). For control of Black Spot on Roses mix 4½ level tablespoons of this product with 2 gallons of water. For Powdery Mildew and Botyris Gray Mold on flowers & Ornamentals, mix 2½ level tablespoons of this product with 2 gallons of water. Spray to thoroughly cover upper and lower sides of plant foliage. Begin applications when disease first appears and repeat at 10-14 day intervals throughout the growing season ... shorten spray intervals during humid, rainy weather.

LAWNS: For control of Brown Patch & Fusarium Patch, apply 4 ozs. of this product to 1,000 sq.ft. For Dollar Spot apply 2 ozs. of this product to 1,000 sq.ft. Use a sufficient amount of water to obtain a thorough coverage(2-5 gallons per 1,000 sq.ft.) As a preventive measure, apply at the earliest appearance of disease and repeat every 10-14 days. When disease conditions are unusually favorable(during high humidity and rainy weather) reduce spray intervals to 5-7 days. As a curative measure(when disease conditions are severe), apply at the first sign of disease and repeat as needed. Addition of a spreader-sticker to the spray mixture enhances curative action of the fungicide and improves distribution of the spray on hard-to-wet leaf surfaces such as roses.

NOTE: If treatment is not effective following the use of this product as recommended, a tolerant strain of fungi may be present; consideration should be given to prompt use of other unrelated fungicides.

Hi-Yield BENOMYL SYSTEMIC FUNGICIDE, when used according to directions, will satisfactorily solve those fungus problems listed on this label.

STORAGE & DISPOSAL

Never allow this product to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of the product as a fungicide. Keep container closed when not in use. Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS

CAUTION: May irritate eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after using. Keep away from fire and sparks.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Est. No. 7401-TX-1

Hi-Yield Chemical Company

EPA Reg. No. 34911-

Bonham, Texas 75412

Net Contents 1 lb.