

# UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

SEP -7 2006

Ms. Peggy Galloway Landis International

Agent for: SSI Maxim Company Inc. 3185 Madison Hwy, PO Box 5126 Valdosta, GA 31603-5126

Subject:

Sprakil S-5 Brush Control Granules EPA Registration Number 34913-10 Resubmission dated June 28, 2006

Dear Ms. Galloway:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

- 1) Remove the statement "Do not allow domestic livestock to graze in treated areas" from the section NON-CROP WEED CONTROL INDUSTRIAL USE ONLY.
- 2) Revise "to the extent allowed by law" to "to the extent consistent with applicable law" in the section CONDITIONS OF SALE.
- 3) In the section DIRECTIONS FOR USE, add the active ingredient lb equivalent to the rate of product stated wherever a product rate is listed.

  Example: "do not apply more than 80 lbs per acre of product (X lbs a.i.)"
- 4) On page 2, remove the statements "If the treated area is to be used for haying, do not apply more than 80 lbs per acre of product, and do not apply this product more than once per year. Grazing is allowed in areas receiving band or individual plant treatments of 80 lbs per acre or less of product" as these statement are already found on page 1 in the DIRECTIONS FOR USE section.

Submit one (1) copy of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records.

Page 2 EPA Registration Number 34913-10

If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

James A. Tompkins Product Manager 25 Herbicide Branch

Registration Division (7505P)

## NON-CROP WEED CONTROL INDUSTRIAL USE ONLY

SpraKii S-5 is Intended for total vegetation control. It is an extremety active herbicide which will kill trees, shrubs, and other forms of desirable vegetation having roots extending into the treated area. Feeder roots of many species of desirable vagetation extend many feet beyond the drip line of the branches, and a very small amount of SpraKil S-5 in contact with one feeder root of a tree, should on other desirable vegetation may cause serious injury or death to the

An arboriculturist (tree expert) should be consulted to help you to determine if the area of proposed application is free of all roots of desirable vegetation. The effect of SpraKil S-5 on desirable vegetation may be irreversible, and its presence in the soil may prevent growth of other desirable vegetation for some years after application. Read the entire label before using SpraKil S-5 to determine if this product is suitable for the desired purpose.

Do not use SpraKil S-5 on areas such as walks, driveways, streets. lawns, patios, tennis courts, swimming pools, cemeteries, or other landscaped areas, or under esphalt or concrete pavement where future landscaping is planned. Do not apply on or near field crops. Do not apply on any area into which the roots of desirable vegetation may extend. Do not apply in locations where the chemical may be washed by runoff or sheet erosion in contact with roots of desirable vegetation, as severe injury or death may occur. Roots of trees, shrubs, and other desirable vegetation may extend far beyond the drip line of the plant's branches.

Ditchbank Usage: Do not apply Sprakii S-5 to any portion of the ditchbank that will come into direct contact with water as movement of Sprakii S.5 in this water to popularnot plant species may result in the injury or death of those plants. Do not apply on ditches used to transport irrigation or potable water. Keep from contact with other pesticides and seeds.

Thoroughly clean all traces of SpraKil S-5 from application equipment after use. Do not empty residues cleaned from application equipment on areas where they may come in contact with the roots of desirable vegetation or the water source for such vegetation.

Herbaceous Perennials (nonwoody plants): Do not allow domestic livestock to graze in treated areas. Do not cut forage from treated areas for feeding livestock. SpraKil S-5 may injure or suppress certain herbaceous vegetation in the treated area. Do not apply where such injury cannot be tolerated. Injury to most herbaceous perennials is reduced if SpraKit S-5 is applied when this vegetation is dormant.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store product in original container only. away from other pesticides, fertilizer, food, or feed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## **CONDITIONS OF SALE**

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one other than officer of Seller is authorized to make any warrant, guarantee, or direction concerning this product. 2. Because the time, place, rate of application, and other conditions for use are beyond the Seller's control, to the extent allowed by law the Seller's liability from handling, storage, and use of this product is limitled to reptacement of product or refund of the purchase price.



# **KEEP OUT OF REACH OF CHILDREN** CAUTION! Hold eye open and rinse slowly and gently with water for 15-20 minutes.

IF IN EYES: Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye Call a poison control center or doctor for treatment advice. IF ON SKIN Take off contaminated clothing. OR CLOTHING: Rinse skin immediately with plenty of water for 15-20 migutes.

Call a poison control center or doctor for treatment advice.

Call a poison control center or doctor immediately for treatment advice.

· Have person sip a glass of water if able to swallow

 Do not induce vomiting unless told to do so by a polson control center or doctor. . Do not give anything by mouth to an unconscious person.

For emergency information, contact CHEMTREC at 1-800-424-9300 (24 hours).

Have the product container or label with you when calling a polson control center or doctor, or going for treatment. SEE ADDITIONAL PRECAUTIONARY STATEMENTS & DIRECTIONS FOR USE

MANUFACTURED BY:

# SSI MAXIM CO., INC.

P.Q. 1954 1954 Kilgore, TX 75663 • (800) 346-4781

**NET WEIGHT:** 8 x 6 LB (8 x 2.72 KG) 5 

EPA REG. NO. 34913-10 EPA EST. NO. 39578-TX-1

R490

## PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin or swallowed. Avoid contact with eyes, skin or clothing. Prolong or trequently repeated skin contact may cause allergic reaction in some

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

- Applicators and other handlers must wear:
- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material

#### **ENVIRONMENTAL HAZARD**

in case of spills, collect, cover, or incorporate granules/pellets spilled on the soil surfaces to prevent contamination to water. Do not apply to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

#### **GROUND WATER ADVISORY**

This product is known to leach through soft into ground water under certain conditions as a result of registered (rangeland and non-crop) uses. Use of this product in areas where soils have rapid to very rapid permeability, particularly where the water table is shallow, may result in ground water contamination.

#### **Use Restrictions for Ground Water Protection**

Vulnerable Sites: To minimize any movement of tehuthluron to enhantere water in not exceed the application rates energial below on treatment sites where soils have a sand or loamy sand texture throughout the soil profile and all of the following characteristics:

- Rapid to very rapid permeability.
- 2. Absence of well-defined organic layers or a textural B-horizon (restricting layer of fine-textured soil).
- 3. The water table of an underlying aquifer is shallow.

An aquifer is defined as an "underground saturated, permeable geologic formation capable of producing significant quantities of water to a well or spring", it is the ability of the saturated zone, or portion of that zone, to yield water which makes it an aquifer (American Chemical Society, 1983). Local agricultural agencies can provide further information on the type of soil in your area and the location of shallow ground water aquifers.

Do not apply SpraKli S-5 in areas where the water table is predominately shallow (5 feet or less), such as marshy or sub irrigated areas, or areas immediately adjacent to streams or lakes which are periodically flooded, unless such use is allowed under a stateapproved pesticide management program. Note: Also on such areas, woody plants rooted directly in a shallow water table are minimally affected by applications of tebuthiuron, and poor woody plant control will result.

Do not apply SpraKii S-5 within areas identified by state or local authorities as protected ground water recharge zones.

#### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

For vegetation control by broadcast (ground equipment) and banded applications: If the treated area is to be used for having, do not apply more than 80 lbs. per acre of product, and do not apply this product more than once per year. Grazing is allowed in areas receiving band or individual plant treatments of 80 lbs. per acre or less of product. The maximum tabel rate and frequency of application is 40 lbs. SpraKil S-5 per acre once every three years for vulnerable areas (where soils are sandy and depth to water table is shallow.) For all other areas, Sprakil S-5 may be applied one time in a 3-year period at a rate of up to 80 lbs. per acre of product; however, no more that 120 lbs. per acre may be applied in two consecutive treatments in any 6 year period.

Spot treatments (hand-held equipment): SpraKil S-5 may be applied at rates up to the equivalency of 120 lbs./acre when needed.

IF SWALLOWED:

# SPRAKIL S-5 **BRUSH CONTROL GRANULES**

# **DIRECTIONS FOR USE** READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING

Sprakil S-5 Granules is a non-selective herbicide for controlling a wide range of annual and perennial weeds and grasses and broadleaved weeds and woody plants. It is recommended only for non-cropland areas such as, AIRPORT RUNWAYS, UTILITY SUBSTATIONS AND RIGHTS-OF-WAY: TANK FARMS: RAILWAY ROADBEDS AND BALLAST: RAILROAD RIGHTS-OF-WAY: ROAD SHOUL-DERS WHERE NO VEGETATION IS DESIRED: AT THE BASE OF HIGHWAY GUARDRAILS: SIGN-POSTS AND MARKERS AT THE BASE OF TRANSMISSION TOWERS AND POLES: AROUND INDUSTRIAL BUILDINGS; LUMBERYARDS; RAILROAD YARDS; DITCHBANKS; FIREBREAKS; AND FENCE ROWS INCLUDING GRAZED AREAS ON THESE SITES. Effects are slow to appear and will not become apparent until the chemical has been carried into the root zone of the weeds by moisture. Control will last for an extended period. The duration of nonproductivity is dependent upon type of soil, amount of rain, and other conditions. Users should consult state agricultural experimental stations or extension service weed specialists for recommendations as to use in their particular area.

If the treated area is to be used for haying, do not apply more than 80 lbs per acre of product, and do not apply this product more than once per year. Grazing is allowed in areas receiving band or individual plant treatments of 80 lbs per acre or less of product.

How to apply: Use a mechanical spreader or other device which will spread the required amount eventy over the area to be treated. It may be applied when top growth is dense, but better control will be obtained when weeds are short or when excess growth is moved and removed before applying.

When to apply: Best results will be obtained if applied shortly before or at the time plant growt begins. Rains or thorough watering after application will cause the material to move to the root zon and thus speed, up control. Do not use when ground is frozen.

#### WOODY PLANT CONTROL

Apply SpraKil S-5 uniformly to the woody plant infested area or by individual plant treatmen Application can be made anytime during the year except when the ground is frozen. Effects are slo to appear and will not become apparent until sufficient moisture has carried the SpraKil S-5 into the root zone of the target species. The time required to achieve control is dependent on soil type, amount of rainfall and depth of species rooting. Some species may go through several defoliations and refolia tions over a period of approximately two or three years prior to dying.

## INDIVIDUAL (SPOT) WOODY PLANT APPLICATION

Apply SpraKii S-5 by hand tossing 2 ounces of the granules around the base of the unwanted individu al woody stem. Wear protective gloves when applying granules by hand. Repeat this process for eac unwanted woody plant in the area. Caution: Do not use SpraKil \$-5 in this manner in any area wher desirable species are in the vicinity of the plants to be eliminated. SpraKil S-5 will injure or contri other herbaceaous vegetation in the treated area. Do not apply where such injury cannot be tolerated

# **BROADCAST APPLICATION**

Apply SpraKil S-5 using a mechanical spreader or other device calibrated to deliver the recommended amount of material over the area to be treated.

#### **WOODY PLANT CONTROL - APPLICATION RATES**

SpraKij S-5 applied at the rate of 40 pounds per acre will control the following species:

Allanthus aftissima (Tree-of-heaven) Aloysia lycioides (Whitebrush) Artemisia vulgaris (Mugort) Liriodendrom tulipifera (Tuliptree)

Nyssa sylvatica (Black gum) Quercus nigra (Water oak) Ulmus americana (American elm)

SpraKil S-5 applied at the rate of 60 pounds per acre will control the following species:

Carva texana (Black hickory)

Quercus maritandica (Blackiack Oak) — Quercus stetlata (Post oak)

Symphoricarpos orbiculatus (Buckbrush) Ulmus alata (Winged elm)

SpraKii S-5 applied at the rate of 80 pounds per acre will control the following species:

Acacia berlandieri (Guajillo) Acacia rigidula (Blackbrush acacia) Acer negundo (Boxelder)

Cettis occidentalis (Western hackberry) Campsis radicans (Trumpetcreeper) Celtis pallida (Granjeno)

Larrea tridentata (Creosotebush) Myrica cerifera (Southern waxmyrtle)

Rhus copallina (Shining sumac) Rosa multiflora (Multifora Rose) Rubus allegheniensis (Allegheny blackberry)

SpraKil S-5 applied at the rate of 100 pounds per acre will control the following species:

Acer macrophyllum (Bigleaf maple) Alnus rubra (Red atder)

Cornus Iterida (Flowering degwood) Gaultheriua shallon (Salal)

Juniperus virginiana (Eastern redcedar) Pinus echinata (Shortleaf pine)

Pinus virginiana (Virginia pine) Prunus emarginala (Bitter cherry) Pseudotsuga menziesii (Douglas fir) Quercus alba (White oak)

Rubus laciniatus (Evergreen blackberry) Robus occidentalis (Black raspberry)

SpraKil S-5 applied at the rate of 120 pounds per acre will control the following species:

Acer circinatum (Vine maple) Cornus spp. (Dogwood)

Crataegus spp. (Hawthorn) Fraxinus americana (White ash) Haplopappus tenuisectus (Burroweed) Liquidambar styraciflua (Sweetgum)

Prosopis julitlora (Mesquite) Prunus serotina (Black Cherry) Rhus glabra (Smooth sumac) Rhus typhina (Staghorn sumac) Salix spp. (Willow) Smilax rotundifolia (Common greenbriar)

Ulmus parvifolia (Chinese Eim) Ulmus rubra (Slippery elm)



SSI MAXIM CO., INC.

P.O. Box 1954 • Kilgore, TX 75663

R480

HUDSON 845600-63