

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMANS  
& DOMESTIC ANIMALS**

**CAUTION**

May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARD**

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

**DIRECTIONS FOR USE**

It is a violation of federal laws to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE**

To maximize the shelf life of this product, store in dry area where the container is not likely to come in contact with moisture.

**DISPOSAL:**

Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

(For Plastic Buckets) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(For Bags) Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

# SPRAKIL B-10 PELLETS WEED KILLER

**ACTIVE INGREDIENTS:**

Bromacil (5-bromo-3-sec-butyl-6-methyluracil)..... 10%

**INERT INGREDIENTS:**..... 90%

**BEST AVAILABLE COPY**

ACCEPTED

NOV 17 1988

KEEP OUT OF REACH OF CHILDREN

## CAUTION!

### STATEMENT OF PRACTICAL TREATMENT

**If in eyes:** Flush with plenty of water. Get medical attention if irritation persists.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

MANUFACTURED FOR

# SSI INDUSTRIES, INC.

A SUBSIDIARY OF MOBLEY INDUSTRIES, INC.

4711 PIEDMONT ROAD  
HUNTINGTON, W. VA 25704

EPA Registration No. 34913-16  
EPA Establishment No. 20000-01

**NON-CROP BRUSH & WEED CONTROL  
INDUSTRIAL USE ONLY**

Sprakil B-10 Pellets is a pelleted product for dry applications to the soil for the control of brush and many annual and perennial weeds. Effects are slow to appear and will not become apparent until the chemical has been activated by moisture. The degree of control and duration of effect will vary with the amount of chemical applied, soil type, rainfall and other conditions.

Sprakil B-10 Pellets is non-corrosive to equipment, non-volatile, and non-flammable.

**BRUSH CONTROL**

To control undesirable woody plants on non-cropland areas such as RAILROAD RIGHT OF WAYS, STORAGE AREAS, INDUSTRIAL PLANT SITES and DRAINAGE DITCHES: Apply pellets in spring or early summer as a broadcast or basal (spot) treatment; for use on drainage ditches, apply as a basal (spot) treatment only. Growth of herbaceous vegetation will be prevented in the treated area for an extended period of time. Note: For effective brush control and prevention of damage to desirable vegetation: do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use in irrigation ditches nor on right of ways or other sites where marketable timber or other desirable trees or shrubs are immediately adjacent to the treated area.

**BASAL (SPOT) TREATMENT**

To control undesirable woody plants such as cottonwood, hackberry, maple, oak, poplar, red bud, sweet gum, wild cherry, willow and winged elm: apply pellets in spring or early summer by spreading 1 to 2 tablespoonfuls on the ground (cover an area 1/2 to 1 sq. ft.) at the base of each brush stem or clump 2 to 4 inches in basal diameter; on larger clumps apply 1 tablespoonful to each of 3 to 4 spots around the base of the brush. Apply to uphill side of plant if ground is not level.

**BROADCAST TREATMENT**

Apply in spring or early summer: use 60 to 96 lbs. per acre to control oak, willow, sweet gum and pine; use 120 to 240 lbs. per acre to control American elm, winged elm, hackberry and cottonwood.

**WEED CONTROL**

To control most weeds for an extended period of time on non-cropland areas such as EQUIPMENT, STORAGE SITES, DITCH BANKS, AROUND FARM BUILDINGS, or other non-crop areas that are not immediately adjacent to cropland or desirable trees and shrubs: Apply pre-emergence or early post-emergence to weed growth by spreading Sprakil B-10 Pellets uniformly over the surface of the ground to be treated. Apply at the following rates; use the higher levels of the dosage ranges on absorptive soils (those high in organic matter or carbon).

**ANNUAL WEEDS AND GRASSES**, such as foxtail, ryegrass, wild oats, crabgrass, chestgrass, bromegrass, ragweed, lambquarters, puncturevine and turkey mullein: use 30 to 60 lbs. per acre.

**PERENNIAL WEEDS AND GRASSES**, such as smooth brome, Bahiagrass, bluegrass, redtop, purpletop, quackgrass, bromesedge, aster, dandelion, dog fennel, goldenrod, plantain and wild carrot: use 60 to 120 lbs. per acre. Higher rates of 120 to 240 lbs. per acre are required for Johnsongrass and other hard-to-kill perennials such as Bermudagrass, Vasey-grass, Dallisgrass, nutsedge, satygrass, bouncingbet, dogbane, bracken fern and horsetail; apply these high rates only as spot treatments to localized infestations of weeds.

**RETREATMENT**

Apply 30 to 60 lbs. per acre when annual weeds and grasses reappear on sites where weed growth has been controlled.

**FOR SMALL AREAS** - one-half lb per 435 square feet is approximately 50 lbs. per acre.