



SSI MOBLEY CO., INC.

SPRAKIL SK-13 GRANULAR WEED KILLER

ACTIVE INGREDIENTS:	
Tebuthiuron: N-(5-(1, 1-dimethylethyl)-1, 3, 4-thiadiazol-2-yl)-N, N'dimethylurea	1.0%
Diuron: [3-(3, 4-dichlorophenyl)-1, 1-dimethylurea]	3.0%
INERT INGREDIENTS:	96.0%
Total:	100.0%

NON-CROP WEED CONTROL INDUSTRIAL USE ONLY

SpraKil SK-13 Granular is a nonselective herbicide for controlling a wide range of annual and perennial weeds and grasses. It is recommended only for non-cropland areas such as railroad rights-of-way and industrial areas. 150 to 400 pound applications of SpraKil SK-13 Granular per acre usually result in a nonproductive condition of the soil for a period of a year or more. The duration of nonproductivity is dependent upon rainfall, soil type and other conditions.

SpraKil SK-13 Granular should be applied as furnished with a seed spreader, a fertilizer spreader, a shaker type applicator, or with any equipment which will distribute the chemical uniformly over the area to be treated. For the control of annual and most perennial weeds and grasses, apply SpraKil SK-13 Granular at the rate of 150 to 400 pounds per acre on smaller plots 10.35 to 11.0 pound per 100 square feet. Repeat spot treatments may be required when deep rooted perennial weeds are present. To obtain the best results, SpraKil SK-13 Granular should be applied to the ground where it will be absorbed by the roots. Applications made early in the season have been found to give results superior to applications at a later seasonal date. SpraKil SK-13 Granular may, however, be applied at any time of the year except when the ground is frozen. Maximum effectiveness in arid regions is obtained when application is made just prior to the rainy season. Users should consult state agricultural experimental stations or extension service weed specialists for recommendations as to use in their particular area.

IMPORTANT: Do not apply on or near desirable woody or herbaceous plants or on areas where their roots may extend, because of possible injury to such plants. Thoroughly clean spreading equipment with a suitable chemical cleaner before using for other purposes (or do not use same spreading equipment for other purposes). Do not contaminate irrigation ditches or water used for domestic purposes. Do not use on croplands or any land to be used for subsequent cropping. Keep animals off treated areas. Do not re-use this container. Destroy when empty.

**KEEP OUT OF REACH OF CHILDREN
CAUTION!**

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED BY:
SSI MOBLEY CO., INC.
1909 N. Longview Street • Kilgore, TX 75662

STORAGE AND DISPOSAL

STORAGE:
To maximize the shelf life of this product, store in dry area where the container is not likely to come in contact with moisture. Do not store near fertilizers, seeds, insecticides or fungicides.

DISPOSAL:
Do not contaminate water, food or feed by storage or disposal.

Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

(For Plastic Buckets) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill or incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(For Bags) Completely empty bag into application equipment. Then dispose of bag in sanitary landfill or by incineration, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

ACCEPTED
MAY 20 1997
MAY 20 1997
[See attached replacement]

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 34913-15

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one other than officer of Seller is authorized to make any warrant, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

EPA REGISTRATION NO. 34913-15
EPA ESTABLISHED NO. 39578-TX-1

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

May cause irritation of eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

ENVIRONMENTAL HAZARD

In case of spills, collect, cover or incorporate granules/pellets spilled on the soil surfaces to prevent contamination to water. Do not apply to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

GROUND WATER ADVISORY

This chemical is known to leach through soil into ground water under certain conditions as a result of registered (rangeland and non-crop) uses. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

A shallow water table is defined as depth to water table of 30 feet or less. Permeable soils include, but are not limited to, sandy soils.

DIRECTIONS FOR USE:

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

For vegetation control by broadcast (aerial and ground equipment) and banded applications: The maximum label rate and frequency of application is 1-2 lbs. ai/acre once every three years for vulnerable areas (where soils are sandy and depth to water table is shallow). For all other areas, may be applied one time in a 3-year period at rates up to 4 lbs. ai/acre; however, no more than 6 lbs. ai/acre may be applied in two consecutive treatments in any 6-year period.

Spot treatments (hand-held equipment): May be applied at rates up to the equivalency of 6 lbs. ai/acre when needed.