

SSI MAXIM CO., INC.



STORAGE AND DISPOSAL

STORAGE:

To missimize the shelf life of this product, store in dry area where the container is not likely to come in contact with moleture. On not store near fartilizers, seeds, inspecticles, or fundation.

DISPOSAL

Do not contaminate water, food, or feed by storage or discount.

Wastes resulting from this product may be disposed of on sits or at an approved waste disposal facility.

(For Plastic Bucket) Triple these (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary familial or incineration, or, if allowed by state and social authorities, by burning, if burned, stay out of arnoke.

(For Bage) Completely empty bag into application equipment, Depose of bag in a sanitary lendfill or by inchemition, or, if allowed by state and local authoriises, by burning, if burned, stay out of smoke,

EAUTION!

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Fissis with plenty of water. Gal medical attention if initiation persists.

UF ON SIGN: Wash with planty of soap and water, Get medical attention if irritation persists.

SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vontiling by fourthing back of throat with finger. Do not induce vomiting or give engiting by mouth to an unonactous person.

SEE ADDITIONAL PRECAUTIONALLY STATEMENTS

Weed Blast 4-G Weed Killer

| | ive ingredients: | |
|---|--|---|
| 1 | Bromacii (5-brome-3-sec-butyi-6-mathyluracii) | , |
| | Diuron: [3-(3,4-dichlorophenyi)-1, 1-dimethylurea]2.0% | |
| | INERT INGREDIENTS: 95.0% | |
| 1 | Total: | , |

NON-CROP WEED CONTROL INDUSTRIAL USE ONLY

Weed Blast 4-G is a nonsetective herbicide for controlling a wide range of annual and perennial weeds and grasses. It is recommended only for non-cropland areas such as relificad, utility and highway rights-of-way, and industrial areas. 200 to 400 pound applications of Weed Blast 4-G per acre usually result in a nonproductive condition of the soil for a period of a year or more. The duration of nonproductivity is dependent upon rainfall, soil typs, and other conditions.

Weed Blast 4-G should be applied as furnished with a seed spreader, a fertilizer spreader, or with any equipment which will distribute the chemical uniformly over the area to be freated. For the control of annual and most perennial weeds and grasses, apply Weed Blast 4-G at the rate of 200 to 400 pounds per acre; on smaller plots, 1/2 to 1 pound per 100 square feet. Repeal spot treatment may be required when deep rooted perennial weeds are present. To obtain the best results, Weed Blast 4-G should be applied to the ground where it will be absorbed by the roots. Applications made early in

the season have been found to give results superior to applications at a later seasonal date. Weed Blast 4-G may, however, be applied at any time of the year except when the ground is frozen. Maximum effectiveness in and regions is obtained when application is made just prior to the rainy season. Users should consult state agricultural experimental stations or extension service weed specialists for recommendations as to use in their particular area.

ENTRY RESTRICTIONS: Oo not enter or allow others to enter the treated area until dusts have settled.

SMPORTANT: Do not apply on or near desirable woody or herbeceous plants or on areas where their roots may extend because of possible injury to such plants. Thoroughly clean spreading equipment with a suitable chemical cleaner before using for other purposes (or do not use same spreading equipment for other purposes). Dispose of finance according to "Dispose!" instructions found under "Storage and Disposal" heading. Do not use on croplands or any land to be used for subsequent cropping. Keep animals off freated areas.

PRECAUTIONARY STATEMENTS

HAZARO TO HUMANS & DOMESTIC ANIMALS

CAUTION -

Hammful if swallowed. May cause initiation of eyes, nose, throat, and skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

Do not apply directly to water, or to areas where surface water is present, or to intertidat areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of faderal law to use this product in a manner inconsistent with its labeling.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for use as directed on the label. No one other than officer of Seller is authorized to make any warrant, guarantee, or disaction concerning this product. 2. Because the time, place, rate of application, and other conditions of use are beyond Seller's control, Seller's liability from handling, storage, and use of this product is limited to replacement of product or retaind of purchase price.

MANUFACTURED BY: 861 MAXIM CO., IMC. P.O. Box 1954 • Kilgore, TX 76683

> NET WEIGHT: 20 LB (9.07 KQ)

EPA REG. NO. 34913-19 EPA EST. NO. 39578-TX-1 023

WB40/20KPley, 10-00)

HUOSON-LONGVIEW MS00

ACCEPTED

1 - NOV-3 0-2000-

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No 3 1912-19