10f5



U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W. Washington, D.C. 20460 EPA Reg. Number: 34913-22

SEP 2 1998

NOTICE OF PESTICIDE:

x Registration
Reregistration

tunder FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:

SSI Maxim Topsite 2.5G Herbicide

Name and Address of Registrant (include ZIP Code):

SSI Maxim Company, Inc.

P.O. Box 1954

Kilgore, TX 75663

Sote: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- 2. Add the phrase "EPA Registration No. 34913-22" to your label before you release the product for shipment.
- 3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

Signature of Approving Official:

Jug L

Date:

SEP 2 1933

EFA Form 8570-6

1

page 2 EPA Reg. No. 34913-22

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

James A. Tompkins Product Manager (25) Herbicide Branch Registration Division (7505C)

Enclosure



ACCEPTED

2 1998

For Use In Noncrop Areas **And Noncropland Farmsteads**

ACTIVE INGREDIENTS:

Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-	
5-oxo-1 <i>H</i> -imidazol-2-yl]-3-pyridinecarboxylic acid)	0.5%
Diuron (3-[3,4-dichlorophenyl]-1,1-dimethylurea),	2.0%
INERT INGREDIENTS	97.5%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION!/iPRECAUCION!

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

> See Additional Precautionary Statements and Directions For Use Inside



SSI MAXIM CO., INC.

P. O. Box 1954 • Kilgore, TX 75663

EPA Est. No.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION!

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water.

PHYSICAL AND CHEMICAL HAZARDS

TOPSITE 2.5G herbicide should be stored only in lined paper bags, stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT store TOPSITE 2.5G herbicide in unlined steel (except stainless steel) containers.

Keep dry.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment washwaters.

IMPORTANT

DO NOT use on food or feed crops. DO NOT treat irrigation ditches, or water used for crop irrigation or for domestic uses. DO NOT apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result. Keep from contact with fertilizers, insecticides, fungicides and seeds. DO NOT apply or rinse equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use on lawns, walks, driveways, tennis courts, or similar areas. DO NOT use in California.

GENERAL INFORMATION

TOPSITE 2.5G herbicide is recommended for control of many annual and perennial grasses and broadleaf weeds on noncropland areas such as railroad, utility, pipeline and highway rights-of-way, petroleum tank farms, pumping installations, fence rows, storage areas, noncropland farmsteads and other similar areas. TOPSITE 2.5G herbicide may be applied at any time throughout the year; however, for maximum effectiveness, the product should be applied prior to weed emergence or when weeds are small. In addition, TOPSITE 2.5G herbicide should be applied prior to ex-

pected rainfall. The degree and duration of control of Kochia and other weed species will depend upon the amount of product applied, the amount of rainfall, soil texture, and other environmental and biological factors.

CONDITIONS OF SALE

- Seller warrants that this product consists of the ingredients specified and is reasonable fit for use as directed on the label. No one other than an officer of Seller is authorized to make any warrant, guarantee, or direction concerning this product.
- 2. Because the time, place, rate of application, and other conditions of use are beyond Seller's control, Seller's liability from handling, storage, and use of this product is limited to replacement of product or refund of purchase price.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL FOR BAG: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR PAIL: Triple rinse (or equivalent). Then offer for recycling or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATION INSTRUCTIONS

Uniformly apply TOPSITE 2.5G herbicide with properly calibrated ground equipment, including hand-held, spreaders, at a rate of 200 to 300 lbs. product per acre (0.5-0.7 lb./100 sq. ft.).

A preemergent/early postemergent application of TOPSITE 2.5G herbicide will control many weed species, including the following:

GRASSES

Annual bluegrass (Poa annua)

Bahiagrass (Paspalum notatum)

Beardgrass (Andropogon spp.)

Bermudagrass (Cynodon dactylon)

Big bluestem (Andropogon gerardi)

Broadleaf signalgrass (Brachiaria platyphylla)

Browntop millet (Brachiara ramosa)

Canada bluegrass (Poa compressa)

Cattail (Typha spp.)

Centipedegrass (Eremochloa ophiuroides)

Cheat (Bromus secalinus)

Crabgrass (Digitaria spp.)

Dallisgrass (Paspalum dilatatum)

Downy brome (Bromus tectorum)

Fall panicum (Panicum dichotomiflorum)

Fescue (Festuca spp.)

Foxtail (Setaria spp.)

Goosegrass (Eleusine indica)

Guineagrass (Panicum maximum)

Italian ryegrass (Lolium multiflorum)

Johnsongrass (Sorghum halepense)

Kentucky bluegrass (Poa pratensis)

Lovegrass (Eragrostis spp.)

Orchardgrass (Dactylis glomerata)

Paragrass (Brachiaria mutica)

Prairie cordgrass (Spartina pectinata)

Prairie threeawn (Aristida oligantha)

Quackgrass (Agropyron repens)

Sandbur (Cenchrus spp.)

Sand dropseed (Sporobolus cryptandrus)

Smooth-brome (Bromus inermis)

Timothy (Phleum pratense)

Torpedograss (Panicum repens)

Vaseygrass (Paspalum urvillei)

Wild barley (Hordeum spp.)

Wild iris (Iris versicolor)

Wild oats (Avena fatua)

Wirestern muhly (Muhlenbergia frondosa)

Witchgrass (Panicum capillare)

BROADLEAF WEEDS

Bigleaf aster (Aster macrophyllas)

Blue mustard (Chorispora tenella)

Bull thistle (Cirsium vulgare)

Burdock (Arctium spp.)

Buttercup (Ranunculus testiculatus)

Camphorweed (Heterotheca subaxillaris)

Canada thistle (Cirsium arvense)

Carpetweed (Mollugo verticillata)

Carolina geranium (Geranium carolinianum)

BROADLEAF WEEDS

Clover (Trifolium spp.)

Cocklebur (Xanthium strumarium)

Common chickweed (Stellaria media)

Common ragweed (Ambrosia artemisiifolia)

Dandelion (Taraxacum officinale)

Dock (Rumex spp.)

Dogfennel (Eupatorium capillifolium)

Eclipta (Edipta prostrata)

Filaree (Erodium spp.)

Fleabane (Engeron spp.)

Florida beggarweed (Desmodium tortuosum)

Florida pusley (Richardia scabra)

Giant ragweed (Ambrosia trifida)

Hoary vervain (Verbena stricta)

Horseweed (Conyza canadensis)

Indian mustard (Brassica juncea)

Kochia (Kochia scoparia)

Lambsquarters (Chenopodium album)

Lespedeza (Lespedeza spp.)

Little mallow (Malva parviflora)

Milkweed (Asclepias spp.)

Miners lettuce (Montia perfoliata)

Mullein (Verbascum spp.)

Nettleleaf goosefoot (Chenopodium murale)

Oxeye daisy (Chrysanthemum leucanthemum)

Pepperweed (Lepidium spp.)

Pigweed (Amaranthus spp.)

Plantain (Plantago spp.)

Poison hemlock (Conium maculatum)

Pokeweed (Phytolacca americana)

Primrose (Oenothera kunthiana)

Prostrate knotweed (Polygonum aviculare)

Purslane (Portulaca spp.)

Silverleaf nightshade (Solanum elaeagnifolium)

Smartweed (Polygonum spp.)

Sorrel (Rumex spp.)

Sowthistle (Sonchus spp.)

Sunflower (Helianthus spp.)

Sweet clover (Melilotus spp.)

Tansymustard (Descurainia pinnata)

Texas thistle (Cirsium texanum)

Western ragweed (Ambrosia psilostachya)

White heath aster (Aster pilosus)

Wild carrot (Daucus carota)

Wild lettuce (Lactuca spp.)

Wild parsnip (Pastinaca sativa)

Wild turnip (Brassica campestris)

Woollyleaf bursage (Ambrosia grayii)

Yellow starthistle (Centaurea solstitialis)

Yellow woodsorrel (Oxalis stricta)