


BEST DOCUMENT AVAILABLE

MICROBIOCIDE 610-L



ACTIVE INGREDIENT

Poly(oxyethylene (dimethyliminio) ethylene-
(dimethyliminio) ethylene dichloride) 6.0%

INERT INGREDIENT 94.0%

This product contains 0.5 lb. of active ingredient per gallon
and weighs 8.44 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROBIOCIDE 610-L is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime and other deposits.

An initial slug addition of 10.1 to 25.25 fluid ounces of MICROBIOCIDE 610-L per 1000 gallons of water to provide a concentration of 80 to 200 parts per million of MICROBIOCIDE 610-L, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 2.52 to 25.25 fluid ounces of MICROBIOCIDE 610-L per 1000 gallons of water 20 to 200 parts per million of MICROBIOCIDE 610-L should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

STORAGE & DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. Do not reuse empty container. Triple rinse (or equivalent) and dispose of container in an incinerator or in an approved landfill, or bury in a safe place. Rinse that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by

MARKAY LABORATORIES
 Markay Laboratories
 209 South Van Norman Road
 Montebello, California 90640

EPA REG NO 35135-5
NET CONTENTS

EPA EST. NO. _____