

[35138-3]

DIRECTIONS FOR USE

DILUTION INSTRUCTIONS: Dilute to a 0.25% emulsion by adding 1 fl. oz. of this 2.0% concentrate to 7 fl. oz. of deionized water (or 18 1/3 fl. oz. of concentrate in one gallon of water).

INDOORS — FOR HOUSEFLIES, MOSQUITOES, AND GNATS: Apply as a space spray. Mix well before using. Fill spray tank with undiluted SBP-1382-0. 25% insecticide spray formulation. Adjust pressure and nozzle to deliver a fine mist spray. Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room especially windows and light sources which attract these insects. Keep the spray nozzle at all times at least four feet from fabrics to avoid wetting or staining. Fill the room with mist at the dosage rate of one fluid ounce per 1,000 cu. ft. of space. Then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed and before re-entry.

INDOORS — FOR CRAWLING INSECTS: Apply as a contact spray. Mix well before using. Fill spray tank with undiluted SBP-1382-0. 25% insecticide spray formulation. Keep spray nozzle about 8 to 12 inches from surface to be treated. Adjust pressure and nozzle to deliver a coarse spray with particle size larger than that of the average aerosol. When possible, spray directly on roaches, earwigs, grain mites, sowbugs, spiders and centipedes to reduce infestations. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Apply spray until area is moist, but not to the point of run-off. Spray directly on insects where possible.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FLEAS: Protect dogs and cats from fleas by spraying their resting areas. Hold spray nozzle about 18 inches from area to be treated. Apply spray at the rate of two seconds per linear foot. Direct spray on resting areas and floor around these areas. Treat bedding the same way. Also spray entire inside surface of dog houses. Repeat as necessary.

Also treat dogs and cats with a registered product before they re-enter the area just treated with this product.

NET CONTENTS

WINCO

No. 2 E. C.

ACTIVE INGREDIENTS:

*+(5-Benzyl-3-furyl)methyl 2, 2-dimethyl-3-(2-methylpropenyl)cyclopropanecarboxylate	2.000% w/w
Related compounds	0.274%
INERT INGREDIENTS:	97.726%
	<u>100.000% w/w</u>

*Cis/trans isomers ratio: max. 30% (+)cis and min. 70% (+)trans.
+U. S. Patent No. 3,465,007.

USE ONLY AS DIRECTED

WARNING

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of feed and food-stuffs. Remove pets, birds and cover fish aquariums before spraying. Do not use on edible crops.

**STORE AT TEMPERATURES ABOVE 40°.
MIX WELL BEFORE USING.**

Manufactured by

WINCO CHEMICAL COMPANY, INC.

P. O. Box 10692
JACKSON, MISSISSIPPI 39209

FOOD AREAS:
cessing plants,
mercially prepared
food is exposed

CONTAINERS:
destroy containers
away from water

OUTDOOR AREAS:
as a coarse contact

ANTS: Spray where possible.

FLEAS:

SPIDERS:
as necessary.

WASPS:
insects are at risk
contacting as necessary

E.P.A. Reg. No.

ACCEPTED

JUL 15 1975

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the purpose of registered under EPA Reg. No. 35138-3

NET CONTENTS

WINCO

No. 2 E. C.

ACTIVE INGREDIENTS:

*+(5-Benzyl-3-furyl)methyl 2, 2-dimethyl-3-(2-methylpropenyl)cyclopropanecarboxylate	2.000% w/w
Related compounds	0.274%
INERT INGREDIENTS:	97.726%
	<u>100.000% w/w</u>

* Cis. trans isomers ratio: max. 30% (-) cis and min. 70% (+) trans.
+ U. S. Patent No. 3,465,007.

USE ONLY AS DIRECTED
WARNING
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of feed and food-stuffs. Remove pets, birds and cover fish aquariums before spraying. Do not use on edible crops.

STORE AT TEMPERATURES ABOVE 40°
MIX WELL BEFORE USING.

Manufactured by
WINCO CHEMICAL COMPANY, INC.

P. O. Box 10692
JACKSON, MISSISSIPPI 39209

FOOD AREAS: Do not use in edible products areas of food processing plants, restaurants or other areas while food is being commercially prepared or processed. Do not use in serving areas while food is exposed.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy container by perforating or crushing, and bury in a safe place away from water.

OUTDOORS—TO CONTROL NUISANCE PESTS: Apply as a coarse contact spray. Spray directly on insects.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FLEAS: Same as directed for indoor use.

SPIDERS: Direct short burst of spray on spiders and repeat as necessary.

WASPS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

E.P.A. Reg. No. 35138-3 E.P.A. Est. No. 35138-MS-1

adding 1 fl. oz. of
fl. oz. of concentrate

Apply as a space
SBP-1382-0. 25%
deliver a fine mist
the spray mist to
which attract these
fabrics to avoid
of one fluid ounce
the room closed for
pleted and before

ect spray. Mix well
% insecticide spray
face to be treated.
le size larger than
roaches, earwigs,
s. Spray thoroughly
s around pipes and
ly spray until area
cts where possible.

ants where possible

resting areas. Hold
pray at the rate of
floor around these
face of dog houses

they re-enter the

TED

1975

135