

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid contact with eyes. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Vacate treated areas and ventilate before reoccupying. Remove pets, birds and cover fish aquariums before spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its label.

**RE-ENTRY STATEMENT**

Vacate treated areas and ventilate before reoccupying.

**STORAGE AND DISPOSAL**

Do not reuse empty container. Wrap container and put in trash collection.

**DIRECTIONS FOR USE**


**ON ANIMALS:** To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

**OUTDOOR ADULT MOSQUITO CONTROL:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may nest, low marshy places, stagnant pools or ponds and shore lines. Apply at a rate of 1 1/2 gallons per acre (4 1/2 ounces per 1000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist® models 51C, 91 or 100 HT traveling at 5 m.p.h. to treat a 300 foot swath calibrate equipment to deliver 4.5 gallons per minute.

**BATCH**



**GENERAL PURPOSE INSECTICIDE**

For Use in Homes, Institutions, Restaurants  
Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control  
Apply with Conventional Mechanical or Compressed Air Equipment

<b>ACTIVE INGREDIENTS:</b>	
Pyrethrins.....	0.1%
*Piperonyl Butoxide, Technical.....	1.0%
Petroleum Distillate.....	0.4%
<b>INERT INGREDIENTS:</b> .....	98.5%
	100.0%

\*Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**  
(See side panel for additional precautionary statements)

Manufactured By: Winco Chemical Company, Inc.  
Jackson, Mississippi

EPA ESTABLISHMENT NO.: 35138-MS-1  
EPA REGISTRATION NO.: 35138-38  
Net Contents

**DIRECTIONS FOR USE  
INDOOR APPLICATION**

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

**SURFACE SPRAYING:** To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spiny beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and beetles: Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

**To control carpet beetles:** Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

**To control bedbugs:** Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

**To control fleas and brown dog ticks:** Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, 60-70% behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

**To control clothes moths:** Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

**SPACE SPRAYING:** Close doors and windows and shut off ventilating systems. Apply with a fog machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, stinger flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths: Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 100 cubic feet of space.