

treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after applying this product, however, stop watering before puddling occurs.

(d) Relocate the lawn spreader setting table from page "15" to page "4", to the end of the overall lawn application directions, i.e., after the specific directions for Hyperodes weevil control (and above the "Spot Treatment" section). When relocating this table, the following revisions should be made. Since the table will then be near the pest listing for overall lawn treatment, delete "Insects listed below" from the top of the righthand column and delete the list of pests from the bottom of the table. Also, revise the table heading to read "SPREADER SETTING NO. (To deliver 2 lbs. per 1000 sq. ft.)". Then, in the right column heading, delete "2 lbs" from beneath "Grass or Dichondra lawn".

(e) In the directions for "Band Treatment Around House Foundation", revise the sentence under the pest listing to read "Apply a 5 foot band next to the house foundation."

(f) Delete the directions for control of lesser cornstalk borers on beans. Ciba-Geigy has elected not to support reregistration of post-emergence application of granular diazinon to food crops.

(g) Delete the directions for control of carrot rust flies on carrots. Ciba-Geigy has elected not to support reregistration of in-furrow application of granular diazinon to food crops.

~~(h) For the crop "melons", under the heading "CROP", relocate the closing parenthesis mark to appear after "... and hybrids of these" rather than after "... and hybrids of these and watermelons".~~

(i) For the crop "onions", revise the "TIMING/SPECIAL DIRECTIONS" to read:
Broadcast and mix into the top 3-4 inches of soil just before planting.

NOTE: Diazinon may not control organophosphate-resistant onion maggots.

(j) Based on the results of avian field studies, Ciba-Geigy is not supporting post-emergence application of granular diazinon to corn. Therefore, on page "13", delete "Sweet Corn (Cont'd)", delete the pest claims and associated directions for control of European corn borers, Fall armyworms, and Southwestern corn borers, and delete the "Notes", which deal with picking corn immediately after last application and prohibiting feeding fodder to livestock for 10 days following the last application. Neither of these notes is appropriate to the remaining at-plant soil application uses on sweet corn.

4 of 20

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

5920

AEROCHEM

5% DIAZINON LAWN GRANULES

TREATS UP TO 5000 SQUARE FEET

lawn & garden
insect
control

This product must not be used
on golf courses and sod farms.

Ants
Armyworms
Bermudagrass mites
Billbugs
Brown dog ticks
Chiggers
Chinch bugs
Crickets
Cutworms

Earwigs
Fire ants*
Fleas
Onion maggots
Root maggots
Sod webworms
White grubs of:
 . Japanese beetle
 . European chafer
 . Southern chafer
Wireworms
and others

*Aids in control.

See Directions for Use.

Ready-to-use Granules. Contains
5% diazinon insecticide.

Active Ingredient:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl) phosphorothioate

Inert Ingredients:

Total:

5%

95%

100%

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See additional precautionary statements
on back of bag.

EPA REG. NO. 35138

EPA EST. 35138-MS-1

Ten Pounds
Net Weight

MFG. BY: AEROCHEM, INC.
P.O. BOX 1089
RAYMOND, MS 39154

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 19 1990

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No. 35138-62

(k) For the crop "sweet potatoes", delete the "TIMING/SPECIAL DIRECTIONS" pertaining to post-emergence application. since, based upon the results of their avian field studies, Ciba-Geigy has elected not to support reregistration of granular diazinon for foliar or unincorporated soil surface application. Specifically, delete: "Southeastern U.S.- Broadcast 10 1/2 oz. per 500 sq. ft. over the tops of vines when roots begin to enlarge and do not mix into the soil."

(l) Revise the "Statement of Practical Treatment" and the associated "Note to Physician" to read as follows:

If swallowed - Call physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If inhaled - Remove victim to fresh air and apply artificial respiration if necessary.

If on skin - Wash promptly with plenty of soap and water. Rinse thoroughly.

If in eyes - Rinse eyes with plenty of water and call a physician immediately.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal, and may be administered, but only in conjunction with atropine.

(m) In the Environmental Hazards section, add a comma after "especially waterfowl" in the second sentence. You may elect to consolidate this entire section into a single paragraph instead of the current three paragraphs. Alternatively, you may choose to divide this section into two paragraphs, the second paragraph would only deal with aquatic hazards and would begin "Keep out of lakes, streams ..." and would contain the current last two paragraphs.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling. (NOTE: When preparing your label, do not print the quotation marks that appear around revised text items specified in this Notice of Registration.

In Lawns
kills:

- Ants
- Armyworms
- Bermudagrass mites
- Billbugs
- Brown dog ticks
- Chiggers
- Chinch bugs
- Clover mites
- Crickets
- Cutworms
- Digger wasps
- Earwigs
- Fleas
- Fire ants*
- Lawn moths
- Leafhoppers
- Millipedes
- Sod webworms
- Sowbugs
- Springtails
- White grubs of:
 - . European chafer
 - . Japanese beetle
 - . Southern chafer

*Aids in control.

In Gardens
kills:

- Armyworms
- Carrot rust flies ✓
- Corn borers
- Cutworms
- Mole crickets
- Onion maggots
- Root maggots
- Rootworms/
- Wireworms

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Overall Lawn Application:

Controls the following lawn insects on both grass and dichondra lawns:

- Ants . Armyworms . Bermudagrass mites . Brown dog ticks .
- Chiggers . Chinch bugs . Clover mites . Crickets . Cutworms .
- Earwigs . Fleas . Lawn moths (sod webworms) . Leafhoppers .
- Millipedes . Sowbugs . Springtails (Collembola)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds. Avoid overlapping treated areas. If waterfowl, i.e., ducks, geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 0.25 inch water immediately after application, however, stop watering before puddling occurs.

Apply with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10').

Billbugs; White grubs of Japanese beetle, European chafer, Southern chafer, Ataenius spretulus (dung beetle):

Apply 2 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late Summer or early Fall if a second generation occurs.

Hyperodes weevils:

Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Spot Treatment

Ants: Apply 1 1/2 teaspoons of over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 2 ft. around each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.

Digger wasps: Apply 1 1/2 teaspoons of in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation:

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses.

- | | |
|-----------------|--------------------|
| Ants | Fleas |
| Brown dog ticks | Millipedes |
| Clover mites | Sowbugs (Pillbugs) |
| Crickets | Springtails |
| Earwigs | (Collembola) |

Apply a 5 foot band of next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. Repeat as necessary.

Vegetables - Insect Control

Controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Unless directions under "Timing/Special Directions" state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

| <u>Insect</u> | <u>Depth of Mixing or Incorporation into Soil</u> |
|------------------------|---|
| Surface cutworms* | 2-3 inches |
| Subterranean cutworms* | 3-6 inches |
| Mole crickets | 1-2 inches |
| Root maggots | 3-4 inches |
| Wireworms | 4-8 inches |

*If type of cutworm cannot be identified, mix into soil 4-6 inches deep.

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---|--|--|---|
| Beans (Pole, Snap or Lima) (succulent only) | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Lesser Cornstalk Borers | 3 1/2-7 ozs. per 500 sq. ft. | Apply in a 6-inch band centered over the row when plants are 1/2-1 inch tall. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc. | | |
| Beets, Red (Table) | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Moles Crickets | 3-1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Broccoli | Root Maggots | 7-10 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting or transplanting. |
| | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|----------------------|------------------|-----------------------------------|--|
| Broccoli (Cont'd) | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Brussels Sprouts | Root Maggots | 7-10 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting or transplanting. |
| | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Cabbage | Root Maggots | 7-10 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting or transplanting. |
| | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------|----------------------|--------------------------------|--|
| Carrots | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Carrot Rust Flies | 7 ozs. per 500 sq. ft. | Apply in the seed furrow at planting time. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Cauliflower | Root Maggots | 7-10 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting or transplanting. |
| | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Celery | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------|---------------|--------------------------------|---|
| Collards | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Cucumber | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Endive (Escarole) | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Kale | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--|---------------|--------------------------------|---|
| Lettuce | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Melons (Canta-loupes, Casabas, Crenshaws, Honeydew, Musk-melons, Persians, & Hybrids of these and water-melons) | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Mustard | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| Onions (green and bulb) | Onion Maggots | 7-14 ozs. per 500 sq. ft. | Broadcast just before planting and mix into the top 3-4 inches of soil by disking, harrowing, or other suitable means. NOTE: Diazinon may not control organophosphate-resistant onion maggots. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---|--|--------------------------------|---|
| Parsley | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Peas (succulent only) | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| NOTE: Do not use on dried pea varieties such as dried blackeyed peas. | | | |
| Peppers | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Potatoes | Cutworms (Surface & Subter- ranean) | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| NOTE: Do not use on commercially-grown potatoes which will be hand-harvested. | | | |

15 9 90

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---------------------------|-------------------|--------------------------------|---|
| Radishes | Root Maggots | 5 1/2-7 ozs. per 500 sq. ft. | Apply in the seed furrow at planting (2 1/2-2 4/5 ozs. per 1,000 linear feet of row). |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| Spinach | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Squash, Summer and Winter | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Sweet Corn | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Seed-Corn Maggots | 7-14 ozs. per 500 sq. ft. | Broadcast just before planting. Mix into soil 2 inches |
| | Wireworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|------------------------|-------------------------------------|---------------------------------|---|
| Sweet Corn (Cont'd) | European Corn Borers | 3 1/2-7 ozs. per 500 sq. ft. | When 75% of the plants show first-generation borers feeding, apply over the corn so granules fall into whorls. Make a second application 7 days later if needed.* For second-generation borers, apply as above when there is an average of 1 egg mass per plant. Do not make more than 3 applica- tions per season.* |
| | Fall Armyworms | 3 1/2-7 ozs. per 500 sq. ft. | Apply over corn so granules fall into whorls when larvae first appear. If necessary, applica- tion may be repeated twice at 7-day interval. Do not make more than 3 applications per season.* |
| | South- western Corn Borers | 3 1/2-7 ozs. per 500 sq. ft. | Apply over corn so granules fall into whorls when second- generation larvae first appear. If necessary, application may be repeated at 7-day intervals. Do not make more than 3 applica- tions per season.* |

*NOTES: Corn may be picked immediately after last application.
Do not feed fodder to livestock for 10 days following
last application.

| | | | |
|-------------------|-------------------------------|--------------------------------|--|
| Sweet Potatoes | Wireworms, Flea Beetles | 10 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. Southeastern U.S. - Broadcast 10 1/2 ozs. per 500 sq. ft. over the tops of vines when roots begin to enlarge and do not mix into soil. |
|-------------------|-------------------------------|--------------------------------|--|

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------|---------------|--------------------------------|---|
| Swiss Chard | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Tomatoes | Cutworms | 7-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |
| Turnips | Mole Crickets | 3 1/2 ozs. per 500 sq. ft. | Broadcast and mix into soil 1-2 days before planting. |
| | Wireworms | 10 1/2-14 ozs. per 500 sq. ft. | Broadcast and mix into soil just before planting. |

18 9 20

SPREADER SETTING NO.
(per 1000 sq. ft.)

| Spreader Make and Model | Insects listed below† |
|---|-----------------------------------|
| | Grass or Dichondra Lawn 2 lbs. |
| Central Imperial Mark No. 1662-G | 6 1/2 |
| Central Suburbia Lawn Spreader - No. 1622-0 | 6 1/2 |
| Cyclone Model B1 Rotary | 4 1/2 |
| Greenfield Model 105 | 6 1/2 |
| ORTHO Drop Spreader Model 3000 | 4 1/2 |
| ORTHO Broadcast Spreader Model 4000 | 4 1/2 |
| J. C. Penney Model 8005 | 7 1/4 |
| J. C. Penney Rotary Spreader | 5 |
| Scott Precision Flow Spreader Model PF-4 | 4 |
| Sears Craftsman Model 671.19198 | 6 |

- | | | |
|--------------------|------------------|----------------|
| †Ants | Clover Mites | Lawn Moths |
| Armyworms | Crickets | (Sod Webworms) |
| Bermudagrass Mites | Cutworms | Leafhoppers |
| Brown Dog Ticks | Earwigs | Millipedes |
| Chiggers | Fleas | Sowbugs |
| Chinch Bugs | Hyperodes Weevil | (Pill Bugs) |
| | | Springtails |
| | | (Collembola) |

Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. May cause contact sensitization following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

Statement of Practical Treatment

If swallowed, remove victim to fresh air and apply artificial respiration if indicated.

If on skin, wash promptly with soap and water. Rinse thoroughly.

If in eyes, rinse eyes with plenty of water and call a physician immediately.

Continued

Note to Physician

This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 0.25 inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries.

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.