

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
(and Domestic Animals)**

**DANGER**

**HAZARDOUS LIQUID AND VAPOR  
MAY BE FATAL IF INHALED,  
ABSORBED THROUGH SKIN, OR SWALLOWED  
CAUSES BURNS OF SKIN OR EYES  
COMBUSTIBLE**

Do Not Get in Eyes, on Skin, or on Clothing. Wear Eye and Skin Protection Necessary to Prevent Contact When Handling This Material. Wash Thoroughly after Handling and Before Eating and Smoking. Wash Contaminated Clothing and Air Contaminated Shoes Thoroughly Before Reuse. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Use Only with Adequate Ventilation. Do Not Breathe Vapor.

Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals. Do not contaminate food, foodstuffs, or domestic water supplies.

**ENVIRONMENTAL HAZARDS**

To avoid injury to fish and other wildlife, do not spill or empty TELONE II into streams, ponds, or other bodies of water.

**PHYSICAL OR CHEMICAL HAZARDS: FLAMMABLE LIQUID**

**DIRECTION FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THIS LABELING

**STORAGE AND DISPOSAL**

Rinse equipment and containers and dispose of wastes by burying in non crop lands away from water supplies. Dispose of empty containers by punching holes in them and burying with wastes.

**USE PRECAUTIONS**

**Important — Note Carefully.** Fumigation may temporarily raise the level of ammonia nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acid, or high in organic matter. To avoid injury to plant roots, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation, or both, to crops on high organic soils do not use fertilizers containing ammonium salts and use only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65 F.

Certain crops including cotton, sugarcane, and pineapple are tolerant to ammonia and the above rule does not apply to them. When using high rates of TELONE II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Cornus* sp., *Crataegus* sp., spruce, and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect it is suggested that additional phosphate fertilizer be used on soils where experience indicates a deficiency may occur.

**Attention:** To avoid reinfestation of treated soil do not use irrigation water, transplants, tools, seed pieces, or crop remains that could carry soilborne pests from infested land. Clean equipment carefully before using.

Since TELONE II soil fumigant is corrosive under certain conditions flush all applicators with fuel oil or kerosene immediately after use. DO NOT USE WATER. Do not use containers, pumps, or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions TELONE II may be severely corrosive to such metals. Common protective clothing, such as rubber gloves and boots, etc., may be penetrated readily by this

# TOXO TELONE II

## SOIL FUMIGANT

A CLEAN, CLEAR, NON-CLOGGING LIQUID FOR PREPLANT TREATMENT OF CROP LANDS TO CONTROL PLANT PARASITIC NEMATODES AND CERTAIN OTHER SOIL PESTS.

**ACTIVE INGREDIENT:**

1,3-Dichloropropene ..... 92%

INERT INGREDIENTS: ..... 8%

TOTAL ..... 100%

## KEEP OUT OF REACH OF CHILDREN DANGER

**STATEMENT OF PRACTICAL TREATMENT**

**In case of contact,** immediately remove contaminated clothing and shoes and wash skin with soap and plenty of water; for eyes, flush with flowing water for at least 15 minutes and get medical attention.

**In case of spillage indoors** have available an approved full face mask equipped with a fresh black canister meeting specifications of the U.S. Bureau of Mines for organic vapors, a full face self contained breathing apparatus, or full face air supplied respirator.

**If inhaled,** remove to fresh air. Use artificial respiration if breathing has stopped. Get immediate medical attention. Keep patient absolutely quiet.

**If swallowed,** call a doctor. Induce vomiting by giving an emetic such as 2 tablespoonfuls of table salt in a glass of warm water.

See side panel for additional precautionary statements

MFG. FOR: TOXO SPRAY-DUST, INC.  
TUSTIN, CALIFORNIA

**DIRECTIONS FOR USE**

**WHEN TO TREAT:** Apply TELONE II either in the spring conditions permit. For best results with annual crops, treat in late summer, or early fall treatment before October to early spring crops, especially transplanted crops such as and orchard stock. Early fall treatment permits planting muck soils only in the early fall and plant as late as possible (clay) soils only when they are near or at the wilting point of any type or soil when it is cold and/or wet.

**SOIL PREPARATIONS:** TELONE II gives best results with fusion of the fumigant through the soil and the soil surface. Avoid fumigant loss during the exposure period. The soil should be free of clods and undecomposed plant material, moisture at application temperature between 40° and 80° F at the depth of injection. If present, it should be plowed down and allowed to decompose. Tillage deeper than 12 inches is necessary for good fumigation. Harrow or "plow" pan occurs at plow depth. Where deep tillage may often be modified to simultaneously apply the fumigant to the field twice.

**APPLICATION:** TELONE II may be applied either as overall treatment, using suitable application equipment that will place at least 6 to 8 inches below the final soil surface. For overall treatment use chisel equipment with the chisels spaced 12 inches apart, a depth of 12 inches or more (deep tillage), the chisel application depth but should not exceed 30 inches. Apply in the direction or at an angle to the direction of the planting row. For row application, use chisel equipment with one chisel 12 inches apart to treat only the soil where the crop is to be planted. When used, adjust the fumigant flow rate to distribute about 1.1/3 lbs per acre for overall application. When two chisels are used, the same flow rate per chisel as for overall. In both cases, the flow rate will decrease as the distance between rows is increased. Avoid placing the seed row directly over the fumigant. When a single chisel is used per row, place the seed 3 to 4 inches from the furrow. When two chisels are used, plant in the center of the furrow.

**SEALING:** Immediately after application compact the soil to prevent fumigant loss. After chisel application, use a roller, cultipacker, or disk. After plow sole application, disk the land, then compact it by row application can be accomplished by the tractor wheel, the fumigant will be 12 to 14 inches below the top of the bed. Seal in the fumigant with ring rollers, press sealers, or by re-

**EXPOSURE PERIOD:** After application and sealing, leave the soil for 7 to 10 days. A longer exposure period will be required if the soil is not sealed during the exposure period.

**AERATION AND PREPARATION OF SOIL BEFORE PLANTING:** After the exposure period allow the soil to aerate completely before planting. Complete when the odor of TELONE II is no longer evident. Under conditions, allow one week of aeration time for each 10 gal. of TELONE II used for treating deep rooted trees. For trees, a 30-day aeration period should be allowed. To hasten aeration at low temperatures occur during the exposure period, work the soil. After row treatment use a knife-like chisel in the furrow, reducing possible recontamination of the treated soil. To hasten aeration, plow or deep cultivate to the depth of the treated zone. In northern states after fall fumigation or after row treatment, the soil should be plowed or disked before planting.

**White Potatoes in Northwestern States:** Use TELONE II as a treatment to control quarantine and for suppression of the Colorado potato beetle. Apply as an overall treatment. Apply as an overall treatment.

Time of Treatment	Gallons per Acre
Spring	17 to 25
Fall	25 to 34

**Air in Northwestern States:** Use 39 gallons of TELONE II per acre as an overall treatment in the spring or pre-planting of the damaging effects of Verticillium wilt in disk and plow. After treatment allow at least 7 to 8 weeks before planting. Consult State Agricultural Experiment Station Service Specialists for the use of other practices for control and cultural practices when using TELONE II as an overall treatment.

**NOTICE:** Seller warrants that the product conforms to the specifications for the purposes stated on the label when used under normal conditions of use, but neither this warranty nor the seller's liability for fitness for a particular purpose or for the use of this product contrary to label instructions or under conditions not reasonably foreseeable to seller, shall be such use.

**CAUSES BURNS OF SKIN OR EYES  
COMBUSTIBLE**

Do Not Get in Eyes, on Skin, or on Clothing. Wear Eye and Skin Protection Necessary to Prevent Contact When Handling This Material. Wash Thoroughly after Handling and Before Eating and Smoking. Wash Contaminated Clothing and Air Contaminated Shoes Thoroughly Before Reuse. DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Use Only with Adequate Ventilation. Do Not Breathe Vapor.

Do not allow contamination of seeds, plants, fertilizers, or other pesticide chemicals. Do not contaminate food, foodstuffs, or domestic water supplies.

**ENVIRONMENTAL HAZARDS**

To avoid injury to fish and other wildlife, do not spill or empty TELONE II into streams, ponds, or other bodies of water.

**PHYSICAL OR CHEMICAL HAZARDS: FLAMMABLE LIQUID**

**DIRECTION FOR USE**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH THIS LABELING

**STORAGE AND DISPOSAL**

Rinse equipment and containers and dispose of wastes by burying in non crop lands away from water supplies. Dispose of empty containers by punching holes in them and burying with wastes.

**USE PRECAUTIONS**

**Important — Note Carefully.** Fumigation may temporarily raise the level of ammonia nitrogen and soluble salts in the soil. This is most likely to occur when heavy rates of fertilizer and fumigant are applied to soils that are either cold, wet, acid, or high in organic matter. To avoid injury to plant roots, fertilize as indicated by soil tests made after fumigation. To avoid ammonia injury or nitrate starvation, or both, to crops on high organic soils do not use fertilizers containing ammonium salts and use only fertilizers containing nitrates, until after the crop is well established and the soil temperature is above 65 F.

Certain crops including cotton, sugarcane, and pineapple are tolerant to ammonia and the above rule does not apply to them. When using high rates of TELONE II as required by certain state nursery regulations, liming of highly acid soils before fumigation may stimulate nitrification and reduce the possibility of ammonia toxicity. Certain nursery crops such as citrus seedlings, *Cornus sp.*, *Crataegus sp.*, spruce, and vegetable crops such as cauliflower have shown evidence of phosphorus deficiency following fumigation. To avoid this possible effect, it is suggested that additional phosphate fertilizer be used on soils where experience indicates a deficiency may occur.

**Attention:** To avoid reinfestation of treated soil do not use irrigation water, transplants, tools, seed pieces, or crop remains that could carry soilborne pests from infested land. Clean equipment carefully before using.

Since TELONE II soil fumigant is corrosive under certain conditions flush all applicators with fuel oil or kerosene immediately after use. DO NOT USE WATER. Do not use containers, pumps, or other transfer equipment made of aluminum, magnesium or their alloys, as under certain conditions TELONE II may be severely corrosive to such metals. Common protective clothing, such as rubber gloves and boots, etc., may be penetrated readily by this material. Polyethylene provides a good barrier. For field operations cover shoes and hands with polyethylene bags. Store TELONE II in tightly closed containers in a cool place away from dwellings. In outside storage, store drums on their sides to avoid accumulation of rain water in the top or bottom recessed areas.

**ACCEPTED**  
FEB 19 1976  
Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended by the Pesticide Control Act of 1972, this pesticide is registered under EPA Reg. No. 35296-1

**TELONE II  
SOIL FUMIGANT**

A CLEAN, CLEAR, NON-CLOGGING LIQUID FOR PREPLANT TREATMENT OF CROP LANDS TO CONTROL PLANT PARASITIC NEMATODES AND CERTAIN OTHER SOIL PESTS.

**ACTIVE INGREDIENT:**

1,3-Dichloropropene ..... 92%

INERT INGREDIENTS: ..... 8%

TOTAL ..... 100%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

**In case of contact,** immediately remove contaminated clothing and shoes and wash skin with soap and plenty of water; for eyes, flush with flowing water for at least 15 minutes and get medical attention.

**In case of spillage indoors** have available an approved full face mask equipped with a fresh black canister meeting specifications of the U.S. Bureau of Mines for organic vapors, a full face self-contained breathing apparatus, or full face air-supplied respirator.

**If inhaled,** remove to fresh air. Use artificial respiration if breathing has stopped. Get immediate medical attention. Keep patient absolutely quiet.

**If swallowed,** call a doctor. Induce vomiting by giving an emetic such as 2 tablespoonfuls of table salt in a glass of warm water.

See side panel for additional precautionary statements

MFG. FOP TOXO SPRAY-DUST, INC.  
TUSTIN, CALIFORNIA

EPA EST. NO. 35296-CA-1  
EPA NO. 35296-1  
NET CONTENTS \_\_\_\_\_ GALS.

\*Trademark of Dow Chemical Company

Tillage deeper than 12 inches is necessary for good hard or "plow" pan occurs at plow depth. Where deep may often be modified to simultaneously apply the fur field twice

**APPLICATION:** TELONE II may be applied either treatment, using suitable application equipment that will be applied at least 6 to 8 inches below the final soil surface. For row or chisel equipment with the chisels spaced 12 inches apart a depth of 12 inches or more (deep tillage), the chisel application depth but should not exceed 30 inches. Apply in direction or at an angle to the direction of the planting. For row application, use chisel equipment with one chisel inches apart to treat only the soil where the crop is to be used, adjust the fumigant flow rate to distribute about 1/2 acre than is recommended for overall application. When two same flow rate per chisel as for overall. In both cases, 1/2 acre will decrease as the distance between rows is increased. When planting, avoid placing the seed row directly over the furrow. When a single chisel is used per row place the seed in the center of the furrow. When two chisels are used, plant in the center of the furrow.

**SEALING:** Immediately after application, compact the soil to prevent fumigant loss. After chisel application, use a roller, cultipacker, or similar equipment. After plow-sole application, disk the land, then compact the soil. Row application can be accomplished by the tractor with the fumigant will be 12 to 14 inches below the top of the soil. Seal in the fumigant with ring rollers, press sealers, or similar equipment.

**EXPOSURE PERIOD:** After application and sealing, the soil should be kept moist during the exposure period.

**AERATION AND PREPARATION OF SOIL BEFORE PLANTING:** After exposure period allow the soil to aerate completely before planting. When the odor of TELONE II is no longer evident, allow one week of aeration time for each 1/2 acre. When TELONE II is used for treating deep rooted crops, a longer aeration period should be allowed. To hasten aeration, use a low tillage implement during the exposure period, or use a low tillage implement during the exposure period, or use a low tillage implement during the exposure period. After row treatment use a knife like chisel in the row zone. After row treatment use a knife like chisel in the row zone. After row treatment use a knife like chisel in the row zone. After row treatment use a knife like chisel in the row zone.

**White Potatoes in Northwestern States:** Use TELONE II treatment to control quackgrass and for suppression of weeds in fields to be planted to white potatoes. Apply as a following directions

Time of Treatment	Gallons per Acre
Spring	17 to 25
Fall	25 to 34

**Mint in Northwestern States:** Use 59 gallons of TELONE II per chisel as an overall treatment in the spring, or reduction of the damaging effects of Verticillium wilt in mint production. After treatment allow at least 7 to 8 weeks for the soil before planting. Consult State Agricultural Extension Service Specialists for the use of other practical control and cultural practices when using TELONE II by Verticillium wilt.

**NOTICE:** Seller warrants that the product conforms reasonably fit for the purposes stated on the label when used under normal conditions of use, but neither this warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE nor the use of this product contrary to label instructions or under conditions not reasonably foreseeable to seller.

**GENERAL INFORMATION**

Use TELONE II soil fumigant only as a preplanting soil fumigant to control nematodes (rootknot, meadow lesion), citrus, cyst, burrowing, ring, spiral, sting, pin, stubby root, stylet, and garden centipedes (symphylans). Fumigate land by applying TELONE II under the conditions and DIRECTIONS FOR USE, DOSAGE AND USE PRECAUTIONS. Read the entire label before using TELONE II.

**Vegetable Crops:**

asparagus	cauliflower	horseradish
beans	celery	kale
beets	collards	kohlrabi
blackeyed peas	corn	leeks
broccoli	cowpeas	lettuce
brussels sprouts	cucumbers	melons
cabbage	egg plant	mustard greens
cantaloupe	endive	okra
carrots	garlic	onions

# TOXO TELONE II

## SOIL FUMIGANT

A CLEAN, CLEAR, NON-CLOGGING LIQUID FOR PREPLANT TREATMENT OF CROP LANDS TO CONTROL PLANT PARASITIC NEMATODES AND CERTAIN OTHER SOIL PESTS.

### ACTIVE INGREDIENT:

1,3-Dichloropropene ..... 92%

INERT INGREDIENTS ..... 8%

TOTAL ..... 100%

## KEEP OUT OF REACH OF CHILDREN DANGER

### STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately remove contaminated clothing and shoes and wash skin with soap and plenty of water; for eyes, flush with flowing water for at least 15 minutes and get medical attention.

In case of spillage indoors have available an approved full face mask equipped with a fresh black canister meeting specifications of the U. S. Bureau of Mines for organic vapors, a full face self-contained breathing apparatus, or full face air supplied respirator.

If inhaled, remove to fresh air. Use artificial respiration if breathing has stopped. Get immediate medical attention. Keep patient absolutely quiet.

If swallowed, call a doctor. Induce vomiting by giving an emetic such as 2 tablespoonfuls of table salt in a glass of warm water.

See side panel for additional precautionary statements

MFG. FOR: TOXO SPRAY-DUST, INC.  
TUSTIN, CALIFORNIA

### DIRECTIONS FOR USE

**WHEN TO TREAT:** Apply TELONE II either in the spring or fall, whenever soil type and conditions permit. For best results with annual crops, treat the soil each year. In northern states, late summer, or early fall treatment (before October 15) is best for land to be planted to early spring crops, especially transplanted crops such as celery, tomatoes, and nursery and orchard stock. Early fall treatment permits planting a fall cover crop. **NOTE:** Treat muck soils only in the early fall and plant as late as possible in the spring; treat fine textured (clay) soils only when they are near or at the wilting point. Do not use TELONE II to treat any type of soil when it is cold and/or wet.

**SOIL PREPARATIONS:** TELONE II gives best results when conditions permit rapid diffusion of the fumigant through the soil and the soil surface can be sealed to prevent excessive fumigant loss during the exposure period. The soil should be in good seed bed condition, free of clods and undecomposed plant material, moisture at about one-half of field capacity, and temperature between 40° and 80°F at the depth of injection. If undecomposed plant debris is present, it should be plowed down and allowed to decompose before applying TELONE II. Tillage deeper than 12 inches is necessary for good fumigant penetration in soils where a hard or "plow" pan occurs at plow depth. Where deep tillage is used, the tillage equipment may often be modified to simultaneously apply the fumigant and thus avoid going over the field twice.

**APPLICATION:** TELONE II may be applied either as an overall (broadcast) or row treatment, using suitable application equipment that will ensure placement of the fumigant at least 6 to 8 inches below the final soil surface. For overall application use either plowsole or chisel equipment with the chisels spaced 12 inches apart. When the fumigant is injected at a depth of 12 inches or more (deep tillage), the chisel spacing may be up to twice the application depth but should not exceed 30 inches. Application may be made in the same direction or at an angle to the direction of the planting row, whichever is most convenient. For row application, use chisel equipment with one chisel per row or two chisels spaced 12 inches apart to treat only the soil where the crop is to be planted. When one chisel per row is used, adjust the fumigant flow rate to distribute about 1.5 times the amount of fumigant per chisel than is recommended for overall application. When two chisels are used per row, apply at the same flow rate per chisel as for overall. In both cases, the amount of fumigant required per acre will decrease as the distance between rows is increased and vice versa. At time of planting, avoid placing the seed row directly over the furrow left by the applicator chisel. When a single chisel is used per row, place the seed 3 to 4 inches to one side of the chisel furrow. When two chisels are used, plant in the center of the area between the chisel furrows.

**SEALING:** Immediately after application, compact the soil surface to prevent excessive fumigant loss. After chisel application, use a roller, cultipacker, or similar sealing device. After plow sole application, disk the land, then compact it by floating or rolling. Sealing after row application can be accomplished by the tractor wheel, by listing, or by bedding so that the fumigant will be 12 to 14 inches below the top of the bed. When fumigating listed rows, seal in the fumigant with ring rollers, press sealers, or by re-listing.

**EXPOSURE PERIOD:** After application and sealing, leave the soil undisturbed for 7 to 14 days. A longer exposure period will be required if the soil becomes excessively cold or wet during the exposure period.

**AERATION AND PREPARATION OF SOIL BEFORE PLANTING:** At the end of the exposure period allow the soil to aerate completely before planting the crop. Aeration is usually complete when the odor of TELONE II is no longer evident. Under optimum soil and weather conditions, allow one week of aeration time for each 10 gallons of TELONE II applied per acre. When TELONE II is used for treating deep rooted tree and shrub planting sites, a 3 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rains or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife like chisel in the bed without turning the soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

**White Potatoes in Northwestern States:** Use TELONE II as a spring or preferably a fall treatment to control quackgrass and for suppression of the damaging effects of Verticillium wilt in fields to be planted to white potatoes. Apply as an overall treatment according to the following directions:

Time of Treatment	Gallons per Acre	Fl. Oz. /1000 Ft. Row per Chisel
Spring	17 to 25	50 to 73
Fall	25 to 34	73 to 100

**Mint in Northwestern States:** Use 59 gallons of TELONE II per acre (173 fl. oz. /1000 ft. row per chisel) as an overall treatment in the spring, or preferably in the fall, to aid in the reduction of the damaging effects of Verticillium wilt in disease infested land to be used for mint production. After treatment allow at least 7 to 8 weeks or until the odor of the fumigant has left the soil before planting. Consult State Agricultural Experiment Station or State Extension Service Specialists for the use of other practices such as flaming the stubble, weed control, and cultural practices when using TELONE II as an aid to reducing damage caused by Verticillium wilt.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE express or implied extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Field Crops:			
alfalfa	flax	oats	sorghum
barley	grasses	pasture grass	soybeans
birdsfoot trefoil	hops	peanuts	sugar beets
buckwheat	lespedeza	popcorn	sugarcane
clover	millet	rice	tobacco
corn	milc	rye	vetch
cotton	mint	safflower	wheat

  

Citrus Fruit Tree Planting Sites:					
grapefruit	kumquats	lemons	limes	oranges	tangerines
tangelos					

  

Deciduous Fruit and Nut Tree Planting Sites:			
almonds	dates	olives	plums
apples	figs	peaches	pomegranates
apricots	filberts	pears	prunes
cashew nuts	hazelnuts	pecans	quince
cherries	hickory nuts	persimmons	walnuts
chestnuts	nectarines	pineapple	

  

Bush and Vine Planting Sites:			
blackberries	currants	huckleberries	youngberries
blueberries	dewberries	loganberries	
boysenberries	gooseberries	raspberries	
cranberries	grapes	strawberries	

### DOSAGE AND USE RECOMMENDATIONS to Control Nematodes, Symphylans, and Wireworms†

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage		
			Gallons per Acre <sup>1</sup>	Fl. Oz./1000 Ft. Row Per Chisel	
Shallow Rooted Plants: Field Crops Floral Crops Grasses and Turf	Row (42") <sup>6</sup>	Mineral	4.5 to 6	46 to 62	
		Muck or Peat	9 to 12	93 to 123	
Small Fruits Vegetables Ornamentals	Overall (Broadcast)	Mineral	9 to 15 <sup>2</sup>	26 to 44	
		Muck or Peat	24 <sup>3</sup> to 36	71 to 106	
Strawberries	Overall	Mineral	24 to 36	71 to 106	
Sugar Beets Root-Knot Nematode	Row (42")	Mineral	6	62	
	Overall		12 to 15	35 to 44	
Sugar Beet Nematode	Row (42")	Mineral	9	93	
	Overall		12 to 18	35 to 53	
Pineapple <sup>4</sup>	Row	Mineral	24 to 36	—	
Citrus — Florida <sup>5</sup>	Overall	Mineral	36	106	
Nursery and Field: Citrus Fruit Trees Deciduous Fruit Trees <sup>6</sup> Forest Trees Grapes Nut Trees Ornamentals (Deep rooted)	Overall: Gallons per Acre to Penetrate Various Depths				
	Mineral Soils	3 ft.	4 ft.	5 ft.	6 ft.
	Sand	15	21	27	33
	Sandy Loam	27	30	36	48
	Silt Loam	42	51	63	75
Clay	54	69	84	102	

<sup>1</sup> Use the higher rates in heavier soil.

<sup>2</sup> For cyst forming nematodes increase dosage to 18 gallons per acre (53 fl. oz./1000 ft. row per chisel).

<sup>3</sup> For muck soils containing less than 30 percent organic matter use 18 gallons per acre.

<sup>4</sup> For Hawaiian pineapple, application may be made at time of or just before planting.

<sup>5</sup> For burrowing nematode in citrus inject on 18 inch centers, 12 inches deep. Keep free of plants susceptible to burrowing nematodes for 2 years before replanting to citrus.

<sup>6</sup> Row treatment is not recommended for potatoes in irrigated areas of western and northwestern states.

# TELONE II

## SOIL FUMIGANT

A CLEAN, CLEAR, NON-CLOGGING LIQUID FOR PREPLANT TREATMENT OF CROP LANDS TO CONTROL PLANT PARASITIC NEMATODES AND CERTAIN OTHER SOIL PESTS.

### ACTIVE INGREDIENT:

1,3-Dichloropropene ..... 92%

INERT INGREDIENTS: ..... 8%

TOTAL ..... 100%

## KEEP OUT OF REACH OF CHILDREN DANGER

### STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately remove contaminated clothing and shoes and wash skin with soap and plenty of water; for eyes, flush with flowing water for at least 15 minutes and get medical attention.

In case of spillage indoors have available an approved full face mask equipped with a fresh black canister meeting specifications of the U.S. Bureau of Mines for organic vapors, a full face self-contained breathing apparatus, or full face air-supplied respirator.

If inhaled, remove to fresh air. Use artificial respiration if breathing has stopped. Get immediate medical attention. Keep patient absolutely quiet.

If swallowed, call a doctor. Induce vomiting by giving an emetic such as 2 tablespoonfuls of table salt in a glass of warm water.

See side panel for additional precautionary statements

MFG. FOR: TOXO SPRAY-DUST, INC.  
TUSTIN, CALIFORNIA

EPA EST. NO. 35296-CA-1  
EPA NO. 35296-1  
NET CONTENTS \_\_\_\_\_ GALS.

\*Trademark of Dow Chemical Company

Tillage deeper than 12 inches is necessary for good fumigant penetration in soils where a hard or "plow" pan occurs at plow depth. Where deep tillage is used, the tillage equipment may often be modified to simultaneously apply the fumigant and thus avoid going over the field twice.

**APPLICATION:** TELONE II may be applied either as an overall (broadcast) or row treatment, using suitable application equipment that will ensure placement of the fumigant at least 6 to 8 inches below the final soil surface. For overall application use either plowsole or chisel equipment with the chisels spaced 12 inches apart. When the fumigant is injected at a depth of 12 inches or more (deep tillage), the chisel spacing may be up to twice the application depth but should not exceed 30 inches. Application may be made in the same direction or at an angle to the direction of the planting row, whichever is most convenient. For row application, use chisel equipment with one chisel per row or two chisels spaced 12 inches apart to treat only the soil where the crop is to be planted. When one chisel per row is used, adjust the fumigant flow rate to distribute about 1-1/3 times more fumigant per chisel than is recommended for overall application. When two chisels are used per row, apply at the same flow rate per chisel as for overall. In both cases, the amount of fumigant required per acre will decrease as the distance between rows is increased and vice versa. At time of planting, avoid placing the seed row directly over the furrow left by the applicator chisel. When a single chisel is used per row, place the seed 3 to 4 inches to one side of the chisel furrow; when two chisels are used, plant in the center of the area between the chisel furrows.

**SEALING:** Immediately after application, compact the soil surface to prevent excessive fumigant loss. After chisel application, use a roller, cultipacker, or similar sealing device. After plow sole application, disk the land, then compact it by floating or rolling. Sealing after row application can be accomplished by the tractor wheel, by listing, or by bedding so that the fumigant will be 12 to 14 inches below the top of the bed. When fumigating listed rows, seal in the fumigant with ring rollers, press sealers, or by re-listing.

**EXPOSURE PERIOD:** After application and sealing, leave the soil undisturbed for 7 to 14 days. A longer exposure period will be required if the soil becomes excessively cold or wet during the exposure period.

**AERATION AND PREPARATION OF SOIL BEFORE PLANTING:** At the end of the exposure period allow the soil to aerate completely before planting the crop. Aeration is usually complete when the odor of TELONE II is no longer evident. Under optimum soil and weather conditions, allow one week of aeration time for each 10 gallons of TELONE II applied per acre. When TELONE II is used for treating deep rooted tree and shrub planting sites, a 1 to 6 months aeration period should be allowed. To hasten aeration, especially if heavy rain, or low temperatures occur during the exposure period, work the soil to the depth of the treatment zone. After row treatment use a knife-like chisel in the bed with a turning soil, thus reducing possible recontamination of the treated soil. To hasten aeration after overall treatment, plow or deep cultivate to the depth of the treatment zone. This is especially desirable in northern states after fall fumigation of muck soils.

**White Potatoes in Northwestern States:** Use TELONE II as a spring or preferably a fall treatment to control quackgrass and for suppression of the damaging effects of Verticillium wilt in fields to be planted to white potatoes. Apply as an overall treatment according to the following directions:

Time of Treatment	Gallons per Acre	Fl. Oz. /1000 Ft. Row per Chisel
Spring	17 to 25	50 to 73
Fall	25 to 34	73 to 100

**Mint in Northwestern States:** Use 59 gallons of TELONE II per acre (173 fl. oz. / 1000 ft. row per chisel) as an overall treatment in the spring, or preferably in the fall, to aid in the reduction of the damaging effects of Verticillium wilt in disease infested land to be used for mint production. After treatment allow at least 7 to 8 weeks or until the odor of the fumigant has left the soil before planting. Consult State Agricultural Experiment Station or State Extension Service Specialists for the use of other practices such as flaming the stubble, weed control and cultural practices when using TELONE II as an aid to reducing damage caused by Verticillium wilt.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

### GENERAL INFORMATION

Use TELONE II soil fumigant only as a preplanting soil treatment to control plant parasitic nematodes (rootknot, meadow lesion), citrus, cyst formers (golden, sugar beet, soybean), burrowing, ring, spiral, sting, pin, stubby root, stylet, dagger); also to control wireworms and garden centipedes (symphylans). Fumigate land to be planted to the crops listed below by applying TELONE II under the conditions, and at the rates recommended under DIRECTIONS FOR USE, DOSAGE AND USE RECOMMENDATIONS, and PRECAUTIONS. Read the entire label before using TELONE II.

#### Vegetable Crops:

asparagus	cauliflower	horseradish	parsnips	shallots
beans	celery	kale	peas	spinach
beets	collards	kohlrabi	peppers	squash (summer)
blackeyed peas	corn	leeks	pimentos	squash (winter)
broccoli	cowpeas	lettuce	potatoes	sweet potatoes
brussels sprouts	cucumbers	melons	pumpkins	swiss chard
cabbage	egg plant	mustard greens	radishes	tomatoes
cantaloupe	endive	okra	rutabaga	turnips
carrots	garlic	onions	salsify	watermelons

hazelnuts	pecans	quince
cherries	persimmons	walnuts
chestnuts	pineapple	
<b>Bush and Vine Planting Sites:</b>		
blackberries	currants	huckleberries
blueberries	dewberries	loganberries
boysenberries	gooseberries	raspberries
cranberries	grapes	strawberries
		youngberries

### DOSAGE AND USE RECOMMENDATIONS to Control Nematodes, Symphylans, and Wireworms†

Crops (consult list of individual crops under General Information)	Type of Treatment	Soil Type	Dosage			
			Gallons per Acre <sup>1</sup>	Fl. Oz./1000 Ft. Row Per Chisel		
Shallow Rooted Plants: Field Crops Floral Crops Grasses and Turf Small Fruits Vegetables Ornamentals	Row (42") <sup>6</sup>	Mineral	4.5 to 6	46 to 62		
		Muck or Peat	9 to 12	93 to 123		
Strawberries	Overall (Broadcast)	Mineral	9 to 15 <sup>2</sup>	26 to 44		
		Muck or Peat	24 <sup>3</sup> to 36	71 to 106		
Sugar Beets Root-Knot Nematode	Row (42")	Mineral	6	62		
Overall	12 to 15		35 to 44			
Sugar Beet Nematode	Row (42")	Mineral	9	93		
			Overall	12 to 18	35 to 53	
Pineapple <sup>4</sup>	Row	Mineral	24 to 36	—		
Citrus — Florida <sup>5</sup>	Overall	Mineral	36	106		
Nursery and Field: <sup>7</sup> Citrus Fruit Trees Deciduous Fruit Trees <sup>6</sup> Forest Trees Grapes Nut Trees Ornamentals (Deep rooted)	Overall: Gallons per Acre to Penetrate Various Depths					
	Mineral Soils	3 ft.	4 ft.	5 ft.	6 ft.	
	Sand	15	21	27	33	
	Sandy Loam	27	30	36	48	
	Silt Loam	42	51	63	75	
Clay	54	69	84	102		

<sup>1</sup> Use the higher rates in heavier soil

<sup>2</sup> For cyst forming nematodes increase dosage to 18 gallons per acre (53 fl. oz./1000 ft. row per chisel)

<sup>3</sup> For muck soils containing less than 30 percent organic matter use 18 gallons per acre.

<sup>4</sup> For Hawaiian pineapple, application may be made at time of or just before planting.

<sup>5</sup> For burrowing nematode in citrus inject on 18-inch centers, 12 inches deep. Keep free of plants susceptible to burrowing nematodes for 2 years before replanting to citrus.

<sup>6</sup> Row treatment is not recommended for potatoes in irrigated areas of western and northwestern states.

<sup>7</sup> Tree planting sites prepared by backhoeing may be treated by pouring 27 fl. oz. of TELONE II into the hole during backfilling. For best results prepare and treat sites in the fall and plant in the spring.

<sup>8</sup> Use TELONE II as an aid in the control of bacterial canker and decline of peach trees by application as a preplant, overall treatment of light (sandy) soils at the rate of 24 to 36 gallons per acre (71 to 106 fl. oz. / 1000 ft. row per chisel) preferably in the fall when the soil is warm (50-85°F at 6 inch depth) and moist. Inject the fumigant at a depth of 10 to 12 inches with chisels mounted on 12 inch centers. Follow directions for soil preparation, sealing, exposure, and aeration as specified elsewhere on this label.

†NOTE: To control symphylans (garden centipedes) use only overall at 18 or more gallons per acre, and apply during late summer or early fall when the soil is warm. To control wireworms use the higher dosages recommended for nematodes in overall or broadcast treatments.