

**RESTRICTED USE PESTICIDE**

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certificate.

**PRECAUTIONARY STATEMENT  
Hazard to Humans and Domestic Animals**

**WARNING**

May be fatal if swallowed. Do not breathe dust or fumes. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Remove and wash contaminated clothing before reuse. Do not contaminate food or foodstuffs. Keep away from children and domestic animals. Wear protective clothing or rubber gloves when handling bait. Not for use or storage in or around the home.

**ENVIRONMENTAL HAZARDS**

This product is toxic to wild and tame birds and other wildlife feeding in treated areas may be killed. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. This product shall not be broadcast over bodies of water, roads, roadsides, areas inhabited by livestock or where a hazard exists to rare or endangered species. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of waste.

**DIRECTIONS FOR USE**

**FOR DISTRIBUTION AND USE ONLY IN HAWAII**

**STORAGE AND DISPOSAL**

**PROHIBITIONS**

Do not contaminate water or food by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL**

Dispose of excess bait by ~~burn~~

**CONTAINER DISPOSAL**

Dispose of containers by ~~burn~~

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS**

This product is to be used only for the control of Polynesian rats (Rattus exulans), Norway rats (R. norvegicus), roof rats (R. rattus), house mice (Mus musculus) in sugarcane fields or in macadamia nut orchards. Do not use this product in areas adjacent to these sites or in any other crop or noncrop area. This product may be used only in Hawaii.

**SUGARCANE FIELDS**

**Prebaiting:** To increase acceptance by rodents of treated bait, prebait with 3 pounds of untreated oat groats per acre 1 or 2 weeks prior to scheduled application of toxic bait.

**HGP INC ZINGPHOS OATS**

**ZINC PHOSPHIDE ON OAT GROATS**

For the control of Polynesian Rats (Rattus Exulans), Norway Rats (Rattus Norvegicus), Black Rats (R. Rattus) and House Mice (Musculus)

**FOR DISTRIBUTION AND USE ONLY IN HAWAII**

**ACTIVE INGREDIENTS**

Zinc Phosphide ..... 1.89%

**INERT INGREDIENTS**

..... 98.12%

**TOTAL** ..... 100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

Any persons applying zinc phosphide products and experiencing signs and symptoms such as nausea, abdominal pain, tightness in the chest, or weakness, should be seen by a physician immediately.

If swallowed, call a physician or poison control center. Give 1 or 2 glasses of water and induce vomiting by touching back of throat.

Avoid use of all oils. Have patient lie down and keep warm. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin or in eyes - flush with plenty of water. Get medical attention if irritation persists.

See left side panel for additional precautionary statements.

Manufactured by HGP Inc., P.O. Box 3700, Honolulu, Hawaii 96826  
EPA Establishment No. 1144-HI-01 (and 02)  
EPA SLN No. HI 80010

**NET WEIGHT:**

**Baiting:** Apply up to 5 pounds of toxic bait per acre. Bait may be broadcast by aircraft, ground-driven devices, or by hand in fields of sugarcane that are at least 6 months of age. Do not apply bait within 30 days of harvest. Do not apply bait to adjacent noncrop areas. Wear gloves to avoid possibility of touching bait with bare hands. Do not apply more than 5 pounds per acre per treatment. Do not apply more than 20 pounds of bait per acre per 22-to 36-month crop.

**MACADAMIA NUT ORCHARDS**

**Spot Treatment:** Apply one or two placements of 1 tablespoon (7 to 14 grams) of bait in bait stations located at the base of each tree or in suitable forks in each tree. Remove bait stations from trees prior to shaker harvest and from ground and trees prior to mechanized harvest.

**Burrow Treatment:** Place 1 tablespoon of loose or bagged bait well into each burrow. No preharvest interval is required if nuts falling into burrows are not retrieved.

**Broadcast Treatment:** Apply up to 5 pounds of bait per acre in the orchard by ground-driven devices or by hand. Do not apply bait within 30 days of a round of harvesting. Do not apply bait to adjacent noncrop areas. Wear gloves to avoid possibility of touching bait with bare hands. Do not apply more than 5 pounds per acre treatment. Do not apply more than 20 pounds of bait per acre per year.

**NOTICE:**

Seller makes no warranty, expressed or implied, concerning the use of this product other than that indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

03 113  
11464-2