

19 53

AUG 25 1992

Ms. Pat Kerwin
Foremost Fertilizer, Inc.
P.O. Box 491260
900 Thomas Road
Leesburg, Florida 34749-1260

Subject: St. Augustine Weed and Feed
EPA Reg. No. 35306-4
RE: Transfer from Florida Favorite Fertilizer to Foremost
Fertilizer. Your submission dated January 10, 1992.

Dear Ms. Kerwin:

The above mentioned amendment submitted in connection with registration under FIFRA is accepted. However please adhere to the following changes to update your label.

1. Correct the following typographical error:
 - a. page two "crabgrass"
2. Delete the wording "All crops" on page one. Since this product is for lawn use only "All crops" would be a misleading statement.
3. To update the "Environmental Hazards" section ...
 - a.) replace "Do not apply directly to water" and "Do not apply directly to water or wetlands" with "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."
 - b.) replace "Do not contaminate water by cleaning of equipment or disposal of wastes" with "Do not contaminate water when disposing of equipment washwaters."

A stamped copy is enclosed for your records.

Sincerely,

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch

CONCURRENT REGISTRATION DIVISION (H7505C)							
SYMBOL	H7505C						
SURNAME	HICKS						
DATE	8-25-92						

FOREMOST

COVERS UP TO 5000 SQ. FT.

ST. AUGUSTINE WEED & FEED

16-4-8 + IRON

ACCEPTED
JG 25 1992
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7366-4

KEEP OUT OF THE REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: CAUTION, causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If in eyes, flush with plenty of water. Get medical attention if irritation persists. Harmful if swallowed. Do not contaminate water used for irrigation or domestic purposes. Do not graze treated areas. Do not feed clippings to livestock.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Not for use on or under trees, shrubs, bedding plants, or garden plants. Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to surface and where these soils are very permeable, i.e., well drained. Your local agricultural agencies can provide further information on the type of soil in your area and location of groundwater.

ALL crops- This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACTIVE INGREDIENTS:

*Atrazine(2-chloro-4-ethylamino-6isopropylamino-s-triazine.....	75%
Inert ingredients.....	99.25%
TOTAL.....	100.00%

*Contains 15.01 lbs. of Atrazine per ton

EPA Reg. No. 35306-4
EPA Est. No. 35306-FL-1

Manufactured by,

NET WT. 32 LBS.

FOREMOST FERTILIZER INC.,
900 THOMAS ROAD
LEESBURG, FLORIDA 32748

FRONT

FOREMOST

ST. AUGUSTINE WEED AND FEED

16-4-8 + IRON

GUARANTEED ANALYSIS

1027-

Total Nitrogen.....	16.00%
Nitrate Nitrogen.....	0.00%
Ammoniacal Nitrogen.....	1.70%
Water soluble Organic Nitrogen.....	0.00%
(and / or Urea Nitrogen.....	14.30%
Water Insoluble Nitrogen.....	0.00%
Available Phosphoric Acid.....	4.00%
Soluble Potash (K2O).....	8.00%
Chlorine, not more than.....	8.00%
Primary Plant Nutrient Sources: Sulfate of ammonia, urea	
Triple Superphosphate	
Muriate of Potash	

Secondary Plant Nutrients:

 Iron, as Fe. 2.00%

Secondary Plant Nutrient Sources: Iron Oxide

WHAT THIS PRODUCT DOES

Foremost St. Augustine Weed & Feed controls and prevents weeds while feeding your lawn.

CONTROLS these common lawn weeds:

black medic	chickweed	Florida betony
blue toadflax	dandelion	Florida pusley
buckhorn	dichondra	fragrant cudweed
burclover	dollarweed	(rabbit tobacco)
Carolina geranium (cranesbill)	evening primrose	henbit

lippia (matchweed)	shepherd's purse
oxalis (yellow woodsorrel)	sowthistle
pepperweed	stinging nettle
purslane	Venus looking-glass
red sorrel (sheep sorrel)	wild mustard

rescuegrass

PREVENTS these annual weeds:

craggrass	sedges	spotted spurge	bull paspalum
-----------	--------	----------------	---------------

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product inconsistent with its labeling.

Apply between Jan. 1 and March 31 in Spring and after September in the fall at the rate of 32 pounds on 5,000 sq.ft.

DO NOT OVERAPPLY

In Florida and other Southern states, applications may be made after March 31.

For best results apply before the weeds come up.

Established weeds are harder to control and may require additional applications in the Spring and Fall.

WATER IN THOROUGHLY

SPREADER SETTING:

Cyclone.....5 Scotts.....8 1/2

NOTE: Spreader rates vary with walking speed. Check your spreader over a known area.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds.

PRODUCT DISPOSAL: To dispose of bags which contain unused product securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag and discard in trash.

NOTE: Avoid getting on concrete walks or drives, swimming pools, stucco or asbestos siding as the composition of certain fertilizer elements can cause permanent stains.

Back