

DUSTING SULPHUR

Active Ingredient:

Sulphur

Inert Ingredients

95%

5%

100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause eye irritation. In case of contact, wash eyes with cold water. Do not contaminate water by cleaning of equipment or disposal of waste. Do not re-use container. Container should be disposed of by burying with waste or by burning. Keep out of smoke. Apply this product only as specified on this label.

Sulphur will cause severe injury to sulphur-sensitive crops.

Do Not Use On Sulphur-Sensitive Melons, Apricots, Squash, Cucumbers, Filberts, Spinach, and Walnuts or Other Plants That May Be Injured By Sulphur.

Sulphur may burn certain crops during high temperatures depending on climatic conditions. Do not apply this product when temperatures exceed 90 F. Do not use sulphur within 30 days of an oil spray (60 days on citrus).

Net Contents 50 lbs.

ARIZONA SULPHUR COMPANY

Glendale, Arizona

EPA Reg. No. 35377-1

EPA - Est. No. 35377-AZ-1

IMPORTANT

Consult State Agricultural Experiment Stations for additional information as the Number, Timing, and Rate of application needed may vary with local conditions.

NOTICE

Arizona Sulphur Company makes no warranty, guaranty, or representation, express or implied concerning this material or the use thereof, except the guaranteed analysis of material packaged in this container. All statements made on this label are believed to be true and accurate as possible.

Arizona Sulphur Company shall not be held responsible for any injury or property damage or loss resulting to the buyer from handling, storage, or use of this material, not in accordance with directions. Because of conditions of improper handling, storage, and use which are beyond our control the buyer assumes all risk and liability and uses this material on these conditions.

DIRECTIONS FOR USE

Dusting sulphur should be applied with the specified amount using aircraft or ground equipment depending upon local conditions. Avoid applying dusting sulphur when windy conditions prevail as drifting of product will occur. Except as otherwise shown, apply this product at the first sign of infection and repeat at 7-10 day intervals as needed to maintain control of problem area.

RECOMMENDATIONS FOR USE

Bush and Vine Fruits:

Boysenberries, Raspberries, Strawberries — For the control of powdery mildew use 10-20 pounds per acre. Apply when disease first appears and repeat as necessary.

Grapes — For the control of powdery mildew and rust apply 5-15 pounds per acre in four applications. Make first application when shoots are 6 inches to 8 inches long. Make second application when shoots are 12 inches to 15 inches long and a third application about 14 days later. Repeat applications at 10-14 day intervals thereafter to harvest.

Citrus:

Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines — For the control of Citrus Thrips, Citrus Citrus, Two-spotted Mite, Citrus Rust Mite, Yuma Spider Mite, apply 100-125 pounds per acre. Apply when pests first appear and repeat as necessary. Avoid applications during or preceding high temperatures. DO NOT APPLY WITHIN 2 MONTHS OF AN OIL SPRAY.

Deciduous Fruits and Nuts:

Almonds, Cherries, Nectarines, Peaches, Plums, Prunes — For the control of blossom blight brown rot, powdery mildew, cherry leaf spot, prune rust, use at the rate of 45-50 pounds per acre in pre-bloom, bloom, and post-bloom periods.

Apples — For the control of scab and powdery mildew use 30-40 pounds per acre. Apply in pre-bloom, bloom, and post-bloom periods. USE ON SULPHUR-TOLERANT VARIETIES ONLY.

Field Crops:

Alfalfa — For the control of Two-spotted Mites, Pacific Mites, Atlantic Mites, apply 30-40 pounds per acre. CAUTION: Sulphur may cause petal burn to open flowers.

Clover — For the control of powdery mildew and Spider Mites, apply 15-25 pounds per acre.

Cotton — For the control of Atlantic Mite and Cotton rust, apply 30-40 pounds per acre when mites first appear and repeat as necessary.

Sugar Beets — For the control of Spider Mite, apply 30-40 pounds per acre.

Vegetables:

Asparagus — For the control of Two-spotted Mite, Pacific Mite, Atlantic Mite, apply 30-40 pounds per acre. CAUTION: Sulphur may cause petal burn to open flowers.

Beans — For the control of powdery mildew, rust, Spider Mite, apply 15-30 pounds per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. CAUTION: May cause injury to some varieties of beans.

Beets, Broccoli, Brusselsprouts, Cabbage, Cauliflower, Lettuce — For the control of powdery mildew and Two-spotted Mite, apply 25-35 pounds per acre.

Carrots — For the control of Petrobia Mite, apply 30-35 pounds per acre.

Onions — For the control of Petrobia Mite, apply 25-30 pounds per acre.

Rutabagas — For the control of powdery mildew, apply 15-25 pounds per acre.

Tomatoes — For the control of Tomato Russet Mite, apply 25-40 pounds per acre.

Turnips — For the control of powdery mildew, apply 15-25 pounds per acre.

DUSTING SULPHUR

Active Ingredient:

Sulphur	95%
Inert Ingredients	5%
	<hr/>
	100%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause eye irritation. In case of contact, wash eyes with cold water. Do not contaminate water by cleaning of equipment or disposal of waste. Do not re-use container. Container should be disposed of by burying with waste or by burning. Keep out of reach of children. Apply this product only as specified on label.

Do Not Use On Sulphur-Sensitive Melons, Apricots, Squash, Peaches, Apples, Spinach, and Walnuts or Other Plants Not Labeled for Insecticide Use by Sulphur.

See Container for Full Directions

WELLS & SULPHUR COMPANY

Glendale, California

U.S. Pat. No. 3,137,771

U.S. Pat. No. 3,137,772

WELLS & SULPHUR COMPANY

DIRECTIONS FOR USE

Dusting Sulphur is used to control insects and mites on plants. It is applied as a dust to the foliage and soil around the plants. It is also used to control insects and mites on stored products. It is applied as a dust to the stored products. It is also used to control insects and mites on stored products. It is applied as a dust to the stored products.

vegetation

RECOMMENDATIONS FOR USE

Bush and Vine Fruits

Apply as a dust to the foliage and soil around the plants. It is also used to control insects and mites on stored products. It is applied as a dust to the stored products. It is also used to control insects and mites on stored products. It is applied as a dust to the stored products.

CITRUS

Apply as a dust to the foliage and soil around the plants. It is also used to control insects and mites on stored products. It is applied as a dust to the stored products. It is also used to control insects and mites on stored products. It is applied as a dust to the stored products.