



# WETTABLE SULPHUR

Active Ingredient

Sulphur	97%
Inert Ingredients	3%
	100%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

May cause eye irritation. In case of contact, wash eyes with cold water. Do not contaminate water by cleaning of equipment or disposal of waste. Do not re-use container. Container should be disposed of by burying with waste or by burning. Keep out of smoke. Apply this product only as specified on this label.

Sulphur will cause severe injury to sulphur-sensitive crops. Do Not Use On Sulphur-Sensitive Melons, Apricots, Squash, Cucumbers, Filberts, Spinach, and Walnuts or Other Plants That May Be Injured By Sulphur.

Sulphur may burn certain crops during high temperatures depending on climatic conditions. Do not apply this product when temperatures exceed 90°F. Do not use sulphur within 30 days of an oil spray. (60 days on citrus)

Net Contents 50 lbs.

Manufactured for:

**ARIZONA SULPHUR COMPANY**

Glendale, Arizona

EPA Reg. No. 35377-2

EPA - Est. No. 476-CA-5

## IMPORTANT

Consult State Agricultural Experiment Stations for additional information as the **Number, Timing, and Rate** of application needed may vary with local conditions.

## NOTICE

Arizona Sulphur Company makes no warranty, guaranty, or representation, express or implied concerning this material or the use thereof, except the guaranteed analysis of material packaged in this container. All statements made on this label are believed to be true and accurate as possible.

Arizona Sulphur Company shall not be held responsible for any injury or property damage or loss resulting to the buyer from handling, storage, or use of this material, not in accordance with directions. Because of conditions of improper handling, storage, and use which are beyond our control the buyer assumes all risk and liability and uses this material on these conditions.

Wettable sulphur should be applied with the specified amount using aircraft or ground equipment depending upon local conditions. Using the specified amount of this product shown, wash through filler screen in nearly filled spray tank. Add balance of water to fill tank. This product should be continually kept under agitation during filling and spraying operations. Do not allow mixture to stand.

## RECOMMENDATIONS FOR USE

Bluyenberries - Raspberries - For the control of powdery mildew use 2.5 pounds per 100 gallons. Begin application when shoots are 3 inches long and before blossom open. Repeat at 10 day intervals. Discontinue applications during hot weather.

Strawberries - For the control of powdery mildew use 1.5 pounds per 100 gallons. Begin at first sign of disease and repeat as necessary. NOTE: Sulphur may injure strawberries under certain climatic conditions.

Grapes - For the control of powdery mildew use 2 pounds per 100 gallons when the shoots are 12-15 inches long. This spray should be used particularly if earlier sulphur dustings have been omitted or for any reason mildew has not been held in check with a sulphur dust. Care should be taken to wet all surfaces on which the fungus is to be controlled.

### Citrus:

Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangelos, Tangerines - For the control of citrus flatmite, Six spotted mite, Citrus Rust mite, Yuma Spider mite use 4 pounds per 100 gallons.

### Deciduous Fruits and Nuts:

Almonds, Cherries - For the control of blossom blight brown rot, cherry leaf spot, and powdery mildew use 4-6 pounds per 100 gallons. Begin treatment when disease threatens or in pre-blossom stage. Repeat at petal fall and as necessary thru cover period.

Nectarines, Peaches - For the control of brown rot, Rust, Peach Silver mite and powdery mildew use 5-6 pounds per 100 gallons from popcorn stage up to harvest. For the control of brown rot in summer sprays use up to 10 pounds per 100 gallons starting about 4-5 weeks before harvest and continuing at weekly intervals.

Plums, Prunes - For the control of brown rot, leaf spot, and rust use 5-6 pounds per 100 gallons. Begin treatments when disease threatens or in pre-blossom stage. Repeat at petal fall and as necessary thru cover period.

Apples - For the control of scab and powdery mildew use 4-5 pounds per 100 gallons. For early season scab control use at least 3 sprays beginning at green tip and continuing through pink or petal fall stages. Use on sulphur tolerant varieties only.