

35484-1

09/26/2011

1/13



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Chemical Safety and Pollution Prevention
Office of Pesticide Programs
Registration Division (7504P)
1200 Pennsylvania Ave., N.W.
Washington, DC 20460

EPA Reg. Number:

35484-1

Date of Issuance:

SEP 26 2011

Term of Issuance:
Unconditional

Name of Pesticide Product:

**Gordon's Bordeaux
Mixture**

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Industrias Quimicas del Valles,
S.A. (IQV, SA)

Mailed to:

c/o Crystal Layton
Landis International, Inc.
PO Box 5126
Valdosta, GA 31603-5126

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted by email on July 25, 2011. EPA grants this request under the authority of section 3(c)(5) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. With this accepted labeling, all requirements set forth in the Reregistration Eligibility Decision for coppers have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Signature of Approving Official:

Date:

SEP 26 2011

Tony Kish, Product Manager (22)
Fungicide Branch/Registration Division/OPP/OCSP (7504P)

Notice of Pesticide Reregistration
Gordon's Bordeaux Mixture
EPA Reg. No. 35484-1
Page 2 of 2

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after 12 months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

If you have any questions or comments regarding this letter, please contact Dominic Schuler at (703) 347-0260 or via e-mail at schuler.dominic@epa.gov.

Enclosure:

- Label stamped "Accepted"
- PRB Label Review dated 7/14/2003
- Acute Toxicity Review DP364841 dated 5/11/2009
- Product Chemistry Review DP364840 dated 5/14/2010

3/13

Not for residential use or application to residential sites. This product may not be used in, on, or around any structure, vehicle, article, surface or area associated with the household, including but not limited to areas such as non-agricultural outbuildings, non-commercial greenhouses, pleasure boats and recreational vehicles; in or around any preschool or day care facility or on humans or pets.

GORDON'S BORDEAUX MIXTURE

EPA Reg. No. 35484-1

EPA EST. No.35484-ESP-001


Active Ingredient:

Copper Sulfate Pentahydrate (CAS #7758-99-8) 52.25%

Inert Ingredients: 47.75%

Total: 100.00%

Metallic Copper (Cu²⁺) Equivalent: 13.30%



KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label find someone to explain it to you in detail.)

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Note to Physician: Probable mucosal damage may contraindicate use of gastric lavage.	
Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: www.npic.orst.edu). After 5 p.m. call your poison control center at 1-800-222-1222.	

Industrias Quimicas de Valles, SA	
Rafael Casanova 81	P.O. Box 5126
08100 Mollet del Valles, Spain	Valdosta, GA 31603-5126
229-247-6472	

Net Contents: 50; 2,000 pounds

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER

Corrosive. Causes irreversible eye damage. May be fatal if swallowed. Harmful if absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear protective eyewear such as goggles, face shield, or safety glasses. Wear rubber gloves when handling. Harmful if swallowed.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear the following:

- Long-sleeve shirt,
- Long pants,
- Goggles or faceshield,
- Chemical resistant gloves made of any waterproof material,
- Shoes plus socks.

User Safety Requirements

Follow manufacturer's instructions for cleaning /maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations
 User should wash hands before eating, drinking, chewing gum, tobacco, or using the toilet.
 User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 User should remove PPE immediately after handling this product. As soon as possible wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates, and may contaminate water through runoff. This product has the potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for residential use or application to residential sites. This product may not be used in, on, or around any structure, vehicle, article, surface or area associated with the household, including but not limited to areas such as non-agricultural outbuildings, non-commercial greenhouses, pleasure boats and recreational vehicles; in or around any preschool or day care facility or on humans or pets.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as soil or water is as follows:

- Coveralls,
- Shoes plus socks,
- Chemical-resistant gloves made of any waterproof material,
- Goggles or faceshield.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

Disease control is a combination of prevention and protective sprays. Begin applications at the first signs of disease infection or when disease threatens.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition (approximately 3 to 10 mph), and there are no sensitive areas within 250 feet downwind.

Temperature Inversions If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements Applicators must follow all state and local pesticide drift requirements regarding application of copper compounds. Where states have more stringent regulations, they must be observed.

Equipment All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Do not apply by air.

SPRAY PREPARATION:

The spray suspension is prepared by dissolving the Gordon's Bordeaux Mixture in one-half the required amount of water. After the product is dispersed completely with agitation, add the remainder of the water to the tank. Maintain agitation in the spray tank to prevent settling. Gordon's Bordeaux Mixture should be applied the same day that it is prepared.

Metallic Copper (Cu2+)	Gordon's Bordeaux Mixture
0.5 lb/A	3.75 lb/A
1.0 lb/A	7.5 lb/A
1.5 lb/A	11.3 lb/A
2.0 lb/A	15.0 lb/A
2.5 lb/A	18.8 lb/A
3.0 lb/A	22.5 lb/A
4.0 lb/A	30.0 lb/A
8.0 lb/A	60.0 lb/A

Metallic Copper (Cu2+)	0.5 lb/A	1.0 lb/A	1.5 lb/A	2.0 lb/A	2.5 lb/A	3.0 lb/A	4.0 lb/A	8.0 lb/A
Gordon's Bordeaux Mixture	3.75 lb/A	7.5 lb/A	11.3 lb/A	15.0 lb/A	18.8 lb/A	22.5 lb/A	30.0 lb/A	60.0 lb/A
Amount of water, gallons	Amount of Gordon's Bordeaux Mixture							
5.0 gallons	0.20 lb	0.38 lb	0.56 lb	0.75 lb	0.94 lb	1.13 lb	1.5 lb	3.0 lb
50 gallons	1.88 lb	3.8 lb	5.6 lb	7.5 lb	9.4 lb	11.3 lb	15.0 lb	30.0 lb
100 gallons	3.75 lb	7.5 lb	11.3 lb	15.0 lb	18.8 lb	22.5 lb	30.0 lb	60.0 lb

Table 3. Use rates of Gordon's Bordeaux Mixture for the listed fruits, vegetables and pecans				
Plant	Disease	Amount of product mixed in water per acre application	Maximum amount of product per acre per year	Minimum retreatment interval
Apples	Late scab, black rot, bitter rot, & blotch	3.75 lb/100 gal/A/app	120 lb/A/yr	5 days
	A protective spray applied during bloom will help prevent infection. Treatments should not be applied after the ¼ inch green stage. May russet certain varieties. Do not spray tender foliage or during the early growing period when the leaf tissue is emerging from the bud.			
Pears	Fire blight	3.75/100 gal/A/app	120 lb/A/yr	5 days
	Bloom sprays are applied to control the blossom blight stage of this disease. Do not add insecticides or exceed this dosage rate. Repeat at 5 to 7 day intervals until the end of the blooming period.			
Apricots, Plums, Sour Cherries	Bacterial leaf spot & brown rot	7.5 to 11.3 lb/100 gal/A/app	135 lb/A/yr	5 days
	Apply at the shuck split stage. Also, the same dosage can be used for the first and second cover sprays. Do not use on Japanese pls.			
Peaches	Peach leaf curl (dormant)	18.8 lb/100 gal/A/app	135 lb/A/yr	7 days
	Use dormant applications after the leaves have dropped in the fall or before the buds swell in the spring. Thorough cover is essential.			
Pecans	Pecan scab	15.0 lb/100 gal/A/app	63.2 lb/A/yr	14 days
	Apply when catkins show. Use three (3) or four (4) applications at 2-3 week intervals			
Grapes Grapes (dormant)	Black rot & powdery mildew	3.75 lb/ 100 gal/A/app	150 lb/A/yr	3 days
	Apply at the early pre-bloom stage, post-bloom stage, and mid-summer stage. For the early bloom stage apply when the shoots are 18 inches long. For the post-bloom stage apply when 90% of blossoms have opened. For the mid-summer stage, apply during July and August depending upon the severity of the disease. This use will exhibit some phytotoxicity on most varieties. Dormant: Apply in spring before bud-swell and green tissue is present.			
Potatoes	Early & late blight	18.8 lb/100 gal/A/app	188 lb/A/yr	5 days
	Apply when the plants are 3 to 4 inches in height. Repeat treatments at the same dosage rate at 5 to 14 day intervals during the entire growing season.			
Strawberry	Leaf spot	7.5 lb/100 gal/A/app	61.5 lb/A/yr	7 days
	Severe Disease	11.3 lb/100 gal/A/app		
Apply at 7 to 14 day intervals from the pre-bloom stage until harvest. Begin treatments at the first sign of leaf spotting. Repeat treatments as long as disease continues.				

Table 3. Use rates of Gordon's Bordeaux Mixture for the listed fruits, vegetables and pecans.

Plant	Disease	Amount of product mixed in water per acre application	Maximum amount of product per acre per year	Minimum retreatment interval
Citrus	Brown rot of lemons, grapefruit scab, melanose of grapefruit & oranges	15.0 lb/gal/A/app	95 lb/A/yr	7 days
	Septoria fruit, leaf spot	3.75 lb/gal/A/app		
Apply two (2) treatments at the same dosage rate. Apply the first treatment in the late winter prior to spring growth. Apply the second treatment after the petals have fallen from the blossoms but before the new fruit is larger than a pea. Apply in October, November or December.				
Tomato	Early blight, late blight, & Septoria leaf spot	3.75 lb/100 gal/A/app	60 lb/A/yr	3 days
	Begin applications when disease threatens and repeat applications at 3 to 10 day intervals using full cover spray.			

For Ornamental Plants (Trees, Shrubs, and Flowers)

Application rate: The maximum amount of metallic copper (Cu²⁺) that may be applied to the listed ornamental plants is 2.0 pounds of metallic copper (Cu²⁺) per acre per application or 15 pounds of product per acre per application.

Minimum re-treatment interval: The minimum number of days between each application is 7 days.

Maximum seasonal rate: The maximum seasonal amount of metallic copper (Cu²⁺) that may be applied to these listed ornamental plants is 20.0 pounds of metallic copper (Cu²⁺) per acre per year or 150 pounds of product per acre per year.

Table 4. Use rates of Gordon's Bordeaux Mixture for the listed ornamental plants (trees shrubs and flowers).

Plant	Disease	Amount of product mixed in water	Maximum amount of product per acre per year	Remarks	Minimum retreatment interval
Ivy	Leaf spot	15.0 lb/100 gal/A/app	150 lbs	Apply full cover spray. Begin applications when the infection begins and repeat application in 14 days if necessary	7 days
Delphinium & Geranium	Leaf spot, blight	11.3 lb/100 gal/A/app	150 lbs	Begin applications when disease infection begins and repeat treatments if necessary	7 days
Peony	Leaf blotch & botrytis blight (gray mold)	15.0 lb/100 gal/A/app	150 lbs	Spray before shoots are 1 ft. tall. Repeat the application at the same dosage rate in 14 days. If any sign of bud blast, spray again. Remove any wilted shoots by cutting below ground, place in paper bag & burn. Remove old stalks and leaves in fall and burn – Do Not Compost.	7 days
Camellia	Dieback	15.0/100 gal/A/app	150 lbs	Apply in early spring.	7 days
Dahlia, Tulip, Gladiolus & lily	Botrytis blight (gray mold)	15.0/100 gal/A/app	150 lbs	Apply when growth begins in the spring. Repeat as necessary to maintain control at 7 to 10 day intervals.	7 days
Iris & Chrysanthemum	Leaf spot & blight	15.0 lb/100 gal/A/app	150 lbs	Apply when growth begins in the spring. Repeat as necessary to maintain control at 7 to 10 day intervals.	7 days
Hollyhock & Phlox	Anthrachnose & leaf spot	15.0 lb/100 gal/A/app	150 lbs	Spray after periods of wet weather.	7 days
Pansy	Downy mildew	15.0 lb/100 gal/A/app	150 lbs	Spray every 14 days during growing season.	7 days
Rose	Powdery mildew & black spot	15.0 lb/100 gal/A/app	150 lbs	Begin application when disease occurs and conditions favor the development of the disease. Apply as a full cover spray in the early spring as leaves expand. Treat at 10 to 14 day intervals. Prune out diseased and dead canes.	7 days
Arborvitae, Cedar, Cypress & Juniper	Coryneum blight, leaf blotch, twig blight, leaf spot & cedar apple rust	15.0 lb/100 gal/A/app	150 lbs	Apply in early spring and fall. Repeat at 14 day intervals as needed.	7 days

Pine	Tip blight & needle cast	15.0 lb/100 gal/A/app	150 lbs	Three applications as a full cover spray are suggested. Spray as new growth starts, as new needles emerge from the sheath, and when needles are 2/3 of mature length.	7 days
Laurel, Rhododendron & Azalea	Leaf spot, leaf blight & <i>Pseudomonas syringae</i>	15.0 lb/100 gal/A/app	150 lbs	Apply as a full cover spray in early spring. A second application at the post-bloom stage is recommended.	7 days
Palm	Anthraxnose, scab & leaf spot	15.0 lb/100 gal/A/app	150 lbs	Apply when disease symptoms appear. Repeat treatments to maintain control.	7 days
Barberry	Bacterial leaf spot	15.0 lb/100 gal/A/app	150 lbs	Apply when new leaves appear. Repeat two (2) to three (3) times at 10 day intervals.	7 days
Boxwood	Nectria Canker & leaf spot	15.0 lb/100 gal/A/app	150 lbs	Begin applications in early spring growth stages. Repeat treatments as needed to maintain control.	7 days
Yew	Twig blight	15.0 lb/100 gal/A/app	150 lbs	Apply in spring just prior to bud break and make two (2) applications at 10 day intervals if needed.	7 days
Blue spruce & Douglas fir	Needle cast	15.0 lb/100 gal/A/app	150 lbs	Apply when the new needles are half developed (1/2 to 1 inch long). Repeat the application when the new needles are full length.	7 days
Sycamore, Linden, Oak, Tulip Tree, Elm, Ash, & Walnut	Anthraxnose, leaf spot & elm leaf curl	15.0 lb/100 gal/A/app	150 lbs	Begin applications when the leaves begin to unfold in the spring to protect the plant before infection begins. Repeat in 14 to 21 days.	7 days
Maple: Red, & Sugar	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Apply as full cover spray as leaves uncurl (bud break). Repeat treatment to maintain control at 14 to 21 day intervals.	7 days
	Nectria canker	15.0 lb/100 gal/A/app	150 lbs	One (1) application in the fall (Oct.) and two (2) to three (3) applications in the spring after the young growth appears may help reduce the infection level.	7 days

Pear, non-Bearing (<i>Pyrus communis</i>)	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Apply as full cover spray as leaves uncurl (bud break). Two (2) to three (3) applications at 14 to 21 day intervals may be needed to maintain control. Dormant applications (before any growth starts in the spring) may aid in the control of <i>Pseudomonas</i> blight and anthracnose.	7 days
Plum, non-Bearing (<i>Prunus</i> sp.)	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Apply as full cover spray as leaves uncurl (bud break). Repeat treatment to maintain control at 14 to 21 day intervals.	7 days
Lilac (<i>Syringa</i>)	Powdery mildew & bacterial blight	15.0 lb/100 gal/A/app	150 lbs	Begin application when disease occurs and conditions favor the development of the disease. Apply as a full cover spray in the early spring. A second application at the post-bloom stage is recommended.	7 days
Japanese dogwood (<i>Cornus kousa</i>) & Flowering dogwood (<i>Cornus florida</i>)	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Begin application at bud break and continue applications at 10 to 14 day intervals until dry weather. Applications during the remainder of the season may be needed if continued periods of wet weather.	7 days
Aspen (<i>Populus</i>)_ & Poplar (<i>Populus</i>)	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Begin application bud break and continue applications at 14 to 21 day intervals.	7 days
Crabapple, non-bearing (<i>Malus</i> sp.)	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Apply as full cover spray as leaves uncurl (bud break). Two (2) to three (3) applications at 14 to 21 day intervals may be needed to maintain control. Dormant applications (before any growth starts in the spring) may aid in the control of <i>Pseudomonas</i> blight and anthracnose.	7 days
Cherry, non-bearing (<i>Prunus</i> sp.)	<i>Pseudomonas</i> blight & anthracnose	15.0 lb/100 gal/A/app	150 lbs	Dormant applications (before any growth starts in the spring) may aid in the control of bacterial canker (<i>Pseudomonas syringae</i>).	7 days

Willow (<i>Salix</i> sp.)	Twig blight (scab)	15.0 lb/100 gal/A/app	150 lbs	Begin applications in the spring when new leaves are visible. Make two (2) to three (3) applications and repeat at 10 to 14 day intervals especially if wet weather occurs.	7 days
----------------------------	--------------------	-----------------------	---------	---	--------

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage Store product in a secure dry place. Keep product dry as product is water soluble. Spilled product should be swept up, used if clean, or disposed of according to the procedures below. Store product in original container. Store pesticide separately to prevent cross-contamination of other pesticides, fertilizers, food and feed.

Pesticide Disposal Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Container Handling Nonrefillable container. Do not reuse or refill this container. (Paper or Plastic Bag) Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. When completely empty, offer for recycling if available, or dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Plastic Container greater than 50 pounds) Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT

CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

APPENDIX

1. **Statements which may appear on different label components depending on packaging configuration.**
 - See next panel for additional Precautionary Statements and First Aid
 - Net Weight: _____
 - EPA Est. No. _____
 - Filled by weight, not by volume, some settling of contents will occur, resulting in container not appearing full. Shake with lid on tightly before using.

2. **Advertising claims that may be presented on container labeling, advertisements, brochures, and other marketing/sales promotional materials:**
 - Mixes with water – Powder.
 - For Disease Control on Evergreens, Shrubs, Shade Trees, Peonies, Flowers, and Fruit Trees.
 - Broad-Spectrum Fungicide