

TURF PRIDE 16-4-8

WITH SURFLAN® PRE-EMERGENCE HERBICIDE

Contains SURFLAN® a Pre-emergent Herbicide for the Control of Crabgrass, Henbit, & Most Other Annual Broadleaf & Grassy Weeds in Lawns, Golf Courses, Parks, & Other Ornamental and Recreational Turf areas.

For Established Southern Lawns
Bermudagrass - Bermudagrass - Centipedegrass
St. Augustinegrass - Tall Fescue - Zoysiagrass

CONTAINS 16-4-8 PLANT FOOD

MAY BE APPLIED IN THE FALL OR SPRING FOR PRE-EMERGENT CONTROL OF ANNUAL GRASSES AND BROADLEAF WEEDS OR WITH A SPLIT APPLICATION FOR LONGER LASTING WEED CONTROL.

ACTIVE INGREDIENT

CHLORANIL, 35 dinitro N₂H₄
dipropyl sulfam, 4.0%
INERT INGREDIENTS

0.2%
2.3%
97.5%

NET WEIGHT 50 POUNDS

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENTS OF PRACTICAL TREATMENT

1. Apply this product to the soil surface of a healthy lawn or golf course in accordance with the directions on the label.
2. Do not apply this product to areas where it is not intended to be used.
3. Do not apply this product to areas where it is not intended to be used.
4. Do not apply this product to areas where it is not intended to be used.
5. Do not apply this product to areas where it is not intended to be used.

NET WEIGHT 50 POUNDS
NET WEIGHT 50 POUNDS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with eyes. Avoid contact with skin. Avoid contact with clothing. Do not eat, drink, or use tobacco products while handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and shoes before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas immediately adjacent to water.

DIRECTIONS FOR USE

For use on lawns and golf courses. For use on lawns and golf courses.

STORAGE AND DISPOSAL

Store in original container. Do not use if container is damaged. Do not use if contents are not as labeled. Do not use if contents are not as labeled. Do not use if contents are not as labeled.

WHERE TO USE

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

GRASSY WEEDS

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

BROADLEAF WEEDS

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

WHEN TO APPLY

FOR BEST RESULTS APPLY AT THE PROPER TIME

Apply only to healthy established turf at least 100 weeks prior to germination of weed seeds since established weeds will be controlled. Annual and annual grasses do not germinate at the same time. Many weeds germinate in the fall. For this reason your application must coincide with their germination period. Apply the early on the later rate weed control will result. See chart for proper timing. WATER THE LAWN AFTER APPLICATION. Best results will be obtained if treated areas are irrigated with 1 to 1 1/2 inch water soon after application. Any cultivation practices that disturb the soil such as verticillaging, should be done prior to treatment.

WINTER ANNUAL WEEDS

For pre-emergence weed control of listed winter annual grasses and broadleaf weeds apply 50 lbs. (7.300 Sq Ft.) of lawn. This is equivalent to 1 lb. herbicide per acre. Apply in the fall as early as Mid-September or early winter, at least one to two weeks prior to warm up weather when a seed for the germination of weed seeds. See chart for timing.

SPRING ANNUAL WEEDS

For control of weeds as listed in late winter or early spring apply the same rate as for winter annual weeds. Apply in the fall as early as Mid-September or early winter, at least one to two weeks prior to warm up weather when a seed for the germination of weed seeds. See chart for timing.

SPLIT APPLICATIONS

For control of weeds as listed in late winter or early spring apply the same rate as for winter annual weeds. Apply in the fall as early as Mid-September or early winter, at least one to two weeks prior to warm up weather when a seed for the germination of weed seeds. See chart for timing.

Although labeled as a pre-emergent herbicide, this product may be applied 4 to 6 weeks after germination of grass weeds. For control of summer annual grasses, apply to the soil surface of lawns and golf courses. For control of winter annual grasses, apply to the soil surface of lawns and golf courses. For control of broadleaf weeds, apply to the soil surface of lawns and golf courses. For control of annual grasses, apply to the soil surface of lawns and golf courses.

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

GUARANTEED FERTILIZER ANALYSIS

16-4-8
16-4-8
16-4-8

Apply to established lawns and golf courses. Apply to established lawns and golf courses. Apply to established lawns and golf courses.

HOWARD FERTILIZER COMPANY
P.O. BOX 13800 • ORLANDO, FL 32859

ACCEPTED

DEC - 2 1988

Under the Federal Insecticide
Fungicide and Rodenticide Act
this product is the pesticide
registered under
EPA Reg. No. 225-111-111

ACCEPTED

DEC - 2 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
Registration No. 100-100-10000
EPA Reg. No. 100-100-10000