

TURF PRIDE 16-4-8 WEEDER

PREMIUM WEED CONTROL WITH COMPLETE LAWN FOOD

50 POUND BAG COVERS 8,000 Sq Ft. of Lawn

Controls Florida Pusley, Lippia, Pigweed and certain Broadleaf Weeds as listed. Prevents Crabgrass, Peppercorn, and Spotted Spurge and other weeds as listed.

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

ACTIVE INGREDIENT

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	0.70%
Related Compounds	0.04%
INERT INGREDIENTS	99.26%
Total 100.00%	

Contains 16 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 35512-RU(14) EPA EST. NO. 35512-FL-1
NET WEIGHT 50 POUNDS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from food or feed products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (leach or leech) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

WEEDS CONTROLLED OR SUPPRESSED

ANNUAL BLUEGRASS (POA ANNUA), ANNUAL BOWTIEWEEED, ANNUAL CLOVER, BLACK MEDIC, BLUE TOADFLAX, BULL PASPALLUM, BUCKHORN PLANTAIN, BUR CLOVER, CAROLINA GERANIUM (CRANEBILL), CHICKWEED (COMMON & MOUSEEAR), COCKLEBUR, COMMON DANDELION, COMMON PURSLANE, CRABGRASS, CROWFOOTGRASS, CUDWEED, CUTLEAF EVENING PRIMROSE, DICHONDRA, DOLLAR WEED, FLORIDA BEGGARWEED, FLORIDA SETONIA (RATTLEBAKE WEED), FLORIDA PUSLEY, FRAGRANT CUDWEED (RABBIT TOBACCO), HENBIT, KNOTWEED, LAMBQUARTER, LESPEDEZA, MAT LIPPIA (MATCHWEED, CREEPING CHARLIE), MONEYWORT, MORNINGGLORY, NARROWLEAF VETCH, PARSLEY-PIERT, OXALIS (COMMON, YELLOW

WEEDS CONTROLLED (Continued)

WOODSORREL, PIGWEED, PURSLANE, RAGWEED, RED SOHREL, RESCUEGRASS, SANDSPUR, SEDGES*, SHEPARD'S PURSE, SICKLE-POD, SPOTTED SPURGE, SPURWEED, STINGING NETTLE, SWINE-CREST, VENUS LOOKING GLASS, VIRGINIA CREEPER, WILD MUSTARD, WOOD SOREL.

*Repeat application are required for control of Nutedge grasses.

WHEN TO APPLY

Best weed control will result if applied before weeds appear since the product acts as a Pre-Emergent AND also as a Post Emergent weed killer.

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a Drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 50 lb bag to treat 8,000 Sq Ft. - 25 lbs. bag treats 4,000 Sq. Ft. of lawn. (equivalent to 5.7 lbs. per 1,000 Sq Ft). Do not exceed recommended rates. Water immediately after application to wash granules off the grass blades.

HOW TO SET YOUR SPREADER CYCLONE 4 SCOTT 6

NOTE: Spreader rate varies with cutting speed. Check spreader over test run.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. Do not use on Bluegrass, Fescue, Bentgrass, Bahia, Fye, Bermuda, or Dichondra lawns. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable garden) in treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.

GUARANTEED FERTILIZER ANALYSIS


Total Nitrogen (N)	16%
Nitrate Nitrogen	0.00%
Ammoniacal Nitrogen	11.20%
Water Soluble Nitrogen	4.80%
(and/or Urea Nitrogen)	
Water Insoluble Nitrogen	0.00%
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%
Chlorine not more than	5.60%

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muricite of Potash, Sulfate-Potash-Magnesium.
Potential Acidity Equivalent to 1245 lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - 1.20%; Iron as Fe - 1.50%; Water Soluble Magnesium Mg - 1.20%; Zinc as Zn - 0.00%; Manganese as Mn - 0.50%; Boron as B 0.02%; Copper as Cu - 0.02%; Molybdenum as Mo - 0.002%.
Derived from: Manganese Sulfate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

NOTICE: Buyer and user assumes all risk of use, storage and/or handling of this material not in accordance with the terms of this label.



Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 35512-14

HOWARD FERTILIZER COMPANY

P.O. BOX 13800 - ORLANDO, FL 32859

