



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
Notification
222455

Application for Pesticide - Section I

1. Company/Product Number 35512-14	2. EPA Product Manager Robert J. Taylor	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted OPP ID # 222455
4. Company/Product (Name) TURF PRIDE WEEDER (0.8% Atrazine)	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) HOWARD FERTILIZER COMPANY P.O. BOX 593800 ORLANDO, FL 32859-3800 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. > N/A Product Name > N/A	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below. Add WPS Exclusionary Stmt.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

>
> Notification Amendment to include WPS exclusionary Statement as follows:
> "Not for use on turf being grown for sale or other commercial use as sod,
> or for commercial seed production or research purposes."
> Found in "Directions For Use" paragraph.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		> <input type="checkbox"/> Metal ----> metal	> <input checked="" type="checkbox"/> Plastic ----> plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container	> <input type="checkbox"/> Glass ----> glass	> <input checked="" type="checkbox"/> Paper ----> paper
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 25;40;50 lb. bag		5. Location of Label Directions <input checked="" type="checkbox"/> On Label -> on label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product Litho or Adhesive				<input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			6. Date Application Received (Stamped)	
Name H. R. MCLANE, INC. By H. R. McLane:	Title Authorized Agent Howard Fertilizer Co	Telephone No. (Include Area Code) 305 661 1706 800 899 3141		
2. Signature 			3. Title Authorized Agent	
4. Typed Name Kathy Larsen, Reg. Affairs James H. Brown, President			5. Date JUL 13 1995	



TURF PRIDE® WEEDER

PREMIUM WEED CONTROL WITH COMPLETE LAWN FOOD

50 POUND BAG COVERS

8,800 Sq.Ft. of Lawn

Controls Florida Pusley, Lippia, Pigweed and certain Broadleaf Weeds as listed. Prevents Crabgrass, Paspalums, and Spotted Spurge and other weeds as listed.

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	- - -	0.76%
Related Compounds	- - -	0.04%
INERT INGREDIENTS	- - -	99.20%
		Total 100.00%

Contains 16 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

WEEDS CONTROLLED OR SUPPRESSED

- | | |
|----------------------------------|---------------------------------------|
| Annual Bluegrass (Poa Annua) | Henbit |
| Annual Sowthistle | Knotweed |
| Clover | Lambsquarter |
| Black Medic | Lespedeza |
| Blue Toadflax | Lippia (Matchweed) |
| Bull Paspalum | Moneywort |
| Buckhorn Plantain | Morning Glory |
| Bur Clover | Narrowleaf Vetch |
| Carolina Geranium (Cranesbill) | Parsley-Piert |
| Chickweed (Common and Mouse Ear) | Oxalis (Common and Yellow Woodsorrel) |
| Cocklebur | Pigweed |
| Common Dandelion | Purslane |
| Common Purslane | Ragweed |
| Crabgrass | Red Sorrel |
| Crowfoot Grass | Rescuegrass |
| | Sandspur |
| | Sedges* |

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner consistent with its labeling. Not for use on turf grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a Pre-Emergent AND also as a Post Emergent weed killer.

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 50 lb bag to treats 8,800 Sq Ft. — 25 lbs. bag treats 4,400 sq. ft. of lawn. (equivalent to 5.7 lbs. per 1,000 Sq Ft). Do not exceed recommended rates. Water immediately after application to wash granules off the grass blades.

HOW TO SET YOUR SPREADER

CYCLONE 4 SCOTT 6

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. Do not use on Bluegrass, Fescue, Bentgrass, Bahia, Rye, Bermuda, or Dichondra lawns. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	- - - - -	16%
Nitrate Nitrogen	0.00%	
Ammoniacal Nitrogen	11.20%	
Water Soluble Nitrogen (and/or Urea Nitrogen)	4.80%	
Water Insoluble Nitrogen	0.00%	
Available Phosphoric Acid (P ₂ O ₅)	- - - - -	4%
Soluble Potash (K ₂ O)	- - - - -	8%
Chlorine not more than	5.65%	

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium Potential Acidity Equivalent to 1245 lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - 1.20%; Iron as Fe - 1.00%; Water Soluble Magnesium Mg - 1.20%; Zinc as Zn - 0.05% Manganese as Mn - 0.50% Boron as B 0.02% Copper as Cu - 0.02%; Molybdenum as Mo - 0.002% Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate

NOTICE: Buyer and user assumes all risk of use, storage and/or handling of this material not in accordance with the terms of this label (Amdmt to remove from scope of WPS11-04-94)

ACTIVE INGREDIENT:
 Atrazine 2-chloro-4-ethylamino-6-isopropylamino-s-triazine - - - 0.76%
 Related Compounds - - - - - 0.04%
INERT INGREDIENTS: - - - - - 99.20%
 Total 100.00%

Contains 16 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

WEEDS CONTROLLED OR SUPPRESSED

- | | |
|-----------------------------------|---------------------------------------|
| Annual Bluegrass (Poa Annua) | Henbit |
| Annual Sowthistle | Knotweed |
| Clover | Lambsquarter |
| Black Medic | Lespedeza |
| Blue Toadflax | Lippia (Matchweed) |
| Bull Paspalum | Moneywort |
| Buckhorn Plantain | Morning Glory |
| Bur Clover | Narrowleaf Vetch |
| Carolina Geranium (Cranesbill) | Parsley-Piert |
| Chickweed (Common and Mouse Ear) | Oxalis (Common and Yellow Woodsorrel) |
| Cocklebur | Pigweed |
| Common Dandelion | Purslane |
| Common Purslane | Ragweed |
| Crabgrass | Red Sorrel |
| Crowfoot Grass | Rescuegrass |
| Cudweed | Sandspur |
| Cutleaf Evening Primrose | Sedges* |
| Dichondra | Shepards Purse |
| Dollar Weed | Sicklepod |
| Florida Beggarweed | Spotted Spurge |
| Florida Betony (Rattlesnake Weed) | Spurweed |
| Florida Pusley | Stinging Nettle |
| Fragrant Cudweed (Rabbit Tobacco) | Swinecrest |
| | Venus Looking Glass |
| | Virginia Creeper |
| | Wild Mustard |
| | Wood Sorrel |

*Repeat application are required for control of Nutsedge grasses

application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 50 lb bag to treats 8,800 Sq Ft. — 25 lbs. bag treats 4,400 sq ft. of lawn. (equivalent to 5.7 lbs. per 1,000 Sq.Ft). Do not exceed recommended rates. Water immediately after application to wash granules off the grass blades.

HOW TO SET YOUR SPREADER

CYCLONE 4 SCOTT 6

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. Do not use on Bluegrass, Fescue, Bentgrass, Bahia, Rye, Bermuda, or Dichondra lawns. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - -	16%
Nitrate Nitrogen	0.00%
Ammoniacal Nitrogen	11.20%
Water Soluble Nitrogen (and/or Urea Nitrogen)	4.80%
Water Insoluble Nitrogen	0.00%
Available Phosphoric Acid (P ₂ O ₅) - - -	4%
Soluble Potash (K ₂ O) - - - - -	8%
Chlorine not more than	3.65%

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate-Potash-Magnesium Potential Acidity Equivalent to 1245 lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - 1.20%; Iron as Fe - 1.00%; Water Soluble Magnesium Mg - 1.20%; Zinc as Zn - 0.05% Manganese as Mn - 0.50%; Boron as B 0.02% Copper as Cu - 0.02%; Molybdenum as Mo - 0.002%
 Derived from: Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate.

NOTICE: Buyer and user assumes all risk of use, storage and/or handling of this material not in accordance with the terms of this label.(Amdmt to remove from scope of WPS\11-04-94)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

EPA REG. NO. 35512- 14 EPA EST. NO. 35512-FL-1
 NET WEIGHT 50 POUNDS

HOWARD FERTILIZER COMPANY

P.O. BOX 13800 - ORLANDO, FL 32859