

35512-14

5/26/2004

Page 172



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAY 26 2004

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Rebecca M. Horton
Consultant for:
Howard Fertilizer & Chemical Company, Inc.
118 1/2 E. Main Street, Suite 1
Salem, VA 24153

Dear Ms. Horton:

Subject: Turf Pride Weeder
EPA Registration Number 35512-14
Application dated March 2, 2004

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended is acceptable, provided you make the following changes before you release the product for shipment.

1. Add an "If Inhaled" section to the First Aid statements after If Swallowed, with the following text:
If Inhaled:
-Move person to fresh air.
-If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.
-Call a poison control center or doctor for further treatment advice

Submit three (3) copies of final printed labeling incorporating the above changes before you release the product for shipment. Amended labeling will supercede all previously accepted ones. A stamped copy of labeling is enclosed for your records.

If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

A handwritten signature in black ink, appearing to read "James A. Tompkins".

James A. Tompkins
Product Manager 25
Herbicide Branch
Registration Division (7505C)

Contains 16 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

TURF PRIDE® is a Registered Trade Mark of Howard Fertilizer & Chemical Co., Inc.

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact x-xxx-xxx-xxxx for emergency medical treatment.

If Swallowed:	<ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annua)	Henbit
Annual Sowthistle	Knotweed
Clover	Lambsquarter
Black Medic	Lespedeza
Blue Toadflax	Lippia (Matchweed)
Bull Paspalum	Moneywort
Buckhorn Plantain	Morning Glory
Bur Clover	Narrowleaf Vetch
Carolina Geranium (Cranesbill)	Parsley-Plant
Chickweed (Common and Mouse Ear)	Oxalis (Common and Yellow Woodsorrel)
Cocklebur	Pigweed
Common Dandelion	Purslane
Common Purslane	Ragweed
Crabgrass	Red Sorrel
Crowfoot Grass	Rescuegrass
Cudweed	Sandspur
Cutleaf Evening Primrose	Sedges*
Dichondra	Shepards Purse
Dollar Weed	Sicklepod
Florida Beggarweed	Spotted Purge
Florida Betony (Rattlesnake Weed)	Spurweed
Florida Pusley	Stinging Nettle
Fragrant Cudweed (Rabbit Tobacco)	Swinecrest
	Venus Looking Glass
	Virginia Creeper
	Wild Mustard
	Wood Sorrel

*Repeat applications are required for control of Nutsedge grasses.

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAY 26 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

35512-14

Do not allow the watering-in to be complete and the surface is dry.
APPLICATION EQUIPMENT RESTRICTIONS
Do not apply granular lawn products by hand, do not allow product to come in contact with hands. Granular lawn products applied by hand-held devices (e.g., belly grinder) can only be applied as a spot treatment.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a Pre-Emergent AND also as a Post Emergent weed killer.

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Betony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 50 lb. bag to treat 8,800 Sq. Ft. - 25 lb. bag treats 4,400 Sq. Ft. of lawn. (equivalent to 5.7 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Water immediately after application to wash granules off the grass blades.

HOW TO SET YOUR SPREADER

CYCLONE 4 SCOTT 8

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. Do not use on Bluegrass, Fescue, Bentgrass, Bahia, Rye, Bermuda, or Dichondra lawns. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N).....	xx%
Nitrate Nitrogen	xx%
Ammoniacal Nitrogen	xx%
Water Soluble Nitrogen	xx%
(and/or Urea Nitrogen)	
Water Insoluble Nitrogen	xx%
Available Phosphoric Acid (P ₂ O ₅)....	xx%
Soluble Potash (K ₂ O).....	x%
Chlorine not more than	xx%

Derived from xxxxxxxx.

Potential Acidity Equivalent to xx lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

XX
F387

NOTICE: Buyer and user assumes all risk of use, storage and/or handling of this material not in accordance with the terms of this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

For Residential Use Product Disposal:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never placed unused product down any indoor or outdoor drain.

EPA REG. NO. 35512-14

EPA EST. NO. 35512-FL-1

NET WEIGHT 50 POUNDS (22.7 kg)

HOWARD FERTILIZER & CHEMICAL CO., INC.

P.O. BOX 628202 - ORLANDO, FL 32862-8202