

[2]

6. Submit three (3) copies of your final printed labeling, before you release the product for shipment. Refer to the A-79 previously mailed to you for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

BEST AVAILABLE COPY

CONCURRENCES							
SYMBOL	H7505C						
SURNAME	Robbins						
DATE	10/2/91						

WITH TRIMEC HERBICIDE

A TRIPLE ACTION BROADLEAF HERBICIDE WITH LAWN FOOD

50 lbs. TREATS 5,500 SQ.FT. of Turfgrass

For Use In Control of Broadleaf Weeds In Home Lawns, Apartment and Condominium Turf, In Parks, Playgrounds and other Ornamental & Recreational Turfgrass

A BROADLEAF WEED KILLER FOR SOUTHERN LAWNS

BALANCED LAWN FOOD GIVES TURF **FREE** WITH EXTENDED FEEDING

CONTROLS: Dandelions, Chickweed, Knotweed and Many Other Weeds as Listed For Use On St. Augustine Grass, Centipede, Zoysiagrass, Bahiagrass & Bermudagrass; Bluegrass, Rye Grass, and Fescue Varieties.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2, (2-methyl-4-chlorophenoxy) propionic acid 0.2400%
**Dimethylamine Salt of 2,4-Dichlorophenoxy acetic acid 0.0720%
***Dimethylamine Salt of Dicamba (3,6 dichloro-p-anisic) acid 0.0308%

INERT INGREDIENTS:

Total 100.0000%

*Equivalent to .2080% 2(2-methyl-4-chlorophenoxy propionic acid)
**Equivalent to .0600% 2,4-Dichlorophenoxy acetic acid
***Equivalent to .0255% (3,6 dichloro-p-anisic) acid (Isomers Specific by AQAC 14th Edition #6.321)

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF INHALED:** Remove victim to fresh air and apply artificial respiration if indicated. Get medical attention. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention immediately. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists.

See Side/Back Panel For Additional Precautionary Statements

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust. Wear face shield or goggles, long pants, long-sleeved shirt, shoes and gloves when mixing, loading or applying this product.

ENVIRONMENTAL HAZARD

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

RE-ENTRY & WORKER SAFETY STATEMENT

Do not re-enter or permit workers to reenter treated areas until dusts have settled.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **DISPOSAL:** Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

BROADLEAF WEEDS CONTROLLED

A PARTIAL LIST OF WEEDS CONTROLLED WITH THIS PRODUCT

Bedstraw	Croton	Knotweed	Pigweed	Vervain
Black Medic	Curly Dock	Koohia	Plantains	Velvetleaf
Blue Lettuce	Dandelion	Lambsquarters	Poleon Ivy/Oak	Vetch
Buckhorn	English Daisy	Lespedeza	Povertyweed	Wild Carrot
Bull Thistle	Fleabane	Mallow, Venice	Purslane	Wild Garlic
Burdock	Florida Pusley	Morning glory	Ragweed	Wild Lettuce
Buttercup	Ground Ivy	Mustard	Shepherdspurse	Wild Onion
Chickweed	Hawthorn	Oxalis	Sheep Sorrel	Yarrow
Clover	Heelall	Pennywort	Smartweed	
Cookiebur	Henbit	Peppergress	Spurge	

and many other Broadleaf Weeds

This product contains a Complete Lawn Food and TRIMEC Herbicide, which gives triple action Broadleaf Weed control.

WHERE TO USE

This product may be used on all varieties of St. Augustine Grass, Zoysiagrass, Bermudagrass, Bahiagrass, and Centipede Grass, Perennial and Annual Varieties of Rye, Rough and Fine leaved Fescues, and Bluegrass. It should not be used where desirable clovers are present. Apply at a time when the air is calm. Do not use on newly seeded lawns until after the second cutting. After application delay seeding for at least three weeks and increase the rate of seed put down by 25%.

WHEN TO USE

Apply when weeds are young and actively growing in the spring, summer, or fall. Best results will be obtained when temperatures range from 65 to 90 degrees F. and weeds are in a vigorous state of growth from rain and/or watering. Results will be erratic if applied during drought periods. Apply when the air is calm. Lawns that are heavily over-run by hard-to-kill weeds will benefit from an additional treatment in 30 to 45 days.

HOW TO USE

Select an appropriate Spreader and distribute 50 lbs. per 5,500 Sq. Ft. (9.1 lbs./1,000 Sq. Ft.) of turf for control of the many weeds as listed and to supply the lawn with a complete NPK fertilization at the same time. This application applies a total of 1.2 lbs. herbicide per acre.

HOW TO MAKE SURE YOUR SPREADER IS OPERATING CORRECTLY

Using a Scott Spreader for broadleaf weeds place 4.5 lbs. granules in your spreader using No. 7 setting and apply to 500 Sq. Ft. With a 20 inch spreader this will equal 3 spreader widths wide (3 passes) by 100 feet long. If you run out before you cover the pre-measure area close the spreader setting by 1/2 the number. If you have granules left over, open (increase) the setting by 1/2 the number. For further help refer to your Spreader Guide Book.

Poor weed control will result if your spreader applies less than recommended. Make sure to apply the total pounds per sq. ft. as shown in the following chart.

SPREADER SETTINGS		
SPREADER	AMOUNT TO APPLY PER 5,500 SQ. FT.	SETTING
Scott	50 lbs.	#7
Cyclone	50 lbs.	#4

When using a 25 lb. bag, spreader settings will be the same as for the 50 lb. bag but it will treat half the Sq. Ft. These settings are based on a One Time Over application with a New spreader.

GUARANTEED FERTILIZER ANALYSIS

Nitrogen (N) variable %
Sulfur Coated Urea variable %
Ammoniated Nitrogen variable %
Available Phosphoric Acid (P₂O₅) variable %
Soluble Potash (K₂O) variable %
Derived from Ammonium Sulfate, Sulfur Coated Urea, Ammoniated Phosphate, Muriate of Potash (as variable fertilizer inerts)

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label (09-19-91 revision of 04-05-91 version)

EPA REG. NO. 35512-EN(20)

EPA EST. NO. 35512-FL-1

Net Weight 50 Pounds

HOWARD FERTILIZER COMPANY

ORLANDO, FL 32859-3800

BEST AVAILABLE COPY

