

Jacket

TURF PRIDE

with .8% OFTANOL INSECTICIDE

50 lbs. TREATS UP TO 17,000 SQ.FT. OF TURF
CONTAINS 0.8% OFTANOL INSECTICIDE + LAWN FOOD

FOR CHINCHBUGS, SOD WEBWORMS,
BILLBUG LARVAE, WHITE GRUB LARVAE,
AND FIRE ANTS

ACTIVE INGREDIENT:

1-Methylethyl 2-[(ethoxy(1-methylethyl)- amino)phosphinothioyl]oxy)benzoate*.....	0.8%
INERT INGREDIENTS:	99.2%
Total	100.0%

*U.S Patent No. 3,621,082 Canadian Patent No. 828,198 Orlando is a Reg. TM of the Parent Company of Fertilizatoren Bayer GmbH, Leverkusen

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention **IF ON SKIN:** Wash affected area with soap and warm water. **IF INHALED:** Remove to fresh air.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. May cause eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Wash contaminated clothing with soap and hot water before re-use. Keep children and pets off treated areas until insecticide has been washed to soil level and grass is dry.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwater. This pesticide is toxic to fish and wildlife. Do not apply directly to water or wetlands. Keep out of lakes, streams and ponds.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original container in a cool, dry, isolated place out of reach of children. PESTICIDE DISPOSAL: (except household) Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: (except household) Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. Burned stay out of smoke. HOUSEHOLD: Do not re-use empty bag. Wrap and put in trash.

NOTICE: Buyer and user assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	%
% Ammoniacal Nitrogen	
% Sulfur Coated Urea Nitrogen	
Available Phosphoric Acid (P_2O_5)	%
Soluble Potash (K_2O)	%

Derived from: Ammonium Sulfate, Sulfur Coated Urea, Ammonium Phosphate and Muriate of Potash. 4% of the Nitrogen source is sulfur coated to provide % slow release nitrogen.
Calcium Carbonate equivalent lbs./ton.

NET WEIGHT 50 POUNDS

EPA REG. NO. 35512-22

EPA EST. NO. 35512-FL-1

SURFACE FEEDING INSECTS

CHINCHBUGS, SOD WEBWORM, IMPORTED FIRE ANTS
USE 50 POUNDS TO TREAT 8,750 TO 17,000 SQ.FT. OF TURF
(See the higher rate to treat 8,750 sq.ft. for Severe Infestations)

CHINCH BUGS AND SOD WEBWORM LARVAE

Apply 2.05 to 5.7 lbs. / 1,000 Sq.Ft. when insect damage is first observed and immediately apply enough water to wet the soil 1/4 to 1/2 inches deep without causing run-off. Repeat as necessary for pest infestations. Restrictions - Do not apply more than 11.25 lbs./1,000 Sq.Ft. per year or an equivalent amount of 1.48 ounces of Active Ingredient per 1,000 Sq.Ft. regardless of formulation used. Minimum interval between applications must be 30 days.

IMPORTED FIRE ANT

Area Fire Ant Control- Apply 5.7 lbs./1,000 Sq.Ft. of turf uniformly over the area to be treated for isolated control. Water immediately after application using 1/2 to 1 inch of water. If established Fire Ant Mounds are present use the mound drench treatment prior to area application. If a mound drench treatment is made only one area treatment may be made.

ESTABLISHED MOUNDS DRENCH TREATMENTS

WET CONDITIONS-Treat 1 ounce of Ortho per 6 Sq.Ft. & immediately spike gently with 3 gallons of water.

DRY CONDITIONS-Treat as noted for wet conditions, however, to insure penetration apply an additional 3 gallons of water to the treated mound area after treatment. Make only one mound drench treatment per season.

ROOT FEEDING INSECTS

USE 50 POUNDS TO TREAT 8,750 SQ.FT. OF TURF

BILL BUG LARVAE, HYPERODES WEEVIL LARVAE, WHITE GRUB LARVAE such as: CHAFERS; PHYLOPHAGA spp.; BLACK TURFGRASS ATAENIUS, GREEN JUNE BEETLE, JAPANESE BEETLE, ORIENTAL BEETLE, ASIATIC GARDEN BEETLE.

For best results, follow the timing and mixing instructions below for the specific insects you need to control. Apply at the rate of 5.7 lbs. per 1,000 Sq.Ft. using the recommended spreader settings.

BILL BUG LARVAE

Apply 5.7 lbs./1,000 sq.ft. when adult activity is observed in the spring, usually late March to Mid-May, and immediately apply enough water to wet the soil to 1 to 1 1/2 " deep. Avoid causing run-off.

HYPEROIDES WEEVIL LARVAE

Apply 5.7 lbs./1,000 sq.ft. in the spring between full bloom periods of Forsythia and Flowering Dogwood, usually April 15 to May 15, and immediately apply enough water to wet the soil to 1 to 1 1/2 inches deep without causing run-off.

WHITE GRUB LARVAE: (Japanese Beetle, Black Turf Grass Attaenius, Chafer, Phylophaga spp., Green June Beetle, Oriental Beetle and Asiatic Garden Beetle).

FIRST-EVENTIVE APPLICATIONS: Apply 5.7 lbs./1,000 sq.ft. in the spring (April through Mid-May) and immediately apply enough water to wet the soil 1 to 1 1/2 inches deep without causing run-off.

CUMULATIVE APPLICATION: Apply 5.7 lbs./1,000 sq.ft. when visible damage or High Grub populations have been observed and immediately apply enough water to wet the soil to 1 to 1 1/2 inches deep without causing run-off.

SPREADERS AND SETTINGS

For the 5.7# Rate / 1,000 sq.ft.

CYCLONE - No. 4; Ortho (Rotary) - No. 4

Ortho (Drop) - 6.5 Scotts PF-1 - No. 5.5

For 2.85# Rate reduce setting by 1 full number. Settings may vary with the age and condition of your spreader. Ask your dealer for settings to use on other brands of spreaders.

IMPORTANT RESTRICTIONS

For Billbugs, Hyperodes Weevils and White Grub Larvae do not apply more than 11.25 lbs. of this product/1,000 Sq.Ft. per year or an equivalent amount of 1.48 ounces of Active Ingredient per 1,000 Sq.Ft. regardless of formulation used. Minimum interval between applications must be 30 days.

Do not collect clippings after first mowing date/dethatching. Do not grow treated areas or use clippings from treated areas for feed. Do not allow material to come in intimate contact with bare skin. Do not allow children and pets on treated area until foliage is dry. Do not apply more than twice per year. Do not mow treated turf areas before mowed or irrigation has been applied as per recommendations.

APPLICATION EQUIPMENT

For best results apply uniformly with a spreader designed to apply granular herbicides and insecticides to turf. Both drop type and rotary type spreaders may be used. Avoid using spreaders that apply in narrow concentrated bands. Calibrate equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and is distributing granules uniformly.

APPLICATION TECHNIQUES

1. For maximum benefit apply the full recommended amount. Apply when the wind is calm to prevent missed areas. Follow watering instructions after applying.
Name Change And. 07-01-01)

HOWARD FERTILIZER COMPANY
P. O. BOX 13800 - ORLANDO, FL 32859

