

Reg # 35512-25. 7:29 PM-23 181

TURF PRIDE. with 1.50% RONSTAR.

50 lb. Bag Treats up to 16,300 Sq.Ft.

Ask for a Copy of the Supplemental Label Available For The Use Of This Product On Ornamental Shrubs, Vines & Trees

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY

FOR COMMERCIAL USE ON PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS

A PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, BERMUDAGRASS, TALL FESCUE & ZOYSIAGRASS

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed & Other Weeds

ACTIVE INGREDIENT:

Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) 1,2,1,3,4-oxadiazolin-5-one]. 1.50%

INERT INGREDIENTS: 98.50%
Total 100.00%

Contains RONSTAR® Herbicide

a Registered Trade Mark of Rhone Poulenc Inc.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water & touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap & water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA Reg. No. 35512-25 EPA EST. NO. 35512-FL-1
NET WEIGHT 50 POUNDS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt & gloves when handling. Launder exposed clothing before reuse. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

FOR LAWNS AND ORNAMENTAL TURF

NOT FOR USE ON HOME LAWNS. FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF & LANDSCAPE PERSONNEL ONLY

When applied according to Directions this product can be used safely on the following established turfgrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS VARIETIES, ST. AUGUSTINE GRASS VARIETIES, TALL FESCUE, & ZOYSIAGRASS lawns & ornamental turf. It may be used for Apartment and Condominium complexes, Parks, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turfgrass areas.

NOTIFICATION CONTROLS GRASSY WEEDS & BROADLEAF WEEDS

Bluegrass (Poa Annual), Barnyard Grass, Crabgrass, Fall Florida Pusley, Goose grass, Green Foxtail, Oxalis, Stinging Nettle, Carpet grass, Pigweed.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not overlap spreader applications, on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After applications if rain is not expected within 24 hours sprinkle the treated turf with 1/4 to 1/2 inch of water to get maximum weed control.

HOW MUCH TO USE

FOR CRABGRASS: Apply 50 lbs./16,300 Sq.Ft. when turf has been in a yearly maintenance program (equal to 2 lbs. A.I. herbicide per acre.) Apply 50 lbs. to treat 11,000 Sq.Ft. of lawn if it has been heavily infested with Crabgrass; equal to 3 lb. A.I. herbicide per acre. Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When To Apply" Chart.

GOOSEGRASS & PROSTRATE SPRUGE: These weeds do not usually germinate until late spring or early summer. Apply 50 lbs. to treat 8,350 Sq.Ft. (4 lbs. A.I./acre) & apply 1 to 2 weeks prior to seed germination. See Chart for When To Apply. Observe maximum rate for St. Augustine grass.

FOR ANNUAL BLUEGRASS (POA ANNUA): Apply 50 lbs. per 8,350 to 9,675 Sq.Ft. of Lawn; equal to 3.5 to 4 lbs. active herbicide per acre. Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control annual bluegrass present at time of application; therefore only germinating seeds. Refer to chart for proper timing in your locale. Annual BARNYARDGRASS and PROSTRATE SPURGE may also be controlled by this same rate. Observe maximum rate for St. Augustine grass.

FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CARPETGRASS: Where these weeds are a problem apply 50 lbs. to treat 8,350 to 9,675 Sq.Ft. of Lawn; (equal to 3.5 to 4 lbs. Active Herbicide per acre) In early spring or 1 to 2 weeks before expected weed seed germination. Will not control of emerged and growing weeds; only effective against germinating weed seeds. Observe maximum rate for St. Augustine grass.

SPREADER SETTINGS FOR SQUARE FOOT COVERAGE

	AMT. OF for 16,300 Sq.Ft.	1,000 Sq.Ft.	8,350 Sq.Ft.
SPREADER PRODUCT	=2#A.I./Acra	=3#A.I./Acra	=4#A.I./Acra
CYCLONE B-1 50 LBS.	3 1/2	4 1/2	5 1/2
SCOTTS #35 50 LBS.	3 1/2	4	4 1/2

NOTE: Spreader rate varies with walking speed
Check spreader over known area.

NOTICE: Buyer and user assume all risk and liability for use of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N).....	38%
Nitrate Nitrogen	0%
Ammoniacal Nitrogen	0%
Water Soluble Nitrogen	
and/or Urea Nitrogen	10%
Water Insoluble Nitrogen	28%
Available Phosphoric Acid (P ₂ O ₅)....	0%
Soluble Potash (K ₂ O).....	0%

Derived from Methylene Diurea

(A Urea Formaldehyde Nitrogen Fertilizer Source)

or handling of this product not in accordance with the terms of the label.

WHEN TO APPLY

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

JAN. FEB. MAR. APR. MAY JUNE JULY AUG. SEPT. OCT. NOV. DEC.

1. For CRABGRASS & other Grasses & Broadleaf Weeds germinating in the Spring in Northern turf.
2. For GOOSEGRASS & other Late Germinating Annual Grasses & Broadleaf Weeds such as PROSTRATE SPURGE. 2a. GOOSEGRASS in Southern Turf.
3. For POA ANNUA in Northern Turf. 3a. For POA ANNUA in Southern Turf.
4. For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.

PRECAUTIONS IN USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Centipede grass since damage will occur. Do not use on Putting greens or "Tees" due to varying cultural practices used. Use only the rate of 50 lbs. to treat 11,000 Sq.Ft. on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended on the Ornamental Use Supplemental Labeling available for this product. For newly seeded lawns do not apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

06-26-91

HOWARD FERTILIZER COMPANY

P.O. BOX 593800 - ORLANDO, FL 32859-3800