

PM 23

35512-26

1 of 3

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: ☒ REGISTRATION ☐ REREISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended.)</i>	EPA REGISTRATION NO. 35512-26 DATE OF ISSUANCE JAN 5 1990 TERM OF ISSUANCE Indefinite NAME OF PESTICIDE PRODUCT Diflufenican 15-0-12.5 (100% EC)	
NAME AND ADDRESS OF REGISTRANT (Include ZIP Code) ROVER, JAMES E. JR. P.O. Box 1000 Orlando, FL 32812		
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.		
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.		
Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by other.		
1. Submit one (1) copy of your final printed labeling before you release the product for shipment. 2. Delete all references to previously issued EPA Registration Numbers.		
3. Submit one (1) copy of your final printed labeling before you release the product for shipment.		
☐ ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL <i>[Signature]</i>		DATE 1/5/90

EPA Form 8570-6 (Rev. 5-78)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED

58839:I:CR-31:Ikeda:MF-15:KENCO:01/04/90:02/05/90:CL:sw:vo:ek:dg

-2-

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Acting Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosure

TURF PRIDE. 12-0-12 with RONSTAR.

50 lb. Bag Treats up to 8,150 Sq Ft.
See Supplemental Label For Use On Ornamental Shrubs, Vines & Trees

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY,
TURF AND LANDSCAPE PERSONNEL ONLY

A. PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR
PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS
ST. AUGUSTINE GRASS, BERMUDAGRASS
TALL FESCUE & ZOYSA

FOR COMMERCIAL USE ON PARKS, GOLF COURSES, ATHLETIC
FIELDS AND OTHER RECREATIONAL AND ORNAMENTAL
TURFGRASS AREAS

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley,
Oxalis, Stinging Nettle,
Carpetgrass, Pigweed and Other Weeds as Listed

ACTIVE INGREDIENT

Oxadiazon: [2-tert-butyl-4-(2,4-dichloro-5-
isopropoxyphenyl)-6-(2,1,3,4-oxadiazol-5-yl)] 0.75%
INERT INGREDIENTS 99.25%

Total 100.00%

Contains RONSTAR® herbicide
a Registered Trade Mark of Rhone-Poulenc Inc.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately.
Induce vomiting by giving victim 1 or 2 glasses of water & touching back
of throat with finger. Do not induce vomiting or give anything by mouth to
an unconscious or comatose person. IF IN EYES: Flush eyes with plenty
of water. Get medical attention if irritation persists. IF ON SKIN: Wash with
plenty of soap & water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA Reg. No. 35512- EPA EST. NO. 35512-FL-1
NET WEIGHT 50 POUNDS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes
or clothing. May be fatal if inhaled. Thoroughly wash face and hands with
soap and water after exposure and before eating or smoking. Wear
protective clothing including long-sleeved shirt & gloves when handling.
Laundry exposed clothing before reuse. Do not allow domestic animals
to graze treated areas. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not apply
when weather conditions favor drift from treated areas. Do not apply where
runoff is likely to occur. Do not contaminate water when disposing of
equipment washwaters. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsis-
tent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in its original bag in a cool, dry, locked place
out of reach of children. In case of spill, sweep up, and use according to
directions or dispose of as follows:

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of
excess pesticide, or residue is a violation of Federal law. If these wastes
cannot be disposed of by use according to label instructions, contact your
State Pesticide or Environmental Control Agency, or the Hazardous Waste
representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equip-
ment. Then dispose of empty bag in a sanitary landfill or by incineration,
or, if allowed by State and local authorities, by burning. If burned stay out
of smoke.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	12%
Nitrate Nitrogen	0%
Ammoniacal Nitrogen	12%
Water Soluble Nitrogen	
and/or Urea Nitrogen	0%
Water Insoluble Nitrogen	0%
Available Phosphoric Acid (P ₂ O ₅)	0%
Soluble Potash (K ₂ O)	12%

Derived from Ammonium Sulfate & Muriate of Potash

Potential Acidity Equivalent to 1257 lbs. of Calcium Carbonate / ton

See Supplemental Label For Use On Ornamental Shrubs, Vines & Trees

FOR LAWNS AND ORNAMENTAL TURF

NOT FOR USE ON HOME LAWNS. FOR SALE TO, USE AND STORAGE
BY COMMERCIAL NURSERY, TURF & LANDSCAPE PERSONNEL ONLY

When applied according to Directions this product can be used safely on the
following established turfgrasses varieties: BERMUDAGRASS, PERENNIAL
BLUEGRASS, PERENNIAL RYEGRASS VARIETIES, ST. AUGUSTINE
GRASS VARIETIES, TALL FESCUE, & ZOYSA. This product is for use on established
turf. It may be used for Apartment and Condominium complexes, Parks, Golf
Courses, Athletic Fields, Communities and all other ornamental and recrea-
tional turfgrass areas.

PRECAUTIONS IN USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or
Centipede grass since damage will occur. Do not use on Putting greens or
"Tees" due to varying cultural practices used. Use only the rate of 50 lbs. to
treat 8,150 Sq. Ft. on St. Augustinegrass since temporary discoloration will
occur from heavier rates. For flowers and shrubs see Supplemental Label-
ing available for this product. For newly seeded lawns do not apply this
product until after the second or third cutting. For best lawn care, place new
grass seed in the fall of the year. Spring pre-emergence weed control can
then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

CONTROLS GRASSY WEEDS & BROADLEAF WEEDS

Annual Bluegrass (Poa Annua), Barnyard Grass, Crabgrass, all Poa, all
Florida Pusley, Goose grass, Green Foxtail, Oxalis, Stinging Nettle,
Carpet grass, Pigweed.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if
necessary so that granules will reach the soil surface. Measure the area to be
treated and uniformly apply the required amount using a fertilizer spreader
(see Spreader Settings chart). Do not apply to turf when it is wet. Do not over-
lap spreader applications, on St. Augustinegrass since discoloration "bleech-
ing" may temporarily occur from "double rate". After applications if rain is not
expected within 24 hours sprinkle the treated turf with 1/4 to 1/2 inch of water
to get maximum weed control.

HOW MUCH TO USE

FOR CRABGRASS Apply 50 lbs. / 8,150 Sq. Ft. when turf has been in a yearly
maintenance program (equal to 2 lbs. x 1 herbicide per acre). Apply 50 lbs.
to treat 8,150 Sq. Ft. of lawn if it has been heavily infested with Crabgrass
equal to 3 lb. A.I. herbicide per acre. Apply in early spring 1 to 2 weeks prior
to weed seed germination. This product is a pre-emergence herbicide. It will
not control emerged and growing weeds, only germinating weed seeds. Late
applications will control only those weed seeds that germinate after the
application. See Chart.

GOOSEGRASS & PROSTRATE SPURGE These weeds do not usually
germinate until late spring or early summer. Apply 50 lbs. to treat 8,150 Sq. Ft.
(4 lbs. A.I./acre) & apply 1 to 2 weeks prior to seed germination. See Chart for
When To Apply. Observe maximum rate for St. Augustine grass.

FOR ANNUAL BLUEGRASS (POA ANNUA) Apply 50 lbs. per 4,100 to 8,150
Sq. Ft. of lawn, equal to 3.5 to 4 lbs. active herbicide per acre. Apply in late
summer to fall or in early spring 1 to 2 weeks before expected weed seed
germination. Will not control annual bluegrass present at time of application,
therefore only germinating seeds. Refer to chart for proper timing in your
locale. Annual BARNYARDGRASS and PROSTRATE SPURGE may also be
controlled by the same rate. Observe maximum rate for St. Augustine grass.
FOR FLOODED FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CAR-
PETGRASS Where these weeds are a problem apply 50 lbs. to treat 4,100
to 8,150 Sq. Ft. of lawn, (equal to 3.5 to 4 lbs. Active Herbicide per acre) in
early spring or 1 to 2 weeks before expected weed seed germination. Will not
control of emerged and growing weeds, only effective against germinating
weed seeds. Observe maximum rate for St. Augustine grass.

SPREADER SETTINGS FOR SQUARE FOOT COVERAGE

AMT OF 8,150 Sq. Ft. 8,400 Sq. Ft. 4,100 Sq. Ft.
SPREADER PRODUCT = 2# A.I./Acre = 3# A.I./Acre = 4# A.I./Acre

CYCLONE B-1	50 LBS.	6 1/2	6 1/2	7 1/2
SCOTTS #35	50 LBS.	4 1/2	5	5 1/2

NOTE: Spreader rate varies with walking speed

Check spreader over known area.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or
handling of this product not in accordance with the terms of this label.

WHEN TO APPLY

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

JAN FEB MAR APR MAY JUNE JULY AUG SEPT OCT NOV DEC

1. For CRABGRASS & other Grasses & Broadleaf weeds germinating in the Spring in Northern turf.
2. For GOOSEGRASS & other Late Germinating Annuals & Broadleaf Weeds such as PROSTRATE SPURGE in Southern Turf.
3. For POA ANNUA in Northern Turf. 3a. For POA ANNUA in Southern Turf.
4. For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.

12-01-88

HOWARD FERTILIZER COMPANY

P.O. BOX 593800 - ORLANDO, FL 32859-3800

35512-26