

TURF PRIDE. with .75% RONSTAR.

SUPPLEMENTAL LABELING

FOR WOODY ORNAMENTAL SHRUBS, VINES, TREES AND CONTAINER GROWN ORNAMENTALS

50 lb. Bag Treats up to 8,150 Sq.Ft.

A Commercial Weed & Feed "Preemergent + Fertilizer"
FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY,
TURF AND LANDSCAPE PERSONNEL ONLY

SEE BASIC TURF PRIDE® PRODUCT LABEL EPA REG NO. 35512-
FOR COMMERCIAL USE ON PARKS, GOLF COURSES, ATHLETIC
FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL
TURFGRASS AREAS

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley,
Oxalis, Stinging Nettle,
Carpetgrass, Pigweed and Many Other Weeds As Listed

ACTIVE INGREDIENT:

Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-
isopropoxyphenyl) 1,2-1,3,4-oxadiazolin-5-one] 0.75%

INERT INGREDIENTS: 99.25%
Total 100.00%

Contains RONSTAR® Herbicide a Registered Trade Mark of
Rhone Poulenc Inc.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately.
Induce vomiting by giving victim 1 or 2 glasses of water & touching back
of throat with finger. Do not induce vomiting or give anything by mouth to
an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if
irritation persists.

IF ON SKIN: Wash with plenty of soap & water. Get medical attention if
irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA Reg. No. 35512-26 EPA EST. NO. 35512-FL-1
NET WEIGHT 50 POUNDS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes
or clothing. Thoroughly wash face and hands with soap and water after
exposure and before eating or smoking. Wear protective clothing includ-
ing long-sleeved shirt & gloves when handling. Launder exposed clothing
before reuse. Do not allow domestic animals to graze treated areas. Avoid
contamination of Feed and foodstuffs.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not apply
when weather conditions favor drift from areas treated. Do not apply where
runoff is likely to occur. Do not contaminate water when disposing of
equipment washwaters. Apply only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsis-
tent with its labeling

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in its original bag, in a cool, dry, locked place
out of reach of children. In case of spill, sweep up, and use according to

WEED CONTROL AROUND WOODY ORNAMENTAL SHRUBS, VINES & TREES CONTROLS GRASSY WEEDS & BROADLEAF WEEDS

Annual Bluegrass, Barnyardgrass, Bittercress, Carpetweed, Carpetgrass,
Common Groundsel, Common Purslane, Crabgrass, *Evening Primrose,
Green Foxtail, Fall Panicum, Florida Pusley, *Galinsoga, Golden Ragwort,
*Goosegrass, Lambsquarter, Oxalis, Pennsylvania Smartweed, Petty
Spurge, *Prostrate Spurge, Redroot Pigweed, Spotted Catsear, Stinging
Nettle, *Swinecress, Yellow Woodsorrel.

Can be applied to both newly transplanted and established ornamentals
plants where both weed control and fertilization is desired. Rainfall or
overhead irrigation will improve weed control. Do not disturb or cultivate soil
after treatment. Do not apply to wet foliage or under conditions in which
granules will collect on leaves.

HOW MUCH TO USE

For susceptible weeds apply 50 lbs. to treat 8,150 to 5,400 Sq.Ft. of Orna-
mental planted area. This supplies plants with approximately 0.74 to 1.11
lbs. actual (N) Nitrogen plant food per 1,000 Sq.Ft. of plant bed See Chart
WHEN TO APPLY. (50 lbs./5,400 Sq.Ft. is preferred rate.)

*For hard to kill weeds as *indicated use the rate of 50 lbs. for 5,400 Sq.Ft.
or 50 lbs. for 4,100 Sq.Ft. (3 to 4 lbs. A.I./Acre) = 6.13 lbs. to 9.2 lbs. actual
product per 1000 Sq. Ft. of plant bed.

FOR AREAS WHERE WEEDS ARE GROWING

If treatment is desired in non-turf areas where weeds are presently growing
first remove existing weeds. Rake and smooth the soil by "breaking-up"
clumps. Apply at rate as directed for specific weeds. (50 lbs. to treat either
8,150 Sq. Ft. or 5,400 Sq.Ft.) This treatment will only control weeds germi-
nating from seeds. It will not control weeds sprouting from roots! Irrigate the
area after treatment with 1/4 to 1/2 inch water to get maximum weed control.

FOR FIELD GROWN ORNAMENTALS

*Apply at the rate of 50 lbs. per 8,150 or 5,400 Sq.Ft. of area.

This product may be used around the following Established Plants. ARBOR-
VITA, ASH, AZALEA, BARBERRY-Japanese, BIRCH, CAMELLIA,
COTONEASTER, CRABAPPLE, DOGWOOD, EUONYMOUS, FORSYTHIA,
HOLLY, HONEYSUCKLE, JUNIPER, LILAC, MAPLE, OAK, PERIWINKLE,
PINE, PITTOSPORUM, PRIVET, RHODODENDRON, ROSE, RUSSIAN OL-
IVE, SPRUCE, TAXUS, VIBURNUM, XYLOSMA, GROUND COVERS: CAR-
POBROTUS, OSTEOSPERMUM, SEDUM, ALGERIAN IVY, VINCA MAJOR,
ENGLISH IVY, PACHASANDRA, AJUGA, EUONYMOUS.

*Use higher rate for hard to kill weeds.

May be used around the following newly transplanted ornamentals.
CAMELLIA, DOGWOOD, FORSYTHIA, HEDRA, HOLLY, JUNIPER, OAK,
PERIWINKLE, PINE, PRIVET, RHODODENDRON, RUSSIAN OLIVE, SPRUCE,
TAXUS, TSUGA.

To obtain continued season long weed control on the following list of plants
additional applications may be made 90 to 120 days after the previous
application in the following situations:

MAY USE TWO REPEAT TREATMENTS (50 lbs / 5,400 SQ.FT)

ON FOLLOWING PLANTS

Established - Container Grown - Abelia, Common Boxwood, English Ivy,
Holly, Juniper, Pine, Rhododendron. Newly Transplanted - Container Grown
Common Boxwood, English Ivy, Juniper, Pine, Rhododendron.

MAY USE THREE REPEAT TREATMENTS (50 lbs / 5,400 SQ.FT.)

ON FOLLOWING PLANTS

Established - Container Grown - Abelia, Holly, Rhododendron.
Forest Nurseries - For Pines - Loblolly, Slash, Eastern White.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen(N)		12%
Nitrate Nitrogen	0.0%	
Ammonical Nitrogen	12.0%	
Water Soluble Nitrogen		
(and/or Urea Nitrogen	0.00%	
Water Insoluble Nitrogen	0.00%	
Available Phosphoric Acid(P ₂ O ₅)	0%	
Soluble Potash(K ₂ O)	12%	

Derived from Ammonium Sulphate and Muriate of Potash

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label.

WEED AND FEED FOR CONTAINER GROWN ORNAMENTAL PLANTS

This product is recommended on a wide variety of container and field grown woody ornamental shrubs, vines and trees. It may be used on the following list of container grown plants. Use at the rate of 50 lbs. per 4,100 sq.ft. of pot soil surface area. Plants in 14 inch containers set out in 12 foot wide rows X 100 feet long will require 11 1/2 lbs. product per row (Equivalent to 4 lbs. A.I. per acre). Apply with a "shoulder strap held" grass-seeder or granule applicator that will direct the granules to each pot soil surface.

AGAPANTHES, ANDROMEDA, ARBORVITA, ARCTOSTAPHYLOS, AZALEA, BARBERRY, BOXWOOD- Common-Japanese, CALISTEMON, CAMELLIA, CEANOTHUS, CEDAR, COTONEASTER, CYPRESS, DOGWOOD, DOUGLAS FIR, EUCALYPTUS, EUCNYMOUS-Evergreen, FORTSYTHIA, GARDENIA, GINKO, HEATH HEATHER, HEDERA, HOLLY, HONEYSUCKLE, JASMINE, JUNIPER-Varieties, JUSTICA, LEUCOTHOE, MAGNOLIA, MAHONIA, MYRTLE, OSMANTHUS, PERIWINKLE, PHOTINIA, PINE-Varieties, PITTOSPORUM, PODOCARPUS, PRIVET, PYRACANTHA, QUERCUS, RAPHIOLEPUS, RHODODENDRON, ROSE, STAR JASMINE, SYZYGIIUM, VIBURNUM, XYLOSMA.

06-26-91

HOWARD FERTILIZER COMPANY
P.O. BOX 593800 - ORLANDO, FL 32859-3800