177



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

TEB 23 1994

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

H. R. McLane

H.R. MCLANE INC.

Agent for: HOWARD FERTILIZER CO. INC.

7210 Red Road - Suite 206

Miami, FL 33143

Subject:

Label Amendment Submission of 10/25/93 in Response to PR Notice 93-7

EPA Reg. No. 35512-31

TURF PRIDE WITH 0.5% RONSTAR

IMPORTANT NOTICE:

This Letter Requires You to Submit a Notification to the Office

of Compliance Monitoring within 20 Days.

Dear Registrant:


The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments listed on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

SERIOUS ERROR(S) ON SUBMITTED LABELING:

EPA has determined that one or more of the errors on the labeling you submitted to the Agency is categorized as "serious" as defined in PR Notice 93-11. In general, serious errors are those that may create a potential for harm to workers, handlers, other persons, or the environment; or those that prevent the achievement of the basic goals of the Worker Protection Standard (WPS) or FIFRA.

YOU MUST NOT SELL OR DISTRIBUTE (INCLUDING RELEASE FOR SHIPMENT) ANY PRODUCT BEARING THE SUBMITTED LABELING. Sale or distribution of any product with the incorrect labeling is a violation of FIFRA that could result in initiation of an action for civil penalties and/or cancellation of your product. This matter has been referred to the Office of Compliance Monitoring (OCM) for further action.

YOU MUST SUBMIT A NOTIFICATION TO THE OFFICE OF COMPLIANCE MONITORING WITHIN 20 DAYS OF THE DATE OF THIS LETTER.



Page 2

Each notification must include the following information:

- 1. Registrant's name and address.
- 2. EPA Registration number of the product for which the labeling was submitted.
- 3. Whether any product already has been sold or distributed (this includes released for shipment) bearing the submitted labeling.

In addition, the notification must include the following information, IF you have already sold or distributed any product bearing the submitted labeling:

- 1. Quantity of the product that has been sold or distributed.
- 2. EPA Producing Establishment number.
- 3. Location of the product, if known.
- 4. Proposed plan for assuring that product with the incorrect labeling in your control, as well as that in the hands of dealers/distributors/users, will be corrected. The plan must include your proposal for assuring that the labeling is corrected as soon as possible.

Submit the notification to:

Michael Wood, Director
Compliance Division, (7203)
Worker Protection Notification
Office of Compliance Monitoring
U.S. Environmental Protection Agency
401 M Street SW
Washington, DC 20460

Failure to submit a notification also may result in initiation of an action under FIFRA for civil penalties, if you sold or distributed (this includes released for shipment) product bearing the incorrect labeling.

Page 3

IN ADDITION, YOU SHOULD CONTACT IMMEDIATELY THE PERSONS TO WHOM YOU SOLD OR DISTRIBUTED YOUR PRODUCT AND INFORM THEM NOT TO SELL OR DISTRIBUTE ANY PRODUCT WITH THE INCORRECT LABELING.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the WPS labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have neither been reviewed nor accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Submit the required notification. Then by the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling

 BEFORE selling or distributing any product bearing the final printed labeling as amended

AND

• WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

BEST AVAILABLE COP

Page 4

Hand or courier deliveries of final labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mail 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

Jim Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Attachment

cc: Office of Compliance Monitoring

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division

H. R. McLane H.R. MCLANE INC AGENT FOR: HOWARD FERTILIZER CO INC 7210 RED RD, SUITE 206 MIAMI FL 33143

Comment for: EPA Reg Nr.35512-31
TURF PRIDE WITH 0.5% RONSTAR

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

You have incorrect wording in the first paragraph in the Agricultural Use Requirements box on your proposed label. That paragraph must contain the exact wording of the paragraph shown as "2" in Section F on Part 2 of the Product Worksheet in Supplement Three-A to PR Notice 93-7.

The eyewear requirement for early entry in the Agricultural Use Requirements Box on your proposed labeling is missing or incorrect. Remove the incorrect requirement (if any) and add "Protective eyewear". The exact location is shown in Sections F and G on Part 2 of the Product Worksheet in Supplement Three-A to PR Notice 93-7.

Delete the Nonagricultural Use Requirements box on your proposed label or add the appropriate restriction per PR 93-1.

Correct the typographical errors circled on your proposed label.



50 lb. Bag Treats up to 5,500 Sq.Ft. FOR COMMERCIAL USE ON PARKS, GOLF COURSES, ATH-LETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURS-ERY, TURF AND LANDSCAPE PERSONNEL ONLY A PRE-EMERGENT WEED CONTROL

PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS ST. AUGUSTINEGRASS, BERMUDA GRASS, TALL

FESCUE, ZOYSIAGRASS
Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed & Other Weeds Listed

ACTIVE INGREDIENT:

Osadiazon; [2-lert-butyl-4-(2,4-dichloro-5-leoproposyphenyl) A'-1,3,4-osadiazolin-5-one] -INERT INGREDIENTS:-

0.50% 99.50% Total

s RONSTAR® Herbicide a Reg. Trade Mark of Bhose Poul KEEP OUT OF REACH OF CHILDREN

WARNING = AVISO

STATE AT LANGUE AT THE PARTY OF A LEAST AND A CONTROL OF THE PARTY OF A LEAST AND A CONTROL OF THE PARTY STATEMENTS OF PRACTICAL TREATMENT

F SWALLOWED. Cult a physician or Poison Control Center immediately, induce wornling by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce wornling or give anything by mouth to an unconscious or convutsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash with

plenty of scep and water. Get medical affention if irritation persists.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Do not allow domestic animals to graze treated areas, Avoid contamination of feed and foodstuffs.

allow domestic animals to graze treated areas, Avoid consmination of read and constants.

Apply: PERSONAL PROTECTIVE EQUIPMENT (PPE)

Apply: The latter handlers must wear overself over short-sleened aller and short parists, waterprant gloves; chemical relations toolware plus socies; protective systemas; chemical features gloves for overself desposure; chemical relations approximation when cleaning equipment, Politic manufacturer's instructions for cleaning free internal processor. It is such instructions for flationalities, use detergent and hot water. Keep and weeth PPE separately from other

Series Recommendations

Light Michiel their hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if posticide gats tracks. Then weak thoroughly and put on clothing. Remove PPE Immediately after handling this product. Wash the outside of stand bidger immediate. At home as postable, with thoroughly and change into clean clothing.

DIFFIC TIONS FOR USE

It is the product in a way that will contact workers or other partons, where directly of this task that, only producted handling may be in the area during application. For any triple makes the product to four State or Titbe, combut the agency trippenalible for publicate highestical. Od not apply this product through any type of trippenalible for publicate highestical.

AGRICULTURAL USE REQUIREMENTS (SELECTION STATES)

BE RELIFICATION IN PROCESSION WITH Its labeling and with the Worker From the Selection States of Corp. part 170. That states is sometime with its labeling and with the Worker From the Selection of Selections of Corp. part 170. That states is sometime in contract for the selection of Selections is selected as the selection of Selections and Selections of Selections and Selections of Selections and Selections of Selections and Selections of Selections of

ENVIRONMENTAL HAZARDS

This posticide is toxic to fish. For terrestrial uses, do not apply directly to water, or to area: where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions fevor drift from areas treated. Do not apply where runoff is likely to occur. Do not conteminate water when disposing of equipment washnuters. Apply only as seculiarion this labor

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal <u>STORAGE</u>. Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of sull, sweep up, and use according to directions or dispose of as follows. <u>PRETINEM DISPORAL:</u> Positions westers are twice improper disposal of ascesses positive, or related to a whether of Federal tow IT it does reactes security to dispose of to use describes.

WEED CONTROL AROUND WOODY ORNAMENTAL SHRUBS, VINES & TREES Controls Grassy Weeds Controls Broadlest Weeds

Controls Grassy Weeds

Annual Bluegrass *Barnyardgrass Bristly Fordall Carpetgrass Crabgrass Fall Panicum

Goosegrass Green Fortal

Jungle Ricegrass Panama Paspalum

Carpetweed Groundsel, common Common Pursiane Petty Spurge
*Evening Primrose *Prostrate Spurge *Evening Primrose Florida Pueley Galinsoga Golden Ragwort Lambsquarter

Pennsylvania Smartweed Redroot Pigw Speedwell Spotted Catsear
*Swinecress Yellow Woodsorrel

Can be applied to both newly transplanted and established ornemental plants wher both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control. Do not disturb or cultivate soil after treatment. Do not apply t wet foliage or under conditions in which granules will collect on leaves.

HOW MUCH TO USE *For Hard-to-Kill Weeds

For Hard-To-Kill weeds in the above chart (as indicated by an *asterisk) use the rat of 50 lbs. to cover 2,750 Sq.Ft.to 3,666 sq.ft. (= 3 to 4 lbs. A.I./Acre).

For Susceptible Weeds

For weeds listed in the above chart (without asterisk) apply 50 lbs. to treat 3,666 t 5,500 Sq.Ft. (= 2 to 3 lbs. A.I. /acre)of ornamental planted area. This also supplie plants with a balanced plant food.

WHEN TO APPLY

See annual Chart in the basic label for proper time to apply in your area. For best wee control apply at least, two weeks before weed seeds are expected to germinate. Fo general overall pre-emergence weed control of the most common weeds found i lawns the rate of 50 lbs./3,666 Sq.Ft, is preferred, equal to 3 lbs. A.i. per scre.)

FOR AREAS WHERE WEEDS ARE GROWING

If treatment is desired in non-turf areas where weeds are presently growing, firs remove existing weeds. Rake and amouth the soil by "breaking-up" clumps. Apply a rate as directed for specific weeds. (50 lbs. to treat either 5,500 Sq. Ft. or 3,666 Sq. Ft.) This treatment will only control weeds germinating from seeds. It will not control weeds sprouting from roots! Irrigate the area after treatment with 1/4 to 1/2 inch water to get maximum weed control.

FOR FIELD GROWN ORNAMENTALS

*Apply at the rate of 50 lbs. per 5,500 or 3,666 Sq.Ft. of area.

Carpobrulus

Algerian My

Vince Major English My

Sedum

Osleospermum

Established Plants. This product may be used around the following Euonymus Pittosporm GROUND COVERS

Arborvitae Euonymus Forsythia **Privet** Ash Holly Rhododenron Azalea Barberry Honeysuckle Rose Russian Olive Juniper Japanese Birch Lifac Spruce Maple Camellia Taxus Oak Periwinide Cotoneaster Vibumum Xylosma Crabapple

Pachesandra Ajuga Euonymus

Dogwood Pine Eutrymus

*Use higher rate (50 lbs:/3,666 sq.ft.) for hard to till weeds.

Newly transplanted ornamentals

May be used around the following plants; Camelia, Dogwood. Forsythia, Hedera Holly, Juniper, Oak, Periwinide, Pine, Privet, Rhododendron, Russian Olive, Spruce Taxus, Tsugs.

ADDITIONAL APPLICATIONS

To obtain continued season long weed control on the following list of plants additions applications may be made 60 to 120 days after the previous application in the followin situations:

2 REPEAT TREATMENTS 60 lbs / 3,666 SQ.FT ON FOLLOWING PLANTS

Established - Container Grown -- Abelia, Common Bonwood, English My, Holly Juniper, Pine, Rhododendron

Newly Transplanted - Container Grown Common Boxwood, English Ivy, Juniper Pine, Rhododendron

3 REPEAT TREATMENTS 60 lbs / 3,000 SQ.FT. ON FOLLOWING PLANTS

Established - Container Grown - Abelia, Holly, Rhododendron. Forest Hurseries - For Pines - (Loblolly, Slash, Eastern White.)

WEED AND FEED FOR CONTAINER GROWN ORNAMENTAL PLANTS

This product is recommended on a wide variety of container and field grown woody ernaments shrube, vines and frees. It may be used on the following list of container grown plants. Use a the rate of 50 fbs. per 2,750 sq.ft. of pot soft surface area. Plants in 14 inch containers set or in 12 fbst wide rows X 100 feet long will require 10 3r4 fbs. product per row (Equivalent to the A1 per zero). Apply with a "abundar strain habit" gross-conder or granute applicator the

Oxadlazon; [2-tert-butyl-4-(2,4-dichloro-5-laoproponyphenyl) A?-1,3,4-oxadlazolin-5-one] -INERT INGREDIENTS: 0.50%

ine RONSTARO Harbicide a Reg. Trade Mark of Rhor KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

SI usted no inflicte to allowed, buildue a algular pairs que to la continue a ustad en detalle.

(If you do hit understand the label, find correcore to deplace it to you in detail. STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center Immediately. Induce vomitir by giving victim 1 or 2 glasses of water and louching back of threat with finger. Do not induc vomiting or give anything by mouth to an unconectous or convalsing person. FIN EYES: Flush eyes with plenty of weter. Get medical attention if irritation persists. IF ON SKIN; Wash with

my of scap and water. Get medical attention of initiation parasists.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING: Causes eye and skin Irritation, Avoid contact with skin, eyes or clothing. Do not allow domestic seitnels to graze treated areas. Avoid contamination of feed and foodskulfs.

Link of PERSONAL PROTECTIVE EQUIPMENT (PPE)
Applications and Salah Interdes in the water coverable over their 1990 Salah Salah Interdes in the water prior of govers; chemical resistant footoners in the salah Interdes in the salah Interdes

User Safety Recommendations

User Safety Recommendations

User Safety Recommendations

User should wish hands before eating, drinking, chewing gum, using tobacce or using the toilet. Remove clothing immediately it pesticide gets finales. This wish thoroughly and put or closin clothing. Remove PPE immediately after handing this product. Wash the outside of aloves before rimoring. As soon as possible with increasing drinking into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its tabeling. Do not apply this product in a way that will contact workers or other parsons, either directly of through drift. Only protected handlers may be in the area during application. For any requirements apacific to your State or Tribe, consult the against helponsible for pasticide regulation. Do not apply this product through any type of irrigations system.

AGRICULTURAL USE REQUIREMENTS

AGRICUL TURAL USE RECUIRER IN 15

Liè fire productionly in accordance with its labeling and with the Worter Protection Standard, 40 CFR part 170, This Standard contains requirements for the protection of agricultural posticides. It contains requirements for training, decontamination, licellication and emergency assistance. It also contains specific instructions and stacephone pertening to the slatements on this label about personal protective equipment (PPE) notification to workers, and restricted entity intervals. The requirements in this box only apply to seem of this product (nursery grains emanmental should, usee, base and contains grave speciments by that are covered by the Worker Protection Standard.

Do not enter of allow worker entry into treated areas during the Restricted entry interval (REI) of 12 hours.

Of 12 recurs.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that Involves contact with anything that has been treated, such as plants, soil, or water is coveralts over short sleeved shirt and short paritic waterproof gloves, chemical resistant footweer plus socks and chemical resistant headgier for overhead exposure.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. For terrestries uses, do not apply directly to water, or to areas where surface weter is present or to intertidal areas below the mean high water mark. Do not apply when weether conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment weetwaters. Apply only as specified on this label

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or dispos Do not contaminate water, food, or feed by storage or disposar.

<u>STORAGE</u>: Store product in its original beg, in a cool, dry, locked place out of reach of children, in case of spill, sweep up, and use according to direct... is or dispose of as follows.

<u>PESTICIDE DISPOSAL</u>: Peaticide westes are toxic. Improper disposal of evcess pesticide, or rineate is a violation of Federal law. If these weetes cannot be disposal of by use according to label instructions, contact your State Penticide or Environmental Control Agency, or the Hazardous Weste representative at the nearest EPA regional effice for guidence.

CONTRABLE PERECRAL - Commissals amonts have into configurate agreement. Then dispose

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sentiary landfill or by incineration, or, if allowed by State and local authorities,

Sy burning Dansed stay cut of stroke

BEST A CASE TEED TERMS FOR STROKE COMMENTS

Total Nikrogen (N). variable%

Nikrogen (N). variable%

TEED 2 3 190.1

Ammonieral Nitrogen

Water Soluble Nitrogen
and/or Urea Nitrogen
Water Insoluble Nitrogen
Available Phosphoric Acid (P,O.)
Soluble Polash (K,O).

Derived from Ammonium Sulfate, Useafore, Mariete of St.

Polash; Menganese as Mn. 5%; Iron as Fe 1 0%; Chlorine <15%

Delived the Fed ral Inserticide, common Redemicide Act Calistemon warrists and trader trader.

FPA Reg. No. Cearnothus
C edar
Cotonesster

řŒ

Barberry

- Cotoneaster

Agapenthus

Andromeda

Arctostaphylos

Arbondae

Azaba

Dogwood Douglas Fir Eucalyptus Euonymus-Forsyttee Gardenia

Girmo Heath Heather **Pyracanthe** Quercus Raphiolepus Rhododendran Rose Star Jasmine Syzygium Viburnum

EPA Reg. No. 35512-31

EPA Est. No. 35512-FL-1 **NET WEIGHT 50 POUNDS**

Can be applied to both newly transplented and established ornamental plants wher both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control. Do not disturb or cultivate soil after treatment. Do not apply t wet foliage or under conditions in which granules will collect on leeves. **HOW MUCH TO USE** *For Hard-to-Kill Weeds For Hard-To-Kill weeds in the above chart (as indicated by an *asterisk') use the rat

of 50 lbs. to cover 2,750 Sq.Ft.to 3,666 sq.ft. (= 3 to 4 lbs. A.I./Acre).

*Liverwort

Yellow Woodsorrel

*Panama Paspelum

Arborvitae

Ash

Azalea

Birch

Barberry

Camella

For Susceptible Weeds

For weeds listed in the above chart (without asterisk) apply 50 lbs. to treat 3,686 t 5,500 Sq.Ft. (= 2 to 3 lbs. A.I. /acre)of ornamental planted area. This also supplie plants with a belanced plant food.

WHEN TO APPLY

See annual Chart in the basic label for proper time to apply in your area. For best were control apply at least, two weeks before weed seeds are expected to germinate. For general overall pre-emergence weed control of the most common weeds found it lewns the rate of 50 lbs./3,606 Sq.Ft. is preferred, equal to 3 lbs. A.I. per acre.) FOR AREAS WHERE WEEDS ARE GROWING

If treatment is desired in non-turf areas where weeds are presently growing, fire remove existing weeds. Rake and smooth the soil by "breaking-up" clumps. Apply a rate as directed for specific weeds. (50 lbs. to treat either 5,500 Sq. Ft. or 3,666 Sq. Ft.) This treatment will only control weeds germinating from seeds. It will not control weeds sprouting from roots! Irrigate the area after treatment with 1/4 to 1/2 inch water to get maximum weed control.

FOR FIELD GROWN ORNAMENTALS

*Apply at the rate of 50 lbs. per 5,500 or 3,666 Sq.Ft. of area. **Established Plants**

This product may be used around the following Pittosporm Euonymus **GROUND COVERS** Carpobrutus Forsythia Privet Rhododenron Osteospermum Holly Sedum Honeysucide Rose Japanese Juniper Russian Olive Algerian Ivy Litac Vinca Major Spruce Maple English Ivy Taxus Cotoneaster Oak Pachasandra Viburnum Aluga Periwinkle Crabapple Xylosma: Pine **Euonymus**

Dogwood *Use higher rate (50 lbs./3,666 sq.ft.) for hard to kill weeds.

Newly transplanted ornamentals

May be used around the following plants; Carnellia, Dogwood, Forsythia, Hedera
Holly, Juniper, Oak, Periwinkle, Pine, Privet, Rhododendron, Russian Olive, Spruce Taxus, Tsuga.

ADDITIONAL APPLICATIONS

To obtain continued season long weed control on the following list of plants additions applications may be made 60 to 120 days after the previous application in the followin

2 REPEAT TREATMENTS 60 Nbs / 3.666 SQLFT ON FOLLOWING PLANTS

<u>Established - Container Grown</u> – Ahelia, Common Boxwood, English Ivy, Holly Juniper, Pine, Rhododendron

Newly Transplanted - Container Grown Common Boxwood, English Ivy, Juniper Pine, Rhododendron.

3 REPEAT TREATMENTS \$0 lbs / 3,666 SQ.FT. ON FOLLOWING PLANTS

Established - Container Grown - Abelia, Holly, Rhododendron. Forest Nurseries - For Pines -(Lobiolly, Slash, Eastern White.)

WEED AND FEED FOR CONTAINER GROWN ORNAMENTAL PLANTS

This product is recommended on a wide variety of container and field grown woody ornaments shrubs, vines and trees. It may be used on the following list of container grown plants. Use a the rate of 50 lbs. per 2,750 sq.R. of pot soil surface area, Plants in 14 inch containers set set in 12 foot wide rows X 100 feet long will require 10 3/4 lbs. product per row (Equivalent to lbs. A.I. per acre). Apply with a "shoulder strap held" grass-seeder or granule applicator the will direct the granules to each pot soil surface.

May be used on the following Container Grown Established Ornamental Plant Species

Cypress Honeysucide **Podocarpus** Jasmine Privet Juniper-Var. Justica Leucothoe evergreen Magnolia Mahonie Myrtle **Osmenthus** Periwinkle Photina Xylosma Hedera Pine-varieties

Holly **Pittosporum** NOTICE: fluyer and user assume all risk and liability of use, storage and/orhandling of this product not in accordance with the terms of this label (35-31SP PM4 / WPS Certification 10-25-93)

HOWARD FERTILIZER COMPANY

3 1994

P.O. BOX 593800 - ORLANDO, FL 32859-3800