

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460		EPA REGISTRATION NO. 25512-24	DATE OF ISSUANCE JUN 16 1994
NOTICE OF PESTICIDE: ☒ REGISTRATION ☐ REREГИSTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)		TERM OF ISSUANCE	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		NAME OF PESTICIDE PRODUCT	
Howard Fertilizer Co. P.O. Box 597890 Orlando, FL 32809		FERTILIZER, 10-10-10, 10-10-10, 10-10-10	
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.			
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.			
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.			
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.			
This product is conditionally registered in accordance with FIFRA Section 3(c)(7)(A) provided that you:			
1. Submit a copy of your product label, FIFRA Section 3(c)(7)(A) or 3(c)(7)(B), to the Agency prior to release of the product for shipment.			
2. Make the labeling changes listed below, before you release the product for shipment:			
a. Add the phrase "EPA Registration No. 25512-24"			
b. Submit five (5) copies of your final printed label before you release the product for shipment.			
If these conditions are not complied with, this registration will be subject to cancellation in accordance with FIFRA Sec. 6(a). Your release for shipment of the product constitutes acceptance of these conditions.			
3. Submit a copy of the final label to the Agency prior to release of the product for shipment.			
BEST AVAILABLE COPY			
☐ ATTACHMENT IS APPLICABLE		DATE	
SIGNATURE OF APPROVING OFFICIAL [Signature]		[Signature]	



For St. Augustine & Centipede Lawns

WEED CONTROL WITH PREMIUM LAWN FOOD

40 POUND BAG COVERS

5,000 Sq.Ft. of Lawn

Controls Florida Pusley, Lippia, Pigweed and certain Broadleaf Weeds as listed. Prevents Crabgrass, Paspalums, and Spotted Spurge and other weeds as listed.

FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

ACTIVE INGREDIENT:

Atrazine 2-chloro-4-ethyl-6-methyl-2-isopropylamino-s-triazine . . . 0.35%

Related Compounds . . . 0.02%

INERT INGREDIENTS . . . 99.43%
Total 100.00%

Contains 11.4 lbs. of Atrazine per lb. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation or contact with eyes. Wash after handling or using. Keep away from feed or food products.

ENVIRONMENTAL HAZARDS

Atrazine can travel (leach) through soil and enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass	Henbit
(Poa Annua)	Knotweed
Annual Sowthistle	Lambsquarters
Clover	Lespedeza
Black Medic	Lippia (Matchweed)
Blue Toadflax	Moneywort
Bull Paspalum	Morning Glory
Buckhorn Plantain	Narrowleaf Vetch
Bur Clover	Parsley-Piert
Carolina Geranium	Oxalis (Common and Yellow Woodsorrel)
(Cranesbill)	Pigweed
Chickweed (Common and Mouse Ear)	Purslane
Cocklebur	*Ragweed
Common Dandelion	Red Sorrel
Common Purslane	Ricegrass
Crabgrass	Sandspur
Crowfoot Grass	Sedges*
Curtweed	Shepherd's Purse
Cutleaf Evening Primrose	Sicklepod
Dichondra	Spotted Spurge
Dollar Weed	Spurweed
Florida Beggarweed	Slinging Nettle
Florida Beltony	Swinecrest
(Rattlesnake Weed)	Venus Looking Glass
Florida Pusley	Virginia Creeper
Fragrant Cudweed	Wild Mustard
(Rabbit Tobacco)	Wood Sorrel

*Repeat application are required for control of Nutsedge grasses.

EPA REG. NO. 35512-

NET WEIGHT 40 POUNDS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner inconsistent with its labeling. Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for ornamental purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a Pre-Emergent and also as a Post Emergent weed killer.

Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1st up to March 1st for best control otherwise product may be applied up to June 15th with good weed control. A fall application will be needed for control of certain annuals such as Chickweed, Henbit, Rabbit Tobacco, Florida Beltony and Evening Primrose.

In case of heavy weed infestation or low soil moisture, a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle wind drift.

HOW TO APPLY

This product may be applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 40 pounds to treat 5,000 Sq. Ft. - 20 lbs. bag treats 2,500 Sq. Ft. of lawn (equivalent to 8 lbs. per 1,000 Sq. Ft.). Do not exceed recommended rates. Sprinkle the area immediately after application to wash granules off the grass blades.

SPREADER SETTINGS 40 lbs./ 5,000 Sq.Ft.

CYCLONE 45 SCOTT 7

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHERE TO USE

This material is for use on St. Augustine and Centipede lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant "sprigs" within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. **DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA, RYE, PERMUDA, OR DICHONDRA LAWNS.**

USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended, on this basis or injury to those other grasses may result.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	12%
Nitrate Nitrogen	0.00%
Ammoniacal Nitrogen	11.20%
Water Soluble Nitrogen	4.60%
(and or Urea Nitrogen)	
Water Insoluble Nitrogen	0.00%
Available Phosphoric Acid (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%
Chlorine not more than	5.65%

Derived from Sulfate of Ammonia, Ammonium Phosphate, Muriate of Potash, Sulphate Potash Magnesium. Potash Acidity Equivalent to 1245 lbs. of Calcium Carbonate per ton.

Statement of Secondary Plant Foods

Total Magnesium as Mg - 1.20%, Iron as Fe - 1.00%, Water Soluble Magnesium Mg - 1.20%, Zinc as Zn - 0.05%, Manganese as Mn - 0.50%, Boron as B - 0.02%, Copper as Cu - 0.02%, Molybdenum as Mo - 0.002% (Derived from Manganese Sulphate, Iron Oxide, Manganese Oxide, Copper Oxide, Zinc Oxide, Borax, Sodium Molybdate).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Wrap in several layers of newspaper and discard in trash.

NOTICE: Buyer and user assumes all risk of use, storage and/or handling of this material not in accordance with the terms of this label (35-A1767 PM4/06 31-92)

EPA EST. NO. 35512-FL-1

HOWARD FERTILIZER COMPANY

P.O. BOX 13800 - ORLANDO, FL 32859

ACCEPTED

JUL 1 1984

35-A1767 PM4/06 31-92