

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

35512-35

DATE OF ISSUANCE

6/1/95

NOTICE OF PESTICIDE:

REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

TERM OF ISSUANCE

Conditional

NAME OF PESTICIDE PRODUCT

Purr Pride[®] with .67% RUSSIAN[®]

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

HOWARD FERTILIZER COMPANY
8306 S. Grange Avenue
Post Office Box 593800
Orlando, FL 32859-3800

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(2) provided that you:

1. add the phrase "EPA registration no. 35512-35 to the label before you release the product for shipment.
2. submit one (1) copy of the final printed label to EPA before you release the product for shipment.
3. submit and/or cite all data required for registration/reregistration of this product under FIFRA sec. 3(c)(2) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA sec 4.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

RECEIVED

JOANNE I. MILLER
Product Manager (23)
Pesticide-Berkeley Branch
Registration Division (75050)

ATTACHMENT IS APPLICABLE

BEST COPY AVAILABLE

SIGNATURE OF APPROVING OFFICIAL

Joanna I. Miller

DATE

6/1/95



TURF PRIDE® with .67% RONSTAR® FOR COMMERCIAL LAWNS & ORNAMENTAL TURF

50 lb. Bag Treats up to 7,300 Sq.Ft.

FOR USE ON PARKS, GOLF COURSES, ATHLETIC FIELDS,
AND OTHER RECREATIONAL / ORNAMENTAL TURFGRASS

FOR SALE TO, USE AND STORAGE BY COMMERCIAL TURF
AND LANDSCAPE PERSONNEL ONLY

Not for use on turf being grown for sale or other commercial
use as sod, or for commercial seed production, or for research
purposes. Not for use on Home Lawns.

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida
Pusley, Oxalis, Stinging Nettle, Carpetgrass,
Pigweed & Other Weeds Listed

ACTIVE INGREDIENT:

Oxadiazon: [2-tert-butyl-4-(2,4-dichloro-5-
isopropoxyphenyl) N-1,3,4-oxadiazolin-5-one] - . . . 0.67%
99.33%

INERT INGREDIENTS: Total 100.00%

Contains RONSTAR® Herbicide a Reg. Trade Mark of Rhone Poulenc Inc.

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle
(If you do not understand the label, find someone to explain it to you in detail.)

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting
by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce
vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:**
Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash
with plenty of soap and water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Wash
hands before eating, drinking, chewing gum, using tobacco or using the toilet. Do not allow
domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling
Do not apply this product through any type of irrigation system

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water
is present or to intertidal areas below the mean high water mark. Do not apply when weather
conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not
contaminate water when disposing of equipment washwaters. Apply only as specified on this
label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

HAZARD: Store product in its original bag, in a cool, dry, locked place out of reach of
children. In case of spill, sweep up, and use according to directions or dispose of as follows.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide,
or residue is a violation of Federal law. If these wastes cannot be disposed of by use according
to label instructions, contact your State Pesticide or Environmental Control Agency, or the
Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose
of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local
authorities, by burning. If burned stay out of smoke.

ACCEPTED

GUARANTEED FERTILIZER ANALYSIS with COMMENTS in EPA Letter Dated

Total Nitrogen (N)	15%	
Nitrate Nitrogen	0%	
Ammoniacal Nitrogen	12%	
Water Soluble Nitrogen		
and/or Urea Nitrogen	0%	
Water Insoluble Nitrogen	3%	
Available Phosphoric Acid (P ₂ O ₅)	15%	
Soluble Potash (K ₂ O)	15%	
Derived from Ammonium Sulfate, Ureaform		
Potash, Manganese as Mn 5%, Iron as Fe 1.0%, Chlorine 1.5%		

JUN 1 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
registered for the pesticide
EPA Reg. No. 35512-35

CONTROLS WEED GRASSES & BROADLEAF WEEDS

Weed Grasses

- *Annual Bluegrass
- *Barnyardgrass
- Bristly Foxtail
- Carpetgrass
- Crabgrass
- Fall Panicum
- Field Sandbur
- **Goosegrass
- Green Foxtail
- **Jungle Ricegrass
- **Panama Paspalum

Broadleaf Weeds

- Bittercress
- Carpetweed
- *Groundsel, common
- Common Furslane
- *Evening Primrose
- Florida Pusley
- *Galinsoga
- Golden Ragwort
- Lambsquarter
- **Liverwort
- Oxalis
- Pennsylvania Smartweed
- Peltly Spurge
- **Prostrate Spurge
- Redroot Pigweed
- Speedwell
- Spotted Catsear
- Stinging Nettle
- **Swinecress
- Yellow Woodsorrel

50 lbs. treats 7,300 sq. ft. For weeds with asterisks see paragraphs below.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that
granules will reach the soil surface. Measure the area to be treated and uniformly apply the
required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to wet
turf. Do not over-lap spreader applications, on St. Augustinegrass since discoloration
"streaking" may temporarily occur from "double rate". If rain is not expected within 24 hours
sprinkle treated turf with 1/4 to 1/2 inch to increase effectiveness.

HOW MUCH TO USE

FOR CRABGRASS and other WEEDS LISTED ABOVE

For susceptible weeds listed above (without *asterisk) apply 50 lbs. per 7,300 Sq.Ft. when
turf has been in a yearly maintenance program (equal to 2 lbs. A.I. herbicide per acre.) Apply
50 lbs. to treat 4,900 Sq Ft. of lawn if it has been heavily infested with Crabgrass and other
susceptible weeds (equal to 3 lb. A.I. herbicide/acre) Apply in early spring 1 to 2 weeks prior
to weed seed germination. This product is a pre-emergence herbicide. It will not control
growing weeds; only germinating weed seeds. Late applications will control only those weed
seeds that germinate after the application. See "When To Apply" Chart.

* MORE TOLERANT WEEDS

BARNYARDGRASS, EVENING PRIMROSE, GALINSOGA, GROUNDSEL, and SWINECRESS

Where these weeds are a problem apply 50 lbs. to treat 3,650 to 4,900 Sq.Ft. of Lawn; (equal
to 3 to 4 lbs. Active Herbicide per acre) in early spring or 1 to 2 weeks before expected weed
seed germination. Will not control of emerged and growing weeds; only effective against
germinating weed seeds. Observe 50#/4,900 sq.ft. maximum rate for St. Augustine grass.

** MOST TOLERANT WEEDS

GOOSEGRASS, JUNGLERICE, LIVERWORT, PANAMA PASPALUM, PROSTRATE SPURGE,

For Hard-to-control or "late germinating" weeds apply 50 lbs. to treat 3,650 Sq.Ft. (4 lbs. A.I./
acre) and apply 1 to 2 weeks prior to seed germination. See Chart for When To Apply. Observe
50#/4,900 sq.ft. maximum rate for St. Augustine grass.

FOR ANNUAL BLUEGRASS (Poa Annua)

Apply 50 lbs. per 3,650 to 4,800 Sq Ft. of Lawn, equal to 3 to 4 lbs. active herbicide per acre
Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed
germination. Will not control annual bluegrass present at time of application; therefore only
germinating seeds. Refer to chart for proper timing in your locale. Observe 50#/4,900 sq.ft.
maximum rate for St. Augustine grass.

SPREADER SETTINGS FOR SQUARE FOOT COVERAGE

	AMT. OF for	7,300 Sq.Ft.	4,800 Sq.Ft.	3,650 Sq.Ft.
SPREADER PRODUCT	2#A./Acre	3#A./Acre	4#A./Acre	
CYCLONE B-1	50 LBS.	6 1/2	7	8
SCOTTS #35	50 LBS.	4 1/2	5	6

NOTE: Spreader rate varies with walking speed

WHEN TO APPLY

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

	JAN	FEB	MAR	APR	MAY	JUNE	JULY	AUG	SEPT	OCT	NOV	DEC
1								3				
2												
3a												
4 >				2a								
1												
2												
2a												
3												
3a												
4												

1 For CRABGRASS and other weeds germinating in the Spring in Northern turf
2 For GOOSEGRASS and other Late Germinating weeds such as PROSTRATE SPURGE
2a For GOOSEGRASS in Southern Turf
3 For POA ANNUA in Northern Turf
3a For POA ANNUA in Southern Turf
4 For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf

PRECAUTIONS IN USING:

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Carpetweed
grass since damage will occur. Do not use on Putting greens or "Tees" due to varying
cultural practices used. Use only the rate of 50 lbs. to treat 4,900 Sq.Ft. on St.
Augustinegrass since temporary discoloration will occur from heavier rates. Avoid
contact with flowers and shrubs except as recommended on the Ornamental Use
Supplemental Labeling available for this product. For newly seeded lawns do not
apply this product until after the second or third cutting. For best lawn care, place new
grass seed in the fall of the year. Spring pre-emergence weed control can then be
applied.

RESEEDING TREATED AREAS Delay new seeding until 4 months after treatment
NOTICE Buyer and user assume all risk and liability of use, storage and/or handling of this
product not in accordance with the terms of this label (35512-35 PMS / 02-17-95)

EPA Reg. No. 35512- EPA EST. NO. 35512-FL-1

NET WEIGHT 50 POUNDS

HOWARD FERTILIZER COMPANY

P.O. BOX 593800 - ORLANDO, FL 32859-3800

BEST COPY AVAILABLE