

35512-43

7/9/2003

INSERT RONSTAR LOGO HERE

Turf Pride® Fertilizer With Ronstar®

50 lb. Bag Treats up to X,XXX Sq. Ft.

FOR SALE TO, USE AND STORAGE BY TURF MANAGEMENT AND LANDSCAPE PERSONNEL ONLY. NOT FOR USE ON HOME LAWNS. FOR USE ON GOLF COURSES, PARKS, SOD FARMS, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS.

A Pre-emergent Weed Control Product plus a Quality Fertilizer for use on varieties of Perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue and Zoysiagrass.

ACTIVE INGREDIENT: Oxadiazon, [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) A-1,3,4-oxadiazolin-5-one] 0.50% 2.00%
OTHER INGREDIENTS: 99.50% 98.00%
TOTAL 100.00% 100.00%

Turf Pride® is a Reg. Trademark of Howard Fertilizer & Chemical Co., Inc

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

FIRST AID

- Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact a XXX-XXX-XXXX for emergency medical treatment information.
- If swallowed:**
 - Call poison control center or doctor immediately for treatment advice.
 - Have person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- If in eyes:**
 - Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 - Call a poison control center or doctor for treatment advice.
- If on skin or clothing:**
 - Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
- If inhaled:**
 - Move person to fresh air.
 - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
 - Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Wear protective eyewear (goggles, face shield or safety glasses). Causes skin irritation. Do not get in eyes, on skin or clothing. Harmful if swallowed, absorbed through the skin, or inhaled. Avoid breathing dust, contact with skin, eyes or clothing. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead application; chemical resistant apron when cleaning equipment. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system. Apply only as specified on this label.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forest nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; waterproof gloves; chemical resistant footwear plus socks; protective eyewear, and chemical resistant headgear for overhead exposure.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of a spill, sweep up, and use according to directions or dispose of as follows. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	variable %
Nitrate Nitrogen	— %
Ammoniacal Nitrogen	— %
Water Soluble Nitrogen and/or Urea Nitrogen	— %
Water Insoluble Nitrogen	— %
Available Phosphoric Acid (P ₂ O ₅)	variable %
Soluble Potash (K ₂ O)	variable %
Derived from Ammonium Sulfate, Ureaform, Muriate of Potash; Manganese as Mn % from as F ₂ % Chlorine %	

ACCEPTED
JUL 9 2003
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 35512-43

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants only on farms, forests, nurseries, or greenhouses. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Wear protective clothing including long-sleeved shirt and impervious gloves when handling. Launder exposed clothing before reuse.

Controls Crabgrass, Goosegrass, Poa Annua, Annual Sedge and other weeds as listed, on Golf Courses, Commercial Lawns, and other Athletic and/or Recreational Ornamental Turf.

WHERE, HOW MUCH AND WHEN TO USE ON TURFGRASS

This product may be used at full rate or as a "split application in established perennial Bluegrass, Bentgrass (see restrictions), Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysiagrass turf, at apartment/condominium complexes, corporate and public parks, golf courses, sod farms, athletic fields, playgrounds, cemeteries and other ornamental and recreational turfgrass area. Not for use on home lawns. The turf should be free of leaves and thatch and in condition to allow granules to reach the soil surface. Know the square footage of the area to be treated and uniformly apply using a granule spreader. Do not apply to turf when it is wet. When using at full rate do not overlap spreader applications, especially on St. Augustinegrass since discoloration "streaking" may temporarily occur from the "doubled rate". Observe the maximum rate for St. Augustinegrass and Bentgrass of 3 lbs. a.i./acre which is equiv. to XXX lbs. of this product per 1,000 sq. ft. See below for restrictions when using on Bentgrass. After applications if rain is not expected within 24 hours sprinkle irrigate the treated turf with up to a 1/4 inch of water to get maximum weed control and quicker response from the fertilizer nutrients. This product is a pre-emergence herbicide and must be applied prior to weed seed germination. It will not control growing weeds. *See paragraph below for instructions to apply the 3 lb. or 4 lb. Rate as a split application. For the rate of 2 - 4 lbs. a.i./acre, on weedy turf, containing difficult-to-control weeds, apply this product in one application at the 3 or 4 lb. rate.

Weeds Controlled	Pre-emergence Weed Control		Directions -
	lbs. a.i./Acre	lbs. of Product per 1,000 sq. ft.	
Crabgrass (<i>Digitaria</i> , spp.)	3 to 4 lbs. a.i./A	XXX to XXX lbs.	Apply in late winter or early spring prior to weed seed germination. Application made after April or May may not be fully effective.
Goosegrass (<i>Elymus indica</i>)	2 to 4 lbs. a.i./A	XXX to XXX lbs.	
Field Sandbur (<i>Cenchrus incertus</i>)	XXX to XXX lbs. product/acre	XXX to XXX lbs.	Apply in late summer to early fall prior to weed seed germination.
Annual Sedge (<i>Cyperus compressus</i>)	2 to 4 lbs. a.i./A	XXX to XXX lbs.	
Annual Bluegrass (<i>Poa annua</i> spp.) (See Note Below)	2 to 4 lbs. a.i./A	XXX to XXX lbs.	Apply in early spring prior to weed seed germination.
Virginia Buttonweed (<i>Diodia virginiana</i>)	4 lbs. a.i./A	XXX lbs./A	
Pigweed (<i>Amaranthus</i> spp.)	4 lbs. a.i./A	XXX lbs./A	For use in Bluegrass; Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup with Perennial Ryegrass only.
Florida Pusley (<i>Richardia scabra</i>)	4 lbs. a.i./A	XXX lbs./A	
Oxalis (<i>Oxalis</i> spp.)	4 lbs. a.i./A	XXX lbs./A	
Stinging Nettle (<i>Urtica dioica</i>)	4 lbs. a.i./A	XXX lbs./A	
Carpentergrass (<i>Axonopus officinis</i>)	4 lbs. a.i./A	XXX lbs./A	
Crabgrass, Goosegrass and other annual weeds as listed.	3 to 4 lbs. a.i./A	XXX to XXX lbs./A	Season Turf
	XXX to XXX lbs./A	1,000 sq. ft.	For use in Bluegrass; Bentgrass, perennial Ryegrass and Tall Fescue. Make application in the fall after turf has become dormant. For late fall applications, seeding into treated areas should be delayed until 4 months after greenup with Perennial Ryegrass only.
	2 to 4 lbs. a.i./A	XXX to XXX lbs./A	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa annua spp. seed germination. Any Poa annua spp. established when this product is applied will not be controlled. It should not be applied closer than 60 days to overseeding. Do not use where Poa trivialis is present in the overseed mixture. Apply the full 2 lb. rate in one application.
	2 to 4 lbs. a.i./A	XXX to XXX lbs. per 1,000 sq. ft.	Apply in late summer to early fall prior to Poa ann